

Supplementary Materials:

Diversity and Distribution of Mesozooplankton in the Coastal Southwestern Mediterranean Alboran Sea, during Summer: What Are the Driving Factors?

Mohamed Reda Benallal ^{1,2,*}, Ahmed Errhif ¹, Laila Somoue ², Mohamed Laabir ³, Hervé Demarcq ⁴, Mohammed Idrissi ², Aziz Agouzouk ², Yassine Goliat ^{1,2}, Hajar Idmoussi ^{1,2}, Ahmed Makaoui ² and Omar Ettahiri ²

¹ Health and Environment Laboratory, Ain Chock Faculty of Sciences, Hassan II University of Casablanca, Morocco, B.P. 5366, Casablanca 20100, Morocco; aerrhif@gmail.com (A.E.); yassine.goliat@gmail.com (Y.G.); hajar.idmoussi@gmail.com (H.I.)

² National Institute of Fisheries Research, Avenue Sidi Abderrahmane, B.P. 20100, Aïn Diab Casablanca, Casablanca 20180, Morocco; lailasomoue@gmail.com (L.S.); idrissi@inrh.ma (M.I.); oceanosaad@gmail.com (A.A.); makaoui@inrh.ma (A.M.); ettahiri@inrh.ma (O.E.)

³ MARBEC, University of Montpellier, CNRS, Ifremer, IRD, 34095 Montpellier, France; mohamed.laabir@umontpellier.fr

⁴ MARBEC, University of Montpellier, CNRS, Ifremer, IRD, 34200 Sète, France; herve.demarcq@ird.fr

* Correspondence: mohamedredabenallal@gmail.com




Figure S1. Chlorophyll-a concentration from the MODIS sensor during the same survey period.




Figure S2. Distribution of bottom chlorophyll-a ($\mu\text{g L}^{-1}$).




Figure S3. Longitudinal comparison along the Moroccan Mediterranean coast of the mesozooplankton and copepod total abundance (ind. m^{-3}) for coastal and offshore stations.




Figure S4. Longitudinal comparison along the Moroccan Mediterranean coast of the species richness (RS) and the Shannon index (H') for coastal and offshore stations.