

Supporting Information

Spectral Properties of Substituted Coumarins in Solution and Polymer Matrices


Jana Donovalová ^{1,*}, Marek Cigán ¹, Henrieta Stankovičová ¹, Jan Gašpar ¹, Martin Danko ², Anton Gáplovský ¹ and Pavol Hrdlovič ^{1,2}

¹ Faculty of Natural Sciences, Institute of Chemistry, Comenius University, Mlynská dolina CH-2, SK-842 15 Bratislava, Slovak; E-Mails: cigan@fns.uniba.sk (M.C.); stankovh@fns.uniba.sk (H.S.); gaspar@fns.uniba.sk (J.G.); gaplovsky@fns.uniba.sk (A.G.)

² Polymer Institute, Slovak Academy of Sciences, 842 36 Bratislava, Dúbravská cesta 9, Slovak; E-Mails: upoldan@savba.sk (M.D.); upolhrdl@savba.sk (P.H.)

* Author to whom correspondence should be addressed; E-Mail: donovalova@fns.uniba.sk; Tel.: +421-2-6029-6306; Fax: +421-2-6029-6337.

Supporting Figure 1. Absorption spectra of carbaldehydes **1** and **5** in chloroform and methanol at $\sim 10^{-4}$ mol dm⁻³.



Supporting Figure 2. The change in chemical shift of signals for various protons of phenylsemicarbazide group for investigated 7-substituted phenylsemicarbazides **4,6,8** and the basic 7-unsubstituted phenylsemicarbazide **2**.

