

Supplementary materials for:

The substituent effect on the radical scavenging activity of apigenin

Yan-Zhen Zheng^{1,‡}, Da-Fu Chen^{1,*‡}, Geng Deng² and Rui Guo¹

1 College of Bee Science, Fujian Agriculture and Forestry University, Fuzhou 350002, P. R. China;
yanzhenzheng@fafu.edu.cn (Y.Z.Z.); dfchen826@fafu.edu.cn (D.F.C.); rui_0508@163.com (R.G.)

2 Key Laboratory of Bioorganic Phosphorous Chemistry and Chemical Biology (Ministry of
Education), Department of Chemistry, Tsinghua University, Beijing 100084, P. R. China;
dengg13@mails.tsinghua.edu.cn (G.D.)

‡ These authors contributed equally to this work

* Correspondence: dfchen826@fafu.edu.cn (D.F.C.); Tel.: +86-0591-8378-9482

12

13

14 **Table S1.** The linear formulas related with the Hammett sigma constants and the BDE/IP/PA. In the
 15 formulas, y represents the BDE/IP/PA, x represents the Hammett sigma constants

	gas	benzene	water
BDE(4'-OH) vs σ_p	$y = 14.5x + 368.5$	$y = 15.7x + 372.0$	$y = 18.1x + 367.7$
BDE(5-OH) vs σ_p	$y = 7.4x + 431.0$	$y = 6.7x + 425.4$	$y = 8.1x + 391.0$
BDE(7-OH) vs σ_m	$y = 9.9x + 386.7$	$y = 13.2x + 390.1$	$y = 19.9x + 386.5$
IP vs σ_p	$y = 86.4x + 774.5$	$y = 84.3x + 666.6$	$y = 79.3x + 578.2$
PA(4'-OH) vs σ_p	$y = -30.0x + 1355.0$	$y = -21.4x + 400.4$	$y = -10.5x + 148.7$
PA(5-OH) vs σ_m	$y = -46.4x + 1446.1$	$y = -38.2x + 470.3$	$y = -19.0x + 162.4$
PA(7-OH) vs σ_m	$y = -46.9x + 1376.2$	$y = -37.9x + 411.7$	$y = -15.9x + 140.9$


16

17

18

19

20




21

22 **Figure S1.** The energies and distributions of HOMO and LUMO orbitals for the investigated
23 compounds in the gas phase.

24

25

26



27

28 **Figure S2.** DAM for apigenin and its derivatives. The inset is definition of the four regions in DAM.

29