

Supplementary Materials

Single Production of Kojic Acid by *Aspergillus flavus* and the Revision of Flufuran

Antonius R B Ola^{1,2,*}, Gema Metboki¹, Catherine D Lay¹, Yoseph Sugi¹, Philipi De Rozari¹, Dodi Dharmakusuma^{1,2} and Euis Holisotan Hakim³

¹ Chemistry Department, Faculty of Science and Engineering, Nusa Cendana University, Kupang, Indonesia; antonius.ola@staf.undana.ac.id

² Integrated Research Center Laboratory (Biosains), University of Nusa Cendana 2; ola.antonius@gmail.com

³ Department of Chemistry, Institute of Technology Bandung, Jl. Ganesha 10, Bandung 40132

* Correspondence: antonius.ola@staf.undana.ac.id

Table of Contents

¹ H NMR spectrum of 1 in methanol- <i>d</i> ₄	3
¹³ C NMR spectrum of 1 in methanol- <i>d</i> ₄	4
DEPT spectrum of 1 in methanol- <i>d</i> ₄	5

Figure S1. ^1H NMR spectrum of **1** (methanol- d_4)




Figure S2. ^{13}C NMR spectrum of **1** (methanol- d_4)




Figure S3. DEPT spectrum of **1** (methanol-*d*₄)

