

Supporting Information

Stereochemistry of *N*-acyl-5*H*-dibenzo[*b,d*]azepin-7(6*H*)-ones

Arisa Chiba ¹, Ryoko Tanaka ¹, Mayuno Hotta ¹, Kayo Nakamura ¹, Kosho Makino ², Hidetsugu Tabata ³, Tetsuta Oshitari ³, Hideaki Natsugari ⁴ and Hideyo Takahashi ^{1,*}

¹ Faculty of Pharmaceutical Sciences, Tokyo University of Science, 2641 Yamazaki, Noda-shi, Chiba 278-8510, Japan

² Research Institute of Pharmaceutical Sciences, Musashino University, 1-1-20 Shinmachi, Nishitokyo-shi, Tokyo 202-8585, Japan

³ Faculty of Pharma Sciences, Teikyo University, 2-11-1 Kaga, Itabashi-ku, Tokyo 173-8605, Japan

⁴ Graduate School of Pharmaceutical Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan


* Correspondence: hide-tak@rs.tus.ac.jp; Tel.: +81-47-121-4436

Contents


1. ¹ H and ¹³ C NMR Spectra.....	S1
--	----

1. ¹H and ¹³C NMR Spectra



¹H NMR (600 MHz, CDCl₃, ppm) of **4**





¹³C NMR (150 MHz, CDCl₃, ppm) of **4**




¹H NMR (600 MHz, CDCl₃, ppm) of **5**









¹³C NMR (150 MHz, CDCl₃, ppm) of **5**




¹H NMR (600 MHz, CDCl₃, ppm) of **1b**








¹³C NMR (150 MHz, CDCl₃, ppm) of **1b**





¹H NMR (600 MHz, CDCl₃, ppm) of 6




¹³C NMR (150 MHz, CDCl₃, ppm) of **6**





¹H NMR (600 MHz, CDCl₃, ppm) of **2a**




¹³C NMR (150 MHz, CDCl₃, ppm) of **2a**





¹H NMR (600 MHz, CDCl₃, ppm) of **2b**



¹³C NMR (150 MHz, CDCl₃, ppm) of **2b**



¹H NMR (600 MHz, CDCl₃, ppm) of **2c**



¹³C NMR (150 MHz, CDCl₃, ppm) of **2c**

