



'Zaomeiguixiang'



'Moldova'



'Tamina'



'QiuHong'



'Xiangfei'



'Italia'



'Muscat of Alexandria'

Figure S1. Photographs of seven table grape berries at harvest.




Figure S2. Total soluble solids (TSS) and pH of seven table grape cultivars during berry development in 2013 and 2014.

Table S1. Real-time qPCR primers used in the present study.

Gene	Sense	Antisense	Reference	GenBank accession
<i>EF1-α</i>	GTGCGTCATAGTTTCTGCCTTCTTCCTTG	CTCAACCAGTTATCTGCCACCGCCTATC	[27]	EC959059
<i>UBQ-L40</i>	CATAACATTTGCGGCAGATCA	TGGTGGTATTATTGAGCCATCCTT	[27]	EC929411
<i>Actin</i>	GCATCCCTCAGCACCTTCCAGCAG	CCACCTCAACACATCTCCATGTCAACC	[27]	EC969944
<i>VvLOXA</i>	GCAAATCAAAGGGACAACGCTGTATGG	TGCTTCCACTGCGGCTTCC	[38]	FJ858255
<i>VvLOXO</i>	TTCCACCCACTCGCCTGATG	GCACCGCACCTGTTCTTCG	[38]	FJ858257
<i>VvHPL1</i>	AAGTACACCGGCGACATTGAG	AGCTCTTTACCCTGGCGTGTTG	[27]	HM627632
<i>VvADH1</i>	TCCGTTCTCAGAGATCAACAA	ACTCTCTCATCTCAAGATATTCTATGG	[40]	AF194173
<i>VvADH2</i>	ATTCCAGTCGGCATAAGTGT	TTGCAACTGCATAGACATTGTT	[40]	AF194174
<i>VvAAT</i>	TTAATTCAGGTGACCCGATT	TCTCCATACACATGCCATTAG	[27]	AAW22989