

E-selectin, ICAM-1, and ET-1 biomarkers address the concern of the challenging diagnosis of interstitial lung disease in patients with autoimmune diseases

Verónica Pulito-Cueto^{1,2*‡}, Sara Remuzgo-Martínez^{3*}, Fernanda Genre^{3*}, Belén Atienza-Mateo^{1,2}, Víctor M. Mora-Cuesta^{1,4}, David Iturbe-Fernández^{1,4}, Leticia Lera-Gómez⁵, María Sebastián Mora-Gil^{1,2}, Virginia Portilla^{1,2}, Alfonso Corrales^{1,2}, Ricardo Blanco^{1,2}, José M. Cifrián^{1,4,6}, Miguel A. González-Gay^{3,6,7**} and Raquel López-Mejías^{1,2**}

Additional file S1

Table S1. Serum levels of E-selectin, ICAM-1, and ET-1 in all individuals of the study.

	<i>E-selectin serum levels (ng/mL)</i> (Mean ± SD)	<i>ICAM-1 serum levels (ng/mL)</i> (Mean ± SD)	<i>ET-1 serum levels (pg/mL)</i> (Mean ± SD)
AD-ILD⁺	81.42 ± 30.65	532.60 ± 121.90	1.27 ± 0.61
RA-ILD⁺	77.55 ± 22.85	556.40 ± 125.40	1.17 ± 0.33
SSc-ILD⁺	85.11 ± 36.79	507.70 ± 116.20	1.37 ± 0.80
AD-ILD⁻	60.72 ± 19.07	439.0 ± 77.03	0.99 ± 1.09
RA-ILD⁻	57.27 ± 14.55	465.40 ± 80.55	0.86 ± 0.21
SSc-ILD⁻	64.87 ± 23.09	404.30 ± 57.30	1.14 ± 0.39
IPF	76.75 ± 20.81	527.40 ± 75.48	1.09 ± 0.38

ICAM-1: intercellular adhesion molecule 1; ET-1: endothelin 1; SD: standard deviation; AD: autoimmune disease; ILD: interstitial lung disease; RA: rheumatoid arthritis; SSc: systemic sclerosis; IPF: idiopathic pulmonary fibrosis.