

Supplementary Materials

L-asparaginase conjugates from the hyperthermophilic archaea *Thermococcus sibiricus* with improved biocatalytic properties

Natalia V. Dobryakova ^{1,2}, Maria V. Dumina ¹, Alexander A. Zhgun ¹, Marina V. Pokrovskaya ³, Svetlana S. Aleksandrova ³, Dmitry D. Zhdanov ^{1,3} and Elena V. Kudryashova ^{1,2,*}

¹ Group of Fungal Genetic Engineering, Federal Research Center "Fundamentals of Biotechnology of the Russian Academy of Sciences", 117312 Moscow, Russia; natdobryak@gmail.com (N.V.D.); duminamaria@gmail.com (M.V.D.); zzhgun@mail.ru (A.A.Z.)

² Chemical Faculty, Lomonosov Moscow State University, Leninskie Gory St. 1, 119991 Moscow, Russia

³ Laboratory of Medical Biotechnology, Institute of Biomedical Chemistry, 119121 Moscow, Russia, ivan1190@yandex.ru (M.V.P.); v-aleksandrov@yandex.ru (S.S.A.), zhdanovdd@gmail.com (D.D.Z.);

* Correspondence: helenakoudriachova@yandex.ru

Content

1. Figure S1. SDS-PAGE and native electrophoresis for the TsA enzyme.




Figure S1. SDS-PAGE for the TsA enzyme with oxidised glutathione (A) and heating times at 90 °C (B). Native electrophoresis of wild-type TsA and the mutant (C).

2. Figure S2. TIC and UV chromatograms of TsA




Figure S2. (A) Results of total ion current chromatogram (TIC chromatogram), 0.53 µg TsA; (B) UV chromatogram, 0.53 µg TsA, 280 nm.

3. Figure S3. Mass spectra of the TsA.




Figure S3. (A) Extracted mass spectrum of 0.53 µg of TsA over a chromatography interval of 7.6 to 7.9 minutes; (B) Deconvolution of the extracted mass spectrum of 0.53 µg of TsA over a chromatography interval of 7.6 to 7.9 minutes.

4. Figure S4. SDS-PAGE for TsA and TsA-PEG.




Figure S4. SDS-PAGE for TsA and TsA-PEG.