

Supplementary materials

Methyl 2-(1-methyl-3-oxoisoindolin-1-yl)acetate

Antonia Di Mola,^a Antonio Macchia,^a Laura Palombi^b and Antonio Massa^{a,*}

^aDipartimento di Chimica e Biologia "A. Zambelli", Università degli studi di Salerno, Via Giovanni Paolo II, 84084-Fisciano (SA), Italy Tel.: +39-089969565,

E-mail: amassa@unisa.it, <http://docenti.unisa.it/antonio.massa>

^bDipartimento di scienze fisiche e chimiche, Università degli studi dell'Aquila, via Vetoio, 67100 Coppito (AQ).

Contents:

Figure S1: ¹H NMR spectrum (400 MHz, CDCl₃) of compound **2**.

Figure S2: Expansion of the aromatic region of ¹H NMR spectrum (400 MHz, CDCl₃) of compound **2**.

Figure S3: ¹³C NMR spectrum (100 MHz, CDCl₃) of compound **2**.




Figure S1: ¹H NMR spectrum (400 MHz, CDCl₃) of compound 2.




Figure S2: Expansion of the aromatic region of ¹H NMR spectrum (400 MHz, CDCl₃) of compound 2.




Figure S3: ¹³C NMR spectrum (100 MHz, CDCl₃) of compound 2

List of impurities

¹H NMR: 7.26 ppm residual chloroform

1.75 ppm water

1.25 ppm greese

0.81 unknown

¹³C NMR: 29 ppm greese