

Fig. S1. Italian common bean (*Phaseolus vulgaris* L.) landraces used in this study. The letter after the sample name indicates the phaseolin type.



1_Fagiolino_S



2_Da-cucina_S



3_Napoletani_S



4_Cannelini-napoletani_S



5_Tondino_S



6_Faeto_S



7_Gradoli_S



8_Carrubara_S



9_Fagiuline_S



10_Fafodde_S



11_Cannellino_S



12_Fagiolo-napoletano_S



13_Fasule_S



14_Paesani_S



15_Monti-Dauni_S



16_Mangiatutto_S



17_Piattella_S



18_Corno-di-capra_S



19_Zolfino_S



20_Locale-di-Accettura_C



21_Mangiatutto-Sx015_T



22_Mangiatutto-Mx200_T




23_Saporro_T



24_Mangiatutto_T



25_Saluggia_T



26_Cosaruciaru_T



27_Marconi-nero_T



28_Cannellino_T



29_Villata_C



30_Verdolino_T



31_Ciuoto_T



32_Tabacchino_T



33_Riso_C



34_Billò_C



35_Badda-nero_C



36_Fasule_C



37_Ciammarrucoli_C



38_Poverello_C



39_Nasieddu_T



40_Vaianelli_C



41_Tuvagliedda_T



42_Tuvagliedda-rossa_T



43_Fagiolino-rosso_C



44_Drià_C



45_Maruchedda_C



46_San-Michele_T



47_Riso-giallo_T



48_Monaco_C



49_Munachedda_C



50_Pezzati_T