

Correction

## Correction: Structural Studies of the Lipopolysaccharide from the Fish Pathogen *Aeromonas veronii* Strain Bs19, Serotype O16. *Mar. Drugs* 2014, 12, 1298–1316

Anna Turska-Szewczuk <sup>1,\*</sup>, Katarzyna A. Duda <sup>2</sup>, Dominik Schwudke <sup>3</sup>, Agnieszka Pekala <sup>4</sup>, Alicja Kozinska <sup>4</sup> and Otto Holst <sup>2</sup>

<sup>1</sup> Department of Genetics and Microbiology, M. Curie-Sklodowska University, Akademicka 19, Lublin 20-033, Poland

<sup>2</sup> Division of Structural Biochemistry, Research Center Borstel, Leibniz-Center for Medicine and Biosciences, Parkallee 4a/c, Borstel D-23845, Germany; E-Mails: kduda@fz-borstel.de (K.A.D.); oholst@fz-borstel.de (O.H.)

<sup>3</sup> Division of Bioanalytical Chemistry, Research Center Borstel, Leibniz-Center for Medicine and Biosciences, Parkallee 10, Borstel D-23845, Germany; E-Mail: dschwudke@fz-borstel.de

<sup>4</sup> Department of Fish Diseases, National Veterinary Research Institute, Partyzantow 57, Pulawy 24-100, Poland; E-Mails: a.pekala@piwet.pulawy.pl (A.P.); koala@piwet.pulawy.pl (A.K.)

\* Author to whom correspondence should be addressed; E-Mail: aturska@hektor.umcs.lublin.pl; Tel.: +48-81-537-50-18; Fax: +48-81-537-59-59.

Received: 25 June 2014; in revised form: 30 June 2014 / Accepted: 4 July 2014 /

Published: 5 September 2014

---

We found one editorial mistake in our published paper [1]. In Line 2 of Table 4, the same composition of sugars is given for the C4 and C5 species (in the C5 species, one residue: 6dHexNAc has been missed). The correct composition for the C5 species is: [6dHexNAc6dHexHexHexNAc]6dHexNAc. We apologize for any inconvenience caused to our readers. The corrected table is as follows:

**Table 4.** Composition of the main species present in the charge deconvoluted ESI FT-ICR MS (negative ion mode) of the lower molecular mass fraction of the degraded PS isolated from the LPS of *A. veronii* Bs19, recorded with unspecific fragmentation. Mass fragments (marked with capital letters) are labeled according to the nomenclature of Domon and Costello [33].

Species	M <sub>measured</sub>	M <sub>calculated</sub>	Composition
C <sub>4</sub>	716.286	716.284	[6dHexNAc6dHexHexHexNAc]
C <sub>5</sub>	903.372	903.369	[6dHexNAc6dHexHexHexNAc]6dHexNAc
C <sub>6</sub>	1049.433	1049.427	[6dHexNAc6dHexHexHexNAc]6dHexNAc6dHex
C <sub>7</sub>	1211.486	1211.479	[6dHexNAc6dHexHexHexNAc]6dHexNAc6dHexHex
C <sub>8</sub>	1414.576	1414.558	[6dHexNAc6dHexHexHexNAc] <sub>2</sub>
C <sub>9</sub>	1601.669	1601.643	[6dHexNAc6dHexHexHexNAc] <sub>2</sub> 6dHexNAc
C <sub>10</sub>	1747.724	1747.700	[6dHexNAc6dHexHexHexNAc] <sub>2</sub> 6dHexNAc6dHex
C <sub>12</sub>	2112.830	2112.831	[6dHexNAc6dHexHexHexNAc] <sub>3</sub>
Z <sub>10</sub>	1827.612	1827.603	Hep <sub>5</sub> Hex <sub>3</sub> HexNKdo-H <sub>2</sub> O
Z <sub>12</sub>	2192.748	2192.734	[HexHexNAc]Hep <sub>5</sub> Hex <sub>3</sub> HexNKdo-H <sub>2</sub> O
Z <sub>14</sub>	2525.895	2525.876	[6dHexNAc6dHexHexHexNAc]Hep <sub>5</sub> Hex <sub>3</sub> HexNKdo-H <sub>2</sub> O
Z <sub>16</sub>	2891.010	2891.007	[6dHexNAc6dHexHex <sub>2</sub> HexNAc <sub>2</sub> ]Hep <sub>5</sub> Hex <sub>3</sub> HexNKdo-H <sub>2</sub> O
Z <sub>17</sub>	3037.077	3037.065	[6dHexNAc6dHex <sub>2</sub> Hex <sub>2</sub> HexNAc <sub>2</sub> ]Hep <sub>5</sub> Hex <sub>3</sub> HexNKdo-H <sub>2</sub> O
Z <sub>18</sub>	3224.159	3224.145	[6dHexNAc6dHexHexHexNAc] <sub>2</sub> Hep <sub>5</sub> Hex <sub>3</sub> HexNKdo-H <sub>2</sub> O

## Reference

- Turska-Szewczuk, A.; Duda, K.A.; Schwudke, D.; Pekala, A.; Kozinska, A.; Holst, O. Structural Studies of the Lipopolysaccharide from the Fish Pathogen *Aeromonas veronii* Strain Bs19, Serotype O16. *Mar. Drugs* **2014**, *12*, 1298–1316.

© 2014 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).