

Supplementary material

The Comparative Analysis of Antiviral Activity of Native and Modified Fucoidans from Brown Algae *Fucus evanescens* In Vitro and In Vivo

Natalya V. Krylova ^{1,*}, Svetlana P. Ermakova ², Vyacheslav F. Lavrov ³, Irina A. Leneva ³, Galina G. Kompanets ¹, Olga V. Iunikhina ¹, Marina N. Nosik ³, Linna K. Ebralidze ³, Irina N. Falynskova ³, Artem S. Silchenko ² and Tatyana S. Zaporozhets ¹

¹ G.P. Somov Institute of Epidemiology and Microbiology, Vladivostok 690087, Russia; galkom1965@gmail.com (G.G.K.); olga_jun@inbox.ru (O.V.I.); niiem_vl@mail.ru (T.S.Z.)

² G.B. Elyakov Pacific Institute of Bioorganic Chemistry, Vladivostok 690022, Russia; swetlana_e@mail.ru (S.P.E.); artem.silchenko@yandex.ru (A.S.S.)

³ I.I. Mechnikov Research Institute of Vaccines and Sera, Moscow 105064, Russia; v.f.lavrov@inbox.ru (V.F.L.); wnyfd385@yandex.ru (I.A.L.); mnossik@yandex.ru (M.N.N.); lina.lidze@gmail.com (L.K.E.); falynskova@mail.ru (I.N.F.)

* Correspondence: krylovanatalya@gmail.com; Tel.: +7-9084-486-423;




Figure S1. ¹H NMR spectra of native fucoidan FeF from *F. evanescence* (A) and its derivative FeHMP obtained after treatment with fucoidanase FFA1 from marine bacterium *Formosa algae* (B).