

Supplementary Materials

Pharmacokinetics and Metabolism Study of Deep-Sea-Derived Butyrolactone I in Rats by UHPLC-MS/MS and UHPLC-Q-TOF-MS

Liang Wu ¹, Chun-Lan Xie ², Xian-Wen Yang ^{2,*} and Gang Chen ^{1,*}

¹ School of Pharmacy, Fudan University, Shanghai, China; yhdhospital@163.com

² Key Laboratory of Marine Biogenetic Resources, Third Institute of Oceanography, Ministry of Natural Resources, 184 Daxue Road, Xiamen, 361005 Fujian, China; xiechunlanxx@163.com

* Correspondence: yangxianwen@tio.org.cn (X.-W.Y.); hhcy_educy@163.com (G.C.); Tel.: +86-592-219-5319 (X.-W.Y.); +86-21-51980168 (G.C.).

Content

Figure S1. ¹H NMR spectrum of **1** in MeOD.

Figure S2. ¹³C NMR spectrum of **1** in MeOD.

Figure S3. HSQC NMR spectrum of **1** in MeOD.

Figure S4. HMBC NMR spectrum of **1** in MeOD.




Figure S1. ¹H NMR spectrum of **1** in MeOD.




Figure S2. ^{13}C NMR spectrum of **1** in MeOD.




Figure S3. HSQC NMR spectrum of **1** in MeOD.




Figure S4. HMBC NMR spectrum of **1** in MeOD.