

Supporting Information

The Inhibition Effect and Mechanism of Staurosporine Isolated from *Streptomyces* sp. SNC087 Strain on Nasal Polyp

Grace Choi ^{1,*}, Eun-Young Lee ², Dawoon Chung ¹, Kichul Choi ¹, Woon-Jong Yu ¹, Sang-Jip Nam ², Seong-Kook Park ³ and Il-Whan Choi ^{4,*}

¹ Department of Microbial Resources, National Marine Biodiversity Institute of Korea, Seocheon, 33662, Republic of Korea

² Department of Chemistry and Nanoscience, Ewha Womans University, Seoul, 03760, Republic of Korea

³ Department of Otorhinolaryngology-Head & Neck Surgery, Busan Paik Hospital, Inje University College of Medicine, Busan 47392, Republic of Korea

⁴ Department of Microbiology and Immunology, Inje University College of Medicine, Busan 47392, Republic of Korea

* Correspondence: gchoi@mabik.re.kr, cihima@inje.ac.kr

Table of Contents

Figure S1 ^1H NMR spectrum (400 MHz, CDCl_3) of Staurosporine (1)	S3
Figure S2 ^{13}C NMR spectrum (100 MHz, CDCl_3) of Staurosporine (1)	S4

Figure S1 ^1H NMR spectrum (400 MHz, CDCl_3) of Staurosporine (**1**)




Figure S2 ^{13}C NMR spectrum (100 MHz, CDCl_3) of Staurosporine (**1**)

