Prediction of indoor air exposure from outdoor air quality using an artificial neural network model for inner city commercial buildings





Figure S1. Time series of neural network training Mc2 run 2 NO₂.



Response of Output Element 1 for Time-Series 1

Figure S2. Time series of neural network training Mc3 run 1 NO2.




Figure S3. Mc3 run 1 NO₂ regression of trained output data set.




Figure S4. Time series of neural network training Mc3 run 2 NO₂.





Figure S5. M_c3 run 2 NO₂ regression of trained output data set.



Response of Output Element 1 for Time-Series 1

Figure S6. Time series of neural network training Nt2 run 1 NO2.




Figure S7. Nt2 run 1 NO2 regression of trained output data set.




Figure S8. Nt2 run 2 NO2 regression of trained output data set.




Figure S9. Time series of neural network training Mc3 run 1 PM_{2.5}.




Figure S10. M_c3 run 1 PM_{2.5} regression of trained output data set.




Figure S11. Time series of neural network training Mc3 run 2 (Street) PM_{2.5}.




Figure S12. Time series of neural network training Mc3 run 2 (roof) PM2.5.

Response of Output Element 1 for Time-Series 1




Figure S13. Time series of neural network training Mc3 run 2 PM_{2.5}.




Figure S14. Time series of neural network training Nt2 run 1 PM2.5.




Figure S15. Time series of neural network training Nt2 run 2 PM_{2.5}.




Figure S16. PM_{2.5} outdoor concentrations Nt2 run 1 and run 2.




Figure S17. PM_{2.5} indoor concentrations Nt2 run 1 and run 2.

 \bigcirc 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (http://creativecommons.org/licenses/by/4.0/).