

Electronic supplementary information for

Carbon/Graphene-Modified Titania with Enhanced Photocatalytic Activity under UV and Vis Irradiation

Kunlei Wang ¹, Maya Endo-Kimura ¹, Raphaëlle Belchi ^{2,3}, Dong Zhang ⁴, Aurelie Habert ², Johann Bouclé ³, Bunsho Ohtani ^{1,4}, Ewa Kowalska ^{1,4,*} and Nathalie Herlin-Boime ^{2,*}

¹ Institute for Catalysis (ICAT), Hokkaido University, N21 W10, Sapporo 001-0021, Japan; kunlei@cat.hokudai.ac.jp (K.W.); m_endo@cat.hokudai.ac.jp (M.E.-K.); ohtani@cat.hokudai.ac.jp (B.O.)

² IRAMIS—NIMBE UMR 3685, Université Paris Saclay, CEA Saclay, 91191 Gif/Yvette CEDEX, France; raphaelle.belchi@cea.fr (R.B.); aurelie.habert@cea.fr (A.H.)

³ Univ. Limoges, CNRS, XLIM, UMR 7252, F-87000 Limoges, France; johann.boucle@unilim.fr

⁴ Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810, Japan; zhang.d@cat.hokudai.ac.jp

* Correspondence: kowalska@cat.hokudai.ac.jp (E.K.); nathalie.herlin@cea.fr (N.H.-B.)




Figure S1. Raman mapping of the G-TiO₂ sample illustrating the presence of C species in localized places.




Figure S2. RDB-PA spectra of (a) G-TiO₂, (b) TiO₂, (c) P25 (anatase/rutile), and (d) ST01 (anatase).