

Supplementary Materials: Bacterial Infection and Implant Loosening in Hip and Knee Arthroplasty: Evaluation of 209 Cases

Ulrike Dapunt, Stephanie Radzuweit-Mihaljevic, Burkhard Lehner, Gertrud Maria Haensch and Volker Ewerbeck

Table S1. Causative agents detected in patients with prosthetic joint infection of the hip and knee.

Causative Agents Detected	Bacteria Species in Detail	Number
<i>S. aureus</i>	—	43
<i>S. epidermidis</i>	—	53
		20 (in detail)
<i>other staphylococcus species</i>	<i>S. hominis,</i> <i>S. haemolyticus,</i> <i>S. caprae,</i> <i>S. lugdunensis,</i> <i>S. gallolyticus,</i> <i>S. capitis</i>	4 3 1 5 1 6
<i>Streptococcus</i> sp.	<i>St. agalactiae,</i> <i>St. pneumoniae,</i> <i>St. parasanguinis,</i> <i>St. dysgalactiae,</i> <i>St. oralis,</i> <i>St. anginosus,</i> <i>St. canis,</i> <i>St. Group G,</i> <i>St. mitis,</i> <i>S. pyogenes</i>	1 1 1 2 4 2 1 1 1 2
<i>Enterococcus</i> species	—	19
		30 (in detail)
others	<i>Enterobacter cloacae,</i> <i>Parvimonas micra,</i> <i>Escherichia coli,</i> <i>Acinetobacter baumannii,</i> <i>Bacillus</i> sp., <i>Pantoea</i> sp., <i>Sphingobacterium multivorum,</i> <i>Finegoldia magna,</i> <i>Phialemonium</i> sp., <i>Clostridium perfringens,</i> <i>Rothiadentocariosa,</i> <i>Serratia</i> sp., <i>Actinomyces oris,</i> <i>Klebsiella pneumoniae,</i> <i>Pseudomonas aeruginosa,</i> <i>Micrococcus latus,</i> <i>Proteus mirabilis,</i> <i>Propionibacterium</i> sp.	3 2 4 2 3 1 1 1 1 1 1 1 2 2 1 2 1 1