

The plant functional groups and species composition of the study area.

Plant Functional Group	Species Composition
Grasses (IV: 22.31%)	<i>Agropyron cristatum</i> (L.) Gaertn.
	<i>Cleistogenes squarrosa</i> (Trin.) Keng
	<i>Enneapogon borealis</i> (Griseb.) Honda
	<i>Eragrostis minor</i> Host
	<i>Poa sphondylodes</i> Trin.
	<i>Setaria viridis</i> (L.) P.Beauv.
	<i>Stipa capillata</i> L.
Forbs (IV: 77.69%)	<i>Tragus racemosus</i> (L.) All.
	<i>Ajania achilloides</i> Poljak. ex Grubov.
	<i>Allium mongolicum</i> Regel
	<i>Artemisia frigida</i> Willd.
	<i>Artemisia sacrorum</i> Ledeb.
	<i>Asparagus gobicus</i> Ivanova ex Grubov
	<i>Astragalus scaberrimus</i> Bunge
	<i>Bassia dasyphylla</i> (Fisch. & C.A.Mey.) O.Kuntze
	<i>Caragana opulens</i> Kom.
	<i>Caragana roborovskyi</i> Kom.
	<i>Caragana tibetica</i> Kom.
	<i>Convolvulus ammannii</i> Desr.
	<i>Cymbaria mongolica</i> Maxim.
	<i>Dracocephalum heterophyllum</i> Benth.
	<i>Gypsophila capituliflora</i> Rupr.
	<i>Heteropappus altaicus</i> Novopokr.
	<i>Iris tenuifolia</i> Pall.
	<i>Linum perenne</i> L.
	<i>Lycium chinense</i> Mill.
	<i>Oxytropis ochrocephala</i> Bunge
	<i>Peganum harmala</i> L.
	<i>Phlomis dentosa</i> Franch.
	<i>Potentilla chrysantha</i> Trev.
	<i>Potentilla discolor</i> Bunge
	<i>Reaumuria songarica</i> (Pall.) Maxim.
	<i>Salsola collina</i> Pall.
	<i>Torularia humilis</i> (C. A. Mey.) O. E. Schulz
	<i>Zygophyllum mucronatum</i> Maxim.