

Supplementary Materials

Unravelling the Role of Biochemical Compounds within the Hydrothermal Liquefaction Process of Real Sludge Mixtures

Marco Balsamo ¹, Francesca Di Lauro ^{1,2,*}, Maria Laura Alfieri ¹, Paola Manini ¹, Piero Salatino ², Fabio Montagnaro ^{1,*} and Roberto Solimene ³

¹ Department of Chemical Sciences, University of Naples Federico II, Complesso Universitario di Monte Sant'Angelo, 80126 Napoli, Italy; marco.balsamo@unina.it (M.B.); marialaura.alfieri@unina.it (M.L.A.); paola.manini@unina.it (P.M.)

² Department of Chemical, Materials and Production Engineering, University of Naples Federico II, Piazzale V. Tecchio 80, 80125 Napoli, Italy; piero.salatino@unina.it

³ Institute of Sciences and Technologies for Sustainable Energy and Mobility, National Research Council, Piazzale V. Tecchio 80, 80125 Napoli, Italy; roberto.solimene@stems.cnr.it

* Correspondence: francesca.dilauro2@unina.it (F.D.L.); fabio.montagnaro@unina.it (F.M.); Tel.: +39-081-674029 (F.M.)




Figure S1. HTL products separation steps.




Figure S2. ^1H , ^1H -COSY spectrum (DMSO- d_6) of the bio-crude fraction obtained at 350 °C and 10 min from pure TS.




Figure S3. ^1H , ^1H -COSY spectrum (D_2O) of the aqueous fraction obtained at 350 °C and 10 min from pure TS.




Figure S4. ^1H , ^1H -COSY spectrum (DMSO- d_6) of the bio-crude fraction obtained at 350 °C and 10 min from a mixed sludge composed by MS:TS 75:25.




Figure S5. ^1H , ^1H -COSY spectrum (D_2O) of the aqueous fraction obtained at 350 °C and 10 min from a mixed sludge composed by MS:TS 75:25.