

A new method for extracting individual plant biocharacteristics from high-resolution digital images

Saba Rabab^{1,2,*}, Edmond Breen², Alem Gebremedhin³, Fan Shi², Pieter Badenhorst⁴, Yi-Ping Phoebe Chen⁵ and Hans D. Daetwyler^{1,2}


- ¹ School of Applied Systems Biology, La Trobe University, Bundoora, Australia
- ² Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Australia.
- ³ Agriculture Victoria, Grain Innovation Park, Horsham, Australia.
- ⁴ Agriculture Victoria, Hamilton Centre, Hamilton, Australia.
- Department of Computer Science and Information Technology, La Trobe University, Bundoora, Australia.
- * Corresponding Author, Email: saba.rabab@agriculture.vic.gov.au



Supp. Fig. 1. Comparisons between the individual plant values of normalized values of fresh weights and areas for first 100 plants in a same range of [0, 100] for the field trial image taken on 9th May 2017.



Supp. Fig. 2. Comparisons between the individual plant values of normalized values of fresh weights and areas for first 100 plants in a same range of [0, 100] for the field trial image taken on 11th September 2017.



Supp. Fig. 3. Comparisons between the individual plant values of normalized values of fresh weights and areas for first 100 plants in a same range of [0, 100] for the field trial image taken on 20th November 2017.