

Figure S1. Schemes of the treatments used to evaluate MMP9 activity in different conditioned media (A) and HT29 cell viability after 24 h treatment with M2 medium with or without Nem, Cp5 and Cp17 (B).




Table S1. qRT-PCR primers used in this study.

Gene	Sequence forward (F)	Sequence reverse (R)
<i>IL6</i>	ACTCCTTCTCCACAAGCGCC	TGTGGGGCGGCTACATCTT
<i>IL8</i>	AAACCACCGGAAGGAACCAT	CCTTCACACAGAGCTGCAGAAA
<i>TNFα</i>	AAGCCTGTAGCCCATGTTGT	GAGGTACAGGCCCTCTGATG
<i>CCL5</i>	CCTCATTGCTAGGCCCTCT	GGTGTGGTGTCCCGAGGAAT
<i>IL10</i>	AAGACTTTCTTTCAAATGAA	GTTTTACAGGGAAGAAATC
<i>CCL2</i>	GCTCATAGCAGCCACCTTCA	GGACACTTGCTGCTGGTGAT
<i>VEGF</i>	ATGGCAGAAGGAGGAGGGCA	ATCGCATCAGGGGCACACAG
<i>CD206</i>	ACCTTCACAAGTATCCACACCATC	CTTCATCACCACACAATCCT
<i>GAPDH</i>	ACATCAAGAAGGTGGTGAAGCA	GTCAAAGGTGGAGGAGTGGGT