

Table S1. Antiplatelet drug use according to Med-Diet-Adherence.

	Adherence to Mediterranean Diet			$p_{\text{among all}}$	$p_{\text{low vs high}}$
	Low (Score 0-2 pts) (n=67)	Intermediate (Score 3-6 pts) (n=487)	High (Score 7-9 pts) (n=101)		
Aspirin (%)	19.4	14.0	11.9	0.376	0.191
Clopidogrel (%)	1.5	1.2	0	0.515	0.399
Ticlopidine (%)	1.5	1.0	0	0.537	0.399

Table S2. Mean TxB_2 and Pro-C3 according to aspirin, clopidogrel and ticlopidine use.

	Aspirin		p	Clopidogrel		p	Ticlopidine		p
	Yes (n=93)	No (n=562)		Yes (n=7)	No (n=648)		Yes (n=6)	No (n=649)	
TxB_2 (pg/ml)	184.0 \pm 32.7	184.9 \pm 32.0	0.809	188.9 \pm 15.4	184.8 \pm 32.2	0.735	182.9 \pm 4.4	184.8 \pm 32.2	0.882
Pro-C3 (ng/ml)	7.2 \pm 2.9	7.2 \pm 2.6	0.836	7.6 \pm 2.8	7.2 \pm 2.6	0.686	8.7 \pm 3.5	7.2 \pm 2.6	0.149