

Table S2. Inclusion of wheat, corn and soybean meal in trout, tilapia, marine fish and carp diets.

Country, year	Wheat	Corn ¹	Soybean meal
<i>Trout</i>			
UK, 2008	10		15
France, 2009	5–10 (7.5)	5–8 (6.5)	10–15 (12.5)
Greece, 2009	5–15 (10)	5–12 (8.5)	10–35 (22.5)
Norway, 2010	12		12
Average inclusion²	10.6 %	7.5%	15.5%
<i>Tilapia</i>			
Mexico 2009	15–20 (17.5)		30
Venezuela 2008	34		23
Ecuador 2009		20–30 (25)	40
Egypt 2008		30–40 (35)	20–25 (22.5)
Peru 2008		30	40
China 2008			0–25 (12.5)
Taiwan 2007			30–35 (32.5)
Vietnam 2008			30–60 (45)
USA 2008			30–35 (32.5)
Average inclusion	19.9%	30 %	30.9 %
<i>Marine fish</i>			
France 2009	5 – 10 (7.5)	10 – 18 (14)	15- 25 (20)
Greece 2009	5 – 15 (10)	5 – 12 (8.5)	10- 35 (22.5)
Spain 2009	1 – 5 (2.5)	4	
UK 2008	10		15
Average inclusion	7.6 %	8.8 %	19.2 %
<i>Carp³</i>			
China 2008	0-25 (13)	0-25 (13)	0-25 (13)
Egypt 2008	10-25 (17.5)	10-25 (17.5)	20-25 (22.5)
India 2006/7	50	50	5
Average inclusion	27%	27%	13.5 %

¹Corn data refers to corn gluten meal in case of trout and marine species.

²Average inclusion was estimated taking into account the average/country.

³Wheat data refers to wheat bran in case of carp.

