



Article

Modified FOLFIRINOX as a Second-Line Treatment for Patients with Gemcitabine-Failed Advanced Biliary Tract Cancer: A Prospective Multicenter Phase II Study


Yong-Pyo Lee, Sung Yong Oh, Kwang Min Kim, Se-Il Go, Jung Hoon Kim, Seok Jae Huh, Jung Hun Kang and Jun Ho Ji



Supplementary Figure S1. Kaplan–Meier curves of progression-free survival (A) and overall survival (B) for achieving objective responses. Kaplan–Meier curves of progression-free survival (C), and overall survival (D) for obtaining disease control.



Supplementary Figure S2. Kaplan–Meier curves of progression-free survival (A) and overall survival (B) according to the number of mFOLFIRINOX administration.



Supplementary Figure S3. Kaplan–Meier curves of progression-free survival (A) and overall survival (B) according to the type of biliary tract cancer. Kaplan–Meier curves of progression-free survival (C), and overall survival (D) according to the duration of first-line gemcitabine/cisplatin.

Supplementary Table S1. Comparison of dose, response rate, and survival results according to studies.

	Lines of treatment	Study	Disease	Chemotherapies	PFS	OS	ORR	DCR
A. Lamarca, et al.	first-line	phase III	BTC	Oxaliplatin 85 mg/m ² , Leucovorin 350 mg/m ² , 5-FU bolus 400 mg/m ² , 5-FU infusion 2,400 mg/m ² over 46 hours	4.0 months	6.2 months	5%	33%
J. Phelip, et al.	first-line	phase II	BTC	Oxaliplatin 85 mg/m ² , Irinotecan 180 mg/m ² , Leucovorin 400 mg/m ² , 5-FU bolus omitted 5-FU infusion 2,400 mg/m ² over 46 hours	6.2 months	11.7 months	25%	66%
A. Belkouz, et al.	second-line	phase II	BTC	Oxaliplatin 85 mg/m ² , Irinotecan 180 mg/m ² , Leucovorin 400 mg/m ² , 5-FU bolus 400 mg/m ² 5-FU infusion 2,400 mg/m ² over 46 hours	6.2 months	10.7 months	10%	66%
Liu-Fang Ye, et al.	second-line	phase II	BTC	Oxaliplatin 65 mg/m ² , Irinotecan 150 mg/m ² Leucovorin 400 mg/m ² , 5-FU bolus omitted 5-FU infusion 2,400 mg/m ² over 46 hours	6.7 months	13.2 months	26%	80%
Present study	second-line	phase II	BTC	Oxaliplatin 65 mg/m ² , Irinotecan 135 mg/m ² Leucovorin 400 mg/m ² , 5-FU bolus omitted 5-FU infusion 1,000 mg/m ² /day for two days	2.8 months	6.2 months	14%	61%
Se-Il Go, et al.	second-line	phase III	PAC	Oxaliplatin 65 mg/m ² , Irinotecan 135 mg/m ² Leucovorin 400 mg/m ² , 5-FU bolus omitted 5-FU infusion 1,000 mg/m ² /day for two days	5.2 months	9.2 months	15%	67%
				BTC , biliary tract cancer; 5-FU , 5-fluorouracil; PFS , progression-free survival; OS , overall survival; ORR , overall response rate; DCR , disease control rate; PAC , pancreatic adenocarcinoma.				