

Supplementary Materials: CeO₂ for Water Remediation: Comparison of Various Advanced Oxidation Processes

Roberto Fiorenza ¹, Stefano Andrea Balsamo ¹, Luisa D'Urso ¹, Salvatore Sciré ¹, Maria Violetta Brundo ², Roberta Pecoraro ², Elena Maria Scalisi ², Vittorio Privitera ³ and Giuliana Impellizzeri ^{3,*}

¹ Department of Chemical Sciences, University of Catania, Viale A. Doria 6, 95125 Catania, Italy; rfiorenza@unict.it (R.F.); stefano.balsamo@phd.unict.it (S.A.B.); ldurso@unict.it (L.D.) sscire@unict.it (S.S.)

² Department of Biological, Geological and Environmental Science, University of Catania, Via Androne 81, 95124 Catania, Italy; mvbrundo@unict.it (M.V.B.); roberta.pecoraro@unict.it (R.P.); scalisimariaelena90@gmail.com (E.M.S.)

³ CNR-IMM, Via S. Sofia 64, 95123 Catania, Italy; vittorio.privitera@cnr.it

* Correspondence: giuliana.impellizzeri@ct.infn.it; Tel.: +39 095 3785345

S1. Influence of the H₂O₂ concentration




Figure S1. Effect of the H₂O₂ concentration on the imidacloprid degradation after 3 hours with Fenton-like and photo-Fenton like tests utilizing the grey CeO₂ catalyst.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).