

Article

Four-Component Synthesis of 9*H*-Pyrimido[4,5-*b*]indoles Using Ammonium Iodide as the Nitrogen Source

 Yufeng Chen ¹, Ruitong Yang ¹, Fuhong Xiao ^{1,*} , Tong Li ¹, Guojiang Mao ² and Guo-Jun Deng ^{1,*} 
¹ Key Laboratory of Environmentally Friendly Chemistry and Application of Ministry of Education, Hunan Province Key Laboratory of Green Organic Synthesis and Application, College of Chemistry, Xiangtan University, Xiangtan 411105, China

² School of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang 453007, China

* Correspondence: fhxiao@xtu.edu.cn (F.X.); gjdeng@xtu.edu.cn (G.-J.D.)

Abstract: A four-component synthesis of 2-phenyl-9*H*-pyrimido[4,5-*b*]indoles was developed using indole-3-carboxaldehydes, aromatic aldehyde and ammonium iodide as the raw materials under transition-metal-free conditions. The pyrimidine ring was formed in one pot through [4 + 2] annulation reaction. Four C–N bonds were formed in one pot promoted by iodine and iodide additives. This work is highlighted by using two ammonium iodides as the sole nitrogen source.

Keywords: 9*H*-pyrimido[4,5-*b*]indole; aromatic aldehyde; ammonium salt; cycloaddition; transition metal free

1. Introduction

As an important part of nitrogen-containing heterocycles, pyrimidines are widespread in natural products and pharmaceuticals [1–3]. Among them, the 9*H*-pyrimido[4,5-*b*]indole motif has important application in many fields, such as anti-inflammatory, antimicrobial, antimalarial agents and cytotoxic inhibitors [4–6]. Furthermore, the core structure of 9*H*-pyrimido[4,5-*b*]indoles was found in many biologically active molecules, such as HSCex vivo expansion agent, pyruvate dehydrogenase kinases (PDHKs) inhibitors, epidermal growth factor receptor (EGFr) tyrosine kinase inhibitors, and endogenous hormones (Figure 1) [7–11]. Therefore, significant research effort has been focused on the synthesis of 9*H*-pyrimido[4,5-*b*]indole derivatives. According to the reported methods, the synthesis of 9*H*-pyrimido[4,5-*b*]indoles usually started with highly functionalized materials [12–15]. For example, 4-azido-5-phenylpyrimidine, 1,3,5-triazines, benzamidine, guanidine nitrate and *o*-nitrobiphenyl were successfully used as the starting materials for the synthesis of target products [16–19]. Furthermore, most reactions often use stoichiometric amount of strong acids or bases which do not meet the requirement of green chemistry. For example, Shkurko and co-workers developed a multi-step synthesis of 2-phenyl-9*H*-pyrimido[4,5-*b*]indole from indolin-2-one using sodium ethylate prepared from sodium as the base [20]. Therefore, efficient and green synthetic methods for constructing 9*H*-pyrimido[4,5-*b*]indole derivatives in one pot are highly desirable.



Citation: Chen, Y.; Yang, R.; Xiao, F.; Li, T.; Mao, G.; Deng, G.-J.

Four-Component Synthesis of 9*H*-Pyrimido[4,5-*b*]indoles Using Ammonium Iodide as the Nitrogen Source. *Catalysts* **2023**, *13*, 623. <https://doi.org/10.3390/catal13030623>

Academic Editors: Lianghua Zou and Xiaofeng Xia

Received: 20 February 2023

Revised: 17 March 2023

Accepted: 18 March 2023

Published: 20 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).





Figure 1. Representative biologically active 9*H*-pyrimido[4,5-*b*]indoles ((a) [7], (b) [9], (c) [8], (d) [11]).


In the last decades, the multi-component reactions (MCRs) have played an increasingly dominant role in the organic synthesis of wide and diverse compounds which provides atom economy, and featured productivity and easy execution [21–29]. MCRs usually start with simple and commercially available materials as the reaction substrates thus providing a shortcut for the synthesis of functional molecules. Meanwhile, ammonium salts are ideal nitrogen sources for the synthesis of nitrogen-containing heterocycles due to their low cost and ease of handling [30–36]. In 2009, Konakahara and coworkers developed a ZnCl₂-catalyzed three-component coupling reaction for the synthesis of 4,5-disubstituted pyrimidines from ammonium acetate, enamines and triethyl orthoformate (Scheme 1a) [37]. In recent years, our group also developed several MCRs for the synthesis of nitrogen-containing heterocycles using ammonium salts as the nitrogen source [38–40]. For example, in 2018, we reported a four-component procedure for the synthesis of quinazoline using anilines, aromatic aldehydes and ammonium iodide as starting materials (Scheme 1b) [41]. In 2020, we developed a base-promoted aerobic oxidation approach for the synthesis of benzimidazo[1,2-*a*]-1,3,5-triazines involving 2-aminobenzimidazoles, aromatic aldehydes and ammonium iodide [42]. However, in most cases, ammonium salts only provided a single nitrogen atom source in these multi-component reactions. When two nitrogen atom rings were formed, another organic nitrogen source such as amines is usually required. In 2019, Chen's group demonstrated a three-component reaction of ammonium acetate, chalcone and phenyl-methanol using copper (II) as the catalyst providing tri-substituted pyrimidines in moderate yields (Scheme 1c) [43]. In this process, both nitrogen atoms in the pyrimidine ring come from the ammonium acetate. However, for the synthesis of *N*-heterocycles containing multiple nitrogen atoms which are easy to form a stable complex with the transition metals but encounter difficulties in removing the toxic transition metals, it is particularly important to perform the reaction without the aid of transition-metal catalysts. In the pursuit of research on heterocycle construction using ammonium salts as the nitrogen source under transition-metal-free conditions, herein we describe a general four-component reaction for the synthesis of 2-phenyl-9*H*-pyrimido[4,5-*b*]indole from indole-3-carboxaldehydes, aromatic aldehydes and ammonium salt (Scheme 1d). In this reaction, both nitrogen atoms in the pyrimidine ring come from ammonium salt and four C–N bonds were formed in one pot in the absence of a transition metal catalyst.

One nitrogen atom comes from ammonium salt!

(a) Konakahara's work



(b) Our previous work




Both nitrogen atoms come from ammonium salt!

(c) Chen's work



(d) This work



Two equivs of NH₄I as nitrogen source
Transition-metal-free conditions


Four C–N bonds formation
40 examples, yields up to 96%

Scheme 1. *N*-heterocycles synthesis through ammonium salts ((a) [37], (b) [41], (c) [43], (d) this work).

2. Results

In order to explore the optimized reaction conditions, our study was initiated by using indole-3-carboxaldehyde (**1a**), benzaldehyde (**2a**) and ammonium iodide as model substrates and 20 mol% of iodine as the catalyst under oxygen atmosphere (Table 1). To begin with, potassium iodide was used as an additive to the reaction, affording the target product **3aa** in 45% yield (entry 1). Then, various iodide-containing additives were screened to find the appropriate promotor, among which NaIO₄ is the best candidate reagent to provide the product **3aa** in 76% yield (entries 2–5). Moderate yields were obtained when reactions were carried out in PhCl, toluene, mesitylene and PhCF₃ (entries 6–9). DMF, CH₃CN, NMP and pyridine as the reaction media completely quenched the reaction (entries 10–13). Furthermore, oxidants screening among others showed that DMSO was superior to others such as H₂O₂, *tert*-butyl hydroperoxide (TBHP), *di*-*tert*-butyl peroxide (DTBP), K₂S₂O₈, and Na₂S₂O₈ (entries 14–18). Diminished yield was obtained when the reaction was performed without I₂ (entry 19). Lower yield was obtained when the reaction was carried out in air atmosphere (entry 20). Finally, no desired product was detected when using NH₄Cl instead of NH₄I as nitrogen source for this kind of transformation (entry 21). We believed that iodine, periodate and oxygen were used as the oxidant. The iodine anion could be oxidized to iodine, which could further promote the oxidative aromatization process.

Table 1. Optimization of the reaction conditions ¹.




Entry	Additive	Oxidant	Solvent	Yield (%) ²
1	KI	DMSO	<i>o</i> -DCB	45
2	KIO ₃	DMSO	<i>o</i> -DCB	68
3	NaIO ₄	DMSO	<i>o</i> -DCB	76
4	I ₂ O ₅	DMSO	<i>o</i> -DCB	trace
5	NIS	DMSO	<i>o</i> -DCB	trace
6	NaIO ₄	DMSO	PhCl	65
7	NaIO ₄	DMSO	toluene	53
8	NaIO ₄	DMSO	mesitylene	40
9	NaIO ₄	DMSO	PhCF ₃	39
10	NaIO ₄	DMSO	DMF	trace
11	NaIO ₄	DMSO	CH ₃ CN	trace
12	NaIO ₄	DMSO	NMP	trace
13	NaIO ₄	DMSO	Pyridine	trace
14	NaIO ₄	H ₂ O ₂	<i>o</i> -DCB	38
15	NaIO ₄	TBHP	<i>o</i> -DCB	35
16	NaIO ₄	DTBP	<i>o</i> -DCB	30
17	NaIO ₄	K ₂ S ₂ O ₈	<i>o</i> -DCB	trace
18	NaIO ₄	Na ₂ S ₂ O ₈	<i>o</i> -DCB	15
19 ³	NaIO ₄	DMSO	<i>o</i> -DCB	58
20 ⁴	NaIO ₄	DMSO	<i>o</i> -DCB	63
21 ⁵	NaIO ₄	DMSO	<i>o</i> -DCB	0

¹ Reaction conditions: **1a** (0.2 mmol), **2a** (0.4 mmol), NH₄I (0.6 mmol), I₂ (0.04 mmol), additive (0.2 mmol), oxidant (0.4 mmol), solvent (0.6 mL), 150 °C, 16 h, oxygen. ² GC yield based on **1a**. ³ Without I₂. ⁴ Under air. ⁵ With NH₄Cl (0.6 mmol) instead of NH₄I.

With the optimal conditions established, the substrate scope of indole-3-carboxaldehydes was explored (Scheme 2). As depicted, various substitutes of indole-3-carboxaldehyde could participate in current reaction regardless of the electronic nature of substituents incorporated on the C4, C5, C6, C7 positions. The model reaction provided product **3aa** in 73% isolated yield. *N*-methylindole-3-carbaldehyde could also be used as the sub-

strate to give product **3ba** in 66% isolated yield. To our surprise, 4-chloro-1*H*-indole-3-carbaldehyde was the best effective substrate that the corresponding product **3ca** was obtained in 96% yield, whereas 4-bromo substituted indole-3-carbaldehyde afforded **3da** in 71% yield. Moreover, when fluoro, chloro and methyl substituents were located at C5, C6 and C7 positions, the counterparts were obtained in moderate to good yields (**3ea**, **3fa**, **3ga**, **3ia**, **3la** and **3ma**). Similarly, bromo substituent at C5 and C6 positions led to lower isolated yields (**3ha** and **3ja**). It was regrettable that 6-methoxy-1*H*-indole-3-carbaldehyde only provided product **3ka** in 23% yield. Further study revealed that 1*H*-pyrrolo[2,3-*b*]pyridine-3-carbaldehyde was also a viable substrate for this kind of reaction, leading to the corresponding product **3na** in 59% yield. Unfortunately, sulfur containing heterocyclic aldehyde such as benzo[*b*]thiophene-3-carbaldehyde was ineffective for this type of cyclization process.




Reaction conditions: **1** (0.2 mmol), **2a** (0.4 mmol), NH_4I (0.6 mmol), I_2 (0.04 mmol), NaIO_4 (0.2 mmol), DMSO (0.4 mmol), 1,2-dichlorobenzene (0.6 mL), 150°C , 16 h, oxygen atmosphere, isolated yield based on **1**.

Scheme 2. Scope of indole-3-carboxaldehydes.

Additional experiments revealed that the current protocol for 2-aryl-9*H*-pyrimido[4,5-*b*]indole synthesis was very effective and efficient, and various aromatic aldehydes derived from **2a** could also react well with indole-3-carboxaldehyde **1a** (Scheme 3). Electron-donating and electron-withdrawing groups decorated on the aryl ring were well tolerated,

of benzaldehyde with another ammonium iodide affords an imine intermediate **B**. The [4+2] annulations reaction of **A** and **B** formed an intermediate **C**. Oxidative dehydrogenation of the intermediate **C** in the presence of iodine/DMSO affords the final product 2-phenyl-9H-pyrimido[4,5-*b*]indole **3aa**.



Scheme 4. Plausible reaction pathway.

3. Discussion

In summary, we have developed a four-component strategy which can provide an efficient approach to various substituted pyrimido[4,5-*b*]indoles from commercially available indole-3-carboxaldehyde, aromatic aldehydes and ammonium iodide under simple reaction conditions. The C–H bond on the C2 position of indole was directly functionalized and used for a further cyclization process under metal-free conditions. Ammonium iodide was involved in this reaction and was used as two nitrogen sources for the pyrimidine ring construction. The reaction showed good functional group tolerance and wide substrate scope. This four-component mixture could be selectively assembled into the target products in one pot without the use of transition-metal catalyst.

4. Materials and Methods

4.1. General Information

All reactions were carried out under an atmosphere of oxygen unless otherwise noted. Column chromatography was performed using silica gel (200–300 mesh) or neutral alumina. ^1H NMR and ^{13}C NMR spectra were recorded on Bruker-AV (400 and 100 MHz, respectively, Hangzhou, China) instrument internally referenced to tetramethylsilane (TMS) and using chloroform-*d* (CDCl_3) and dimethyl sulphoxide-*d*₆ ($\text{DMSO-}d_6$) as solvent. The obtained spectra can be found in Supplementary Materials. Mass spectra were measured on Agilent 5975 GC-MS instrument (EI) from Agilent Technologies Co., Ltd. (Beijing, China). High-resolution mass spectra (ESI) were obtained with the Thermo Scientific LTQ Orbitrap XL mass spectrometer from Thermo Fisher Scientific (Shanghai, China). Melting points were measured with a YUHUA X-5 melting point instrument from Gongyi Yuhua Instrument Co., Ltd. (Gongyi, China) and were uncorrected. The structures of known compounds were further corroborated by comparing their ^1H NMR, ^{13}C NMR data and MS data with those of literature. All reagents were obtained from commercial suppliers and used without further purification unless otherwise noted.

4.2. General Procedure for the Synthesis of 2-Phenyl-9H-pyrimido[4,5-*b*]indole Derivatives

It was added to an oven-dried reaction vessel with a stir bar including indole-3-carboxaldehyde (0.2 mmol), benzaldehyde (0.4 mmol) and ammonium iodide (0.6 mmol), NaIO_4 (0.2 mmol), DMSO (0.4 mmol), I_2 (0.04 mmol), and 1,2-dichlorobenzene (0.6 mL). The reaction vessel was purged with oxygen for three times and then stirred at 150 °C for 16 h. After cooling to room temperature, the volatiles were removed under reduced

pressure. The residue was purified by column chromatography on neutral alumina to give the desired product **3aa** in 73% yield.

Characterization of products are below:

2-Phenyl-9*H*-pyrimido[4,5-*b*]indole (**3aa**, CAS: 1904604-60-9)

Yellow solid (37.3 mg, yield 76%, petroleum ether/ethyl acetate = 15:1), m.p. 295–297 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.35 (s, 1H), 9.53 (s, 1H), 8.51 (d, *J* = 6.1 Hz, 2H), 8.24 (d, *J* = 7.8 Hz, 1H), 7.60–7.48 (m, 5H), 7.33 (t, *J* = 7.3 Hz, 1H). ¹³C NMR (101 MHz, DMSO) δ 160.1, 156.5, 149.4, 139.5, 138.7, 130.7, 129.1, 128.2, 128.0, 122.1, 121.5, 119.5, 112.8, 112.3.

9-Methyl-2-phenyl-9*H*-pyrimido[4,5-*b*]indole (**3ba**):

Yellow solid (34.2 mg, yield 66%, petroleum ether/ethyl acetate = 15:1), m.p. 190–192 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.46 (s, 1H), 8.56 (d, *J* = 5.9 Hz, 2H), 8.22 (d, *J* = 7.8 Hz, 1H), 7.71–7.48 (m, 5H), 7.36 (t, *J* = 7.5 Hz, 1H), 3.93 (s, 3H). ¹³C NMR (101 MHz, DMSO) δ 160.0, 155.6, 149.1, 140.6, 138.5, 130.8, 129.0, 128.3, 128.0, 122.0, 121.8, 119.0, 112.5, 110.7, 27.9. HRMS (ESI): *m/z* calcd. for C₁₇H₁₄N₃ [M + H]⁺ 260.1182, found 260.1185.

5-Chloro-2-phenyl-9*H*-pyrimido[4,5-*b*]indole (**3ca**):

Yellow solid (53.7 mg, yield 96%, petroleum ether/ethyl acetate = 15:1), m.p. 320–322 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.70 (s, 1H), 9.57 (s, 1H), 8.50 (d, *J* = 7.1 Hz, 2H), 7.56–7.46 (m, 5H), 7.36 (d, *J* = 6.4 Hz, 1H). ¹³C NMR (101 MHz, DMSO) δ 160.6, 156.5, 150.3, 140.6, 138.3, 131.0, 129.1, 129.0, 128.4, 128.3, 121.6, 117.5, 111.8, 111.3. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁ClN₃ [M + H]⁺ 280.0636, found 280.0632.

5-Bromo-2-phenyl-9*H*-pyrimido[4,5-*b*]indole (**3da**):

Yellow solid (55.0 mg, yield 71%, petroleum ether/ethyl acetate = 15:1), m.p. 274–276 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.73 (s, 1H), 9.74 (s, 1H), 8.55–8.47 (m, 2H), 7.63–7.43 (m, 6H). ¹³C NMR (101 MHz, DMSO) δ 160.7, 156.5, 149.9, 140.6, 138.3, 131.0, 129.2, 129.1, 128.4, 124.8, 119.3, 116.6, 112.5, 111.7. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁BrN₃ [M + H]⁺ 324.0131, found 324.0160.

6-Methyl-2-phenyl-9*H*-pyrimido[4,5-*b*]indole (**3ea**):

Yellow solid (36.8 mg, yield 71%, petroleum ether/ethyl acetate = 15:1), m.p. 359–361 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.20 (s, 1H), 9.47 (s, 1H), 8.51 (d, *J* = 6.4 Hz, 2H), 8.02 (s, 1H), 7.58–7.43 (m, 4H), 7.38–7.31 (m, 1H), 2.47 (s, 3H). ¹³C NMR (101 MHz, DMSO) δ 160.0, 156.6, 149.2, 138.8, 137.6, 130.6, 130.4, 129.3, 129.0, 128.2, 121.8, 119.6, 112.7, 112.0, 21.5. HRMS (ESI): *m/z* calcd. for C₁₇H₁₄N₃ [M + H]⁺ 260.1182, found 260.1181.

6-Fluoro-2-phenyl-9*H*-pyrimido[4,5-*b*]indole (**3fa**):

Yellow solid (30.0 mg, yield 57%, petroleum ether/ethyl acetate = 15:1), m.p. 317–319 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.40 (s, 1H), 9.54 (s, 1H), 8.51 (d, *J* = 7.1 Hz, 2H), 8.10 (d, *J* = 8.1 Hz, 1H), 7.60–7.50 (m, 4H), 7.38 (t, *J* = 8.0 Hz, 1H). ¹³C NMR (101 MHz, DMSO-*d*₆) δ 160.6, 158.0 (d, *J* = 235.0 Hz), 157.1, 150.3, 138.6, 135.9, 130.8, 129.1, 128.3, 120.2 (d, *J* = 10.5 Hz), 115.8 (d, *J* = 25.6 Hz), 113.5 (d, *J* = 9.2 Hz), 112.7 (d, *J* = 4.5 Hz), 108.0 (d, *J* = 24.9 Hz). ¹⁹F NMR (376 MHz, DMSO) δ –121.9. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁FN₃ [M + H]⁺ 264.0932, found 264.0929.

6-Chloro-2-phenyl-9*H*-pyrimido[4,5-*b*]indole (**3ga**):

Yellow solid (45.7 mg, yield 82%, petroleum ether/ethyl acetate = 15:1), m.p. 290–292 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.52 (s, 1H), 9.58 (s, 1H), 8.51 (d, *J* = 7.9 Hz, 2H), 8.38 (s, 1H), 7.61–7.49 (m, 5H). ¹³C NMR (101 MHz, DMSO) δ 160.8, 156.9, 150.4, 138.5, 138.0, 130.9, 129.1, 128.3, 127.9, 125.8, 121.8, 120.9, 113.9, 112.2. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁ClN₃ [M + H]⁺ 280.0636, found 280.0632.

6-Bromo-2-phenyl-9*H*-pyrimido[4,5-*b*]indole (**3ha**):

Yellow solid (25.9 mg, yield 40%, petroleum ether/ethyl acetate = 15:1), m.p. 292–294 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.54 (s, 1H), 9.59 (s, 1H), 8.55–8.48 (m, 3H), 7.67 (d, *J* = 8.5 Hz, 1H), 7.54 (d, *J* = 6.9 Hz, 4H). ¹³C NMR (101 MHz, DMSO) δ 160.8, 156.7, 150.5, 138.5, 138.2, 130.9, 130.5, 129.1, 128.3, 124.8, 121.5, 114.3, 113.6, 112.0. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁BrN₃ [M + H]⁺ 324.0131, found 324.0155.

7-Fluoro-2-phenyl-9H-pyrimido[4,5-*b*]indole (3ia):

Yellow solid (37.9 mg, yield 72%, petroleum ether/ethyl acetate = 15:1), m.p. 286–288 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.50 (s, 1H), 9.52 (s, 1H), 8.50 (d, *J* = 6.4 Hz, 2H), 8.27 (dd, *J* = 8.6, 5.5 Hz, 1H), 7.59–7.48 (m, 3H), 7.34 (d, *J* = 7.3 Hz, 1H), 7.19 (t, *J* = 9.2 Hz, 1H). ¹³C NMR (101 MHz, DMSO-*d*₆) δ 162.4 (d, *J* = 242.0 Hz), 159.9, 157.2, 149.3, 140.4 (d, *J* = 12.5 Hz), 138.5, 130.7, 129.1, 128.2, 123.8 (d, *J* = 10.6 Hz), 116.2, 112.5, 109.6 (d, *J* = 24.2 Hz), 99.1 (d, *J* = 26.7 Hz). HRMS (ESI): *m/z* calcd. for C₁₆H₁₁FN₃ [M + H]⁺ 264.0932, found 264.0930.

7-Bromo-2-phenyl-9H-pyrimido[4,5-*b*]indole (3ja):

Yellow solid (27.6 mg, yield 43%, petroleum ether/ethyl acetate = 15:1), m.p. 291–293 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.52 (s, 1H), 9.57 (s, 1H), 8.50 (d, *J* = 8.1 Hz, 2H), 8.21 (d, *J* = 8.3 Hz, 1H), 7.72 (s, 1H), 7.57–7.46 (m, 4H). ¹³C NMR (101 MHz, DMSO) δ 160.6, 156.8, 150.0, 140.4, 138.5, 130.9, 129.1, 128.3, 124.4, 123.9, 120.7, 118.7, 115.0, 112.3. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁BrN₃ [M + H]⁺ 324.0131, found 324.0160.

7-Methoxy-2-phenyl-9H-pyrimido[4,5-*b*]indole (3ka):

Yellow solid (12.4 mg, yield 23%, petroleum ether/ethyl acetate = 10:1), m.p. 244–246 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.27 (s, 1H), 9.39 (s, 1H), 8.49 (d, *J* = 6.5 Hz, 2H), 8.10 (d, *J* = 8.6 Hz, 1H), 7.57–7.46 (m, 3H), 7.05–6.90 (m, 1H), 3.87 (s, 3H). ¹³C NMR (101 MHz, DMSO) δ 160.2, 158.9, 156.8, 147.9, 141.1, 138.8, 130.4, 129.0, 128.0, 123.0, 113.1, 112.8, 110.5, 95.9, 55.9. HRMS (ESI): *m/z* calcd. for C₁₇H₁₄N₃O [M + H]⁺ 276.1131, found 276.1137.

8-Chloro-2-phenyl-9H-pyrimido[4,5-*b*]indole (3la):

Yellow solid (42.9 mg, yield 77%, petroleum ether/ethyl acetate = 15:1), m.p. 247–249 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.81 (s, 1H), 9.57 (s, 1H), 8.51 (d, *J* = 7.1 Hz, 2H), 8.20 (d, *J* = 7.7 Hz, 1H), 7.61–7.50 (m, 4H), 7.31 (t, *J* = 7.8 Hz, 1H). ¹³C NMR (101 MHz, DMSO) δ 160.9, 156.9, 150.4, 138.5, 136.7, 130.9, 129.1, 128.3, 127.4, 122.5, 121.6, 120.8, 116.8, 112.9. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁ClN₃ [M + H]⁺ 280.0636, found 280.0632.

8-Methyl-2-phenyl-9H-pyrimido[4,5-*b*]indole (3ma):

Yellow solid (36.8 mg, yield 71%, petroleum ether/ethyl acetate = 15:1), m.p. 189–191 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.38 (s, 1H), 9.49 (s, 1H), 8.52 (d, *J* = 7.4 Hz, 2H), 8.02 (d, *J* = 7.7 Hz, 1H), 7.53 (q, *J* = 9.8, 8.5 Hz, 3H), 7.30 (d, *J* = 7.3 Hz, 1H), 7.21 (t, *J* = 7.5 Hz, 1H), 2.56 (s, 3H). ¹³C NMR (101 MHz, DMSO) δ 160.0, 156.7, 149.4, 138.8, 138.8, 130.6, 129.1, 128.6, 128.2, 121.9, 121.6, 119.3, 119.2, 113.2, 17.5. HRMS (ESI): *m/z* calcd. for C₁₇H₁₄N₃ [M + H]⁺ 260.1182, found 260.1181.

2-Phenyl-9H-pyrido[3',2':4,5]pyrrolo[2,3-*d*]pyrimidine (3na):

Yellow solid (29.0 mg, yield 59%, petroleum ether/ethyl acetate = 10:1), ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.92 (s, 1H), 9.57 (s, 1H), 8.63 (d, *J* = 7.7 Hz, 1H), 8.57–8.48 (m, 3H), 7.59–7.50 (m, 3H), 7.39 (dd, *J* = 7.8, 4.9 Hz, 1H). ¹³C NMR (101 MHz, DMSO) δ 160.8, 156.3, 152.0, 150.6, 148.3, 138.4, 131.0, 130.6, 129.1, 128.3, 117.9, 112.8, 111.3. HRMS (ESI): *m/z* calcd. for C₁₅H₁₁N₄ [M + H]⁺ 247.0978, found 247.0972.

2-(*p*-Tolyl)-9H-pyrimido[4,5-*b*]indole (3ab):

Yellow solid (39.4 mg, yield 76%, petroleum ether/ethyl acetate = 15:1), m.p. 277–279 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.29 (s, 1H), 9.49 (s, 1H), 8.40 (d, *J* = 7.9 Hz, 2H), 8.22 (d, *J* = 7.8 Hz, 1H), 7.61–7.45 (m, 2H), 7.36–7.27 (m, 3H), 2.38 (s, 3H). ¹³C NMR (101 MHz, DMSO) δ 160.3, 156.6, 149.4, 140.3, 139.4, 136.1, 129.7, 128.2, 127.9, 122.0, 121.4, 119.6, 112.6, 112.3, 21.5. HRMS (ESI): *m/z* calcd. for C₁₇H₁₄N₃ [M + H]⁺ 260.1182, found 260.1181.

2-(4-(*tert*-Butyl)phenyl)-9H-pyrimido[4,5-*b*]indole (3ac):

Yellow solid (47.6 mg, yield 79%, petroleum ether/ethyl acetate = 15:1), m.p. 171–173 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.36 (s, 1H), 9.50 (s, 1H), 8.43 (d, *J* = 8.5 Hz, 2H), 8.22 (d, *J* = 7.6 Hz, 1H), 7.58–7.49 (m, 4H), 7.36–7.28 (m, 1H), 1.33 (s, 9H). ¹³C NMR (101 MHz, DMSO) δ 160.2, 156.6, 153.4, 149.3, 139.4, 136.0, 128.1, 128.0, 125.9, 122.0, 121.4, 119.5, 112.6, 112.3, 35.0, 31.5. HRMS (ESI): *m/z* calcd. for C₂₀H₂₀N₃ [M + H]⁺ 302.1652, found 302.1635.

2-(4-Methoxyphenyl)-9H-pyrimido[4,5-*b*]indole (3ad):

White solid (40.7 mg, yield 74%, petroleum ether/ethyl acetate = 10:1), m.p. 256–258 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.35 (s, 1H), 9.47 (s, 1H), 8.45 (d, *J* = 8.9 Hz, 2H), 8.21 (d, *J* = 7.8 Hz, 1H), 7.56 (d, *J* = 8.0 Hz, 1H), 7.54–7.46 (m, 1H), 7.35–7.26 (m, 1H), 7.08 (d,

$J = 8.9$ Hz, 2H), 3.84 (s, 3H). ^{13}C NMR (101 MHz, DMSO) δ 161.5, 160.2, 156.6, 149.4, 139.3, 131.3, 129.8, 127.8, 121.8, 121.4, 119.6, 114.4, 112.3, 112.2, 55.8. HRMS (ESI): m/z calcd. for $\text{C}_{17}\text{H}_{14}\text{N}_3\text{O}$ $[\text{M} + \text{H}]^+$ 276.1131, found 276.1133.

2-(4-Ethoxyphenyl)-9H-pyrimido[4,5-*b*]indole (3ae):

Yellow solid (49.1 mg, yield 85%, petroleum ether/ethyl acetate = 10:1), m.p. 244–246 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 12.26 (s, 1H), 9.46 (s, 1H), 8.44 (d, $J = 8.9$ Hz, 2H), 8.20 (d, $J = 7.8$ Hz, 1H), 7.58–7.47 (m, 2H), 7.34–7.26 (m, 1H), 7.05 (d, $J = 8.9$ Hz, 2H), 4.09 (q, $J = 6.9$ Hz, 2H), 1.35 (t, $J = 6.9$ Hz, 3H). ^{13}C NMR (101 MHz, DMSO) δ 160.8, 160.2, 156.6, 149.4, 139.3, 131.1, 129.8, 127.8, 121.8, 121.4, 119.6, 114.8, 112.2, 112.2, 63.7, 15.1. HRMS (ESI): m/z calcd. for $\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}$ $[\text{M} + \text{H}]^+$ 290.1288, found 290.1289.

2-(4-Fluorophenyl)-9H-pyrimido[4,5-*b*]indole (3af):

Yellow solid (38.9 mg, yield 74%, petroleum ether/ethyl acetate = 15:1), m.p. 247–249 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 12.36 (s, 1H), 9.51 (s, 1H), 8.58–8.50 (m, 2H), 8.23 (d, $J = 7.8$ Hz, 1H), 7.60–7.48 (m, 2H), 7.40–7.28 (m, 3H). ^{13}C NMR (101 MHz, DMSO- d_6) δ 164.1 (d, $J = 247.4$ Hz), 159.2, 156.5, 149.4, 139.4, 135.2 (d, $J = 2.9$ Hz), 130.5 (d, $J = 8.6$ Hz), 128.1, 122.1, 121.5, 119.5, 116.0 (d, $J = 21.6$ Hz), 112.7, 112.3. ^{19}F NMR (376 MHz, DMSO) δ –111.5. HRMS (ESI): m/z calcd. for $\text{C}_{16}\text{H}_{11}\text{FN}_3$ $[\text{M} + \text{H}]^+$ 264.0932, found 264.0929.

2-(4-Chlorophenyl)-9H-pyrimido[4,5-*b*]indole (3ag):

Yellow solid (39.7 mg, yield 71%, petroleum ether/ethyl acetate = 15:1), m.p. 288–290 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 12.39 (s, 1H), 9.53 (s, 1H), 8.50 (d, $J = 8.6$ Hz, 2H), 8.24 (d, $J = 7.8$ Hz, 1H), 7.63–7.51 (m, 4H), 7.36–7.31 (m, 1H). ^{13}C NMR (101 MHz, DMSO) δ 159.1, 156.4, 149.4, 139.5, 137.6, 135.5, 129.9, 129.2, 128.2, 122.1, 121.6, 119.4, 113.0, 112.4. HRMS (ESI): m/z calcd. for $\text{C}_{16}\text{H}_{11}\text{ClN}_3$ $[\text{M} + \text{H}]^+$ 280.0636, found 280.0632.

2-(4-Bromophenyl)-9H-pyrimido[4,5-*b*]indole (3ah):

Yellow solid (42.8 mg, yield 66%, petroleum ether/ethyl acetate = 15:1), m.p. 284–286 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 12.37 (s, 1H), 9.52 (s, 1H), 8.43 (d, $J = 8.6$ Hz, 2H), 8.24 (d, $J = 7.8$ Hz, 1H), 7.73 (d, $J = 8.6$ Hz, 2H), 7.61–7.50 (m, 2H), 7.38–7.29 (m, 1H). ^{13}C NMR (101 MHz, DMSO) δ 159.2, 156.4, 149.4, 139.5, 137.9, 132.1, 130.2, 128.2, 124.4, 122.2, 121.6, 119.4, 113.0, 112.4. HRMS (ESI): m/z calcd. for $\text{C}_{16}\text{H}_{11}\text{BrN}_3$ $[\text{M} + \text{H}]^+$ 324.0131, found 324.0160.

2-(4-(Trifluoromethyl)phenyl)-9H-pyrimido[4,5-*b*]indole (3ai):

Yellow solid (48.3 mg, yield 78%, petroleum ether/ethyl acetate = 10:1), m.p. 290–292 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 12.48 (s, 1H), 9.58 (s, 1H), 8.69 (d, $J = 8.1$ Hz, 2H), 8.27 (d, $J = 7.8$ Hz, 1H), 7.90 (d, $J = 8.2$ Hz, 2H), 7.63–7.51 (m, 2H), 7.39–7.31 (m, 1H). ^{13}C NMR (101 MHz, DMSO- d_6) δ 158.6, 156.4, 149.5, 142.5, 139.7, 130.5 (q, $J = 31.6$ Hz), 128.8, 126.0 (q, $J = 4.0$ Hz), 124.8 (q, $J = 272.1$ Hz), 122.3, 121.7, 119.3, 113.4, 112.4. ^{19}F NMR (376 MHz, DMSO) δ –61.1. HRMS (ESI): m/z calcd. for $\text{C}_{17}\text{H}_{11}\text{F}_3\text{N}_3$ $[\text{M} + \text{H}]^+$ 314.0900, found 314.0930.

2-(4-(Trifluoromethyl)phenyl)-9H-pyrimido[4,5-*b*]indole (3aj):

Yellow solid (37.6 mg, yield 60%, petroleum ether/ethyl acetate = 10:1), m.p. 285–287 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 12.42 (s, 1H), 9.52 (s, 1H), 8.59 (d, $J = 8.9$ Hz, 2H), 8.24 (d, $J = 7.8$ Hz, 1H), 7.59–7.48 (m, 4H), 7.37–7.29 (m, 1H). ^{13}C NMR (101 MHz, DMSO- d_6) δ 158.8, 156.4, 150.3, 149.4, 139.5, 137.8, 130.2, 128.2, 122.2, 121.6, 121.4, 120.6 (q, $J = 256.8$ Hz), 119.4, 113.0, 112.4. ^{19}F NMR (376 MHz, CDCl_3) δ –51.9. HRMS (ESI): m/z calcd. for $\text{C}_{17}\text{H}_{11}\text{F}_3\text{N}_3\text{O}$ $[\text{M} + \text{H}]^+$ 330.0849, found 330.0876.

4-(9H-Pyrimido[4,5-*b*]indol-2-yl)benzotrile (3ak):

Yellow solid (27.0 mg, yield 50%, petroleum ether/ethyl acetate = 10:1), m.p. 298–300 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 12.47 (s, 1H), 9.56 (s, 1H), 8.62 (d, $J = 8.2$ Hz, 2H), 8.25 (d, $J = 7.8$ Hz, 1H), 7.98 (d, $J = 8.1$ Hz, 2H), 7.62–7.51 (m, 2H), 7.38–7.30 (m, 1H). ^{13}C NMR (101 MHz, DMSO) δ 158.2, 156.3, 149.4, 142.8, 139.7, 133.1, 128.7, 128.5, 122.3, 121.7, 119.3, 119.3, 113.5, 112.8, 112.4. HRMS (ESI): m/z calcd. for $\text{C}_{17}\text{H}_{11}\text{N}_4$ $[\text{M} + \text{H}]^+$ 271.0978, found 271.0976.

2-([1,1'-Biphenyl]-4-yl)-9H-pyrimido[4,5-*b*]indole (3al):

Yellow solid (36.6 mg, yield 57%, petroleum ether/ethyl acetate = 15:1), m.p. 298–300 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.38 (s, 1H), 9.55 (s, 1H), 8.60 (d, *J* = 8.4 Hz, 2H), 8.25 (d, *J* = 7.8 Hz, 1H), 7.85 (d, *J* = 8.2 Hz, 2H), 7.77 (d, *J* = 7.2 Hz, 2H), 7.60–7.47 (m, 4H), 7.41 (t, *J* = 7.4 Hz, 1H), 7.34 (t, *J* = 7.1 Hz, 1H). ¹³C NMR (101 MHz, DMSO) δ 159.9, 156.5, 149.5, 142.1, 140.0, 139.5, 137.8, 129.5, 128.8, 128.3, 128.1, 127.3, 127.2, 122.1, 121.5, 119.5, 112.8, 112.3. HRMS (ESI): *m/z* calcd. for C₂₂H₁₆N₃ [M + H]⁺ 322.1339, found 322.1325.

2-(*m*-Tolyl)-9H-pyrimido[4,5-*b*]indole (3am):

Yellow solid (39.4 mg, yield 76%, petroleum ether/ethyl acetate = 15:1), m.p. 247–249 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.35 (s, 1H), 9.51 (s, 1H), 8.34 (s, 1H), 8.30 (d, *J* = 7.8 Hz, 1H), 8.23 (d, *J* = 7.8 Hz, 1H), 7.60–7.48 (m, 2H), 7.41 (t, *J* = 7.6 Hz, 1H), 7.36–7.28 (m, 2H), 2.42 (s, 3H). ¹³C NMR (101 MHz, DMSO) δ 160.3, 156.5, 149.3, 139.4, 138.6, 138.1, 131.3, 129.0, 128.8, 128.0, 125.5, 122.0, 121.5, 119.5, 112.8, 112.3, 21.6. HRMS (ESI): *m/z* calcd. for C₁₇H₁₄N₃ [M + H]⁺ 260.1182, found 260.1185.

2-(3-Methoxyphenyl)-9H-pyrimido[4,5-*b*]indole (3an):

Yellow solid (50.0 mg, yield 91%, petroleum ether/ethyl acetate = 10:1), m.p. 265–267 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.34 (s, 1H), 9.51 (s, 1H), 8.23 (d, *J* = 7.8 Hz, 1H), 8.11 (d, *J* = 7.8 Hz, 1H), 8.06 (s, 1H), 7.60–7.49 (m, 2H), 7.45 (t, *J* = 7.9 Hz, 1H), 7.33 (t, *J* = 7.3 Hz, 1H), 7.13–7.04 (m, 1H), 3.87 (s, 3H). ¹³C NMR (101 MHz, DMSO) δ 160.0, 159.9, 156.5, 149.3, 140.2, 139.5, 130.1, 128.1, 122.1, 121.5, 120.7, 119.5, 116.7, 113.0, 112.9, 112.3, 55.6. HRMS (ESI): *m/z* calcd. for C₁₇H₁₄N₃O [M + H]⁺ 276.1131, found 276.1137.

2-(3-(Trifluoromethoxy)phenyl)-9H-pyrimido[4,5-*b*]indole (3ao):

Yellow solid (46.7 mg, yield 71%, petroleum ether/ethyl acetate = 10:1), m.p. 280–282 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.47 (s, 1H), 9.56 (s, 1H), 8.53 (d, *J* = 7.8 Hz, 1H), 8.39 (s, 1H), 8.26 (d, *J* = 7.8 Hz, 1H), 7.69 (t, *J* = 8.0 Hz, 1H), 7.61–7.48 (m, 3H), 7.35 (t, *J* = 7.2 Hz, 1H). ¹³C NMR (101 MHz, DMSO-*d*₆) δ 158.4, 156.4, 149.5, 149.3, 141.1, 139.6, 131.3, 128.4, 127.1, 123.1, 122.2, 121.6, 120.7 (q, *J* = 256.4 Hz), 119.9, 119.4, 113.4, 112.4. ¹⁹F NMR (376 MHz, DMSO) δ –56.6. HRMS (ESI): *m/z* calcd. for C₁₇H₁₁F₃N₃O [M + H]⁺ 330.0849, found 330.0844.

2-(3-Fluorophenyl)-9H-pyrimido[4,5-*b*]indole (3ap):

Yellow solid (36.3 mg, yield 69%, petroleum ether/ethyl acetate = 15:1), m.p. 210–212 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.53 (s, 1H), 9.54 (s, 1H), 8.35 (d, *J* = 7.8 Hz, 1H), 8.25 (d, *J* = 7.8 Hz, 1H), 8.22–8.15 (m, 1H), 7.64–7.49 (m, 3H), 7.39–7.29 (m, 2H). ¹³C NMR (101 MHz, DMSO-*d*₆) δ 163.01 (d, *J* = 242.5 Hz), 158.77 (d, *J* = 3.3 Hz), 156.32, 149.39, 141.32 (d, *J* = 7.8 Hz), 139.64, 131.15 (d, *J* = 8.3 Hz), 128.25, 124.20 (d, *J* = 2.6 Hz), 122.18, 121.56, 119.36, 117.41 (d, *J* = 21.1 Hz), 114.47 (d, *J* = 23.1 Hz), 113.20, 112.47. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁FN₃ [M + H]⁺ 264.0932, found 264.0930.

2-(3-Chlorophenyl)-9H-pyrimido[4,5-*b*]indole (3aq):

Yellow solid (31.3 mg, yield 56%, petroleum ether/ethyl acetate = 15:1), m.p. 201–203 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.49 (s, 1H), 9.55 (s, 1H), 8.48 (s, 1H), 8.47–8.42 (m, 1H), 8.25 (d, *J* = 7.8 Hz, 1H), 7.63–7.50 (m, 4H), 7.38–7.30 (m, 1H). ¹³C NMR (101 MHz, DMSO) δ 158.6, 156.4, 149.4, 140.8, 139.6, 134.0, 131.1, 130.4, 128.3, 127.8, 126.7, 122.2, 121.6, 119.4, 113.3, 112.4. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁ClN₃ [M + H]⁺ 280.0636, found 280.0632.

2-(3-Bromophenyl)-9H-pyrimido[4,5-*b*]indole (3ar):

Yellow solid (38.9 mg, yield 60%, petroleum ether/ethyl acetate = 15:1), m.p. 213–215 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.54 (s, 1H), 9.54 (s, 1H), 8.63 (s, 1H), 8.49 (d, *J* = 7.8 Hz, 1H), 8.25 (d, *J* = 7.8 Hz, 1H), 7.70 (d, *J* = 8.0 Hz, 1H), 7.61 (d, *J* = 8.1 Hz, 1H), 7.58–7.47 (m, 2H), 7.33 (t, *J* = 7.5 Hz, 1H). ¹³C NMR (101 MHz, DMSO) δ 158.4, 156.3, 149.4, 141.0, 139.6, 133.2, 131.4, 130.7, 128.3, 127.1, 122.5, 122.2, 121.6, 119.4, 113.2, 112.5. HRMS (ESI): *m/z* calcd. for C₁₆H₁₁BrN₃ [M + H]⁺ 324.0131, found 324.0160.

2-(*o*-Tolyl)-9H-pyrimido[4,5-*b*]indole (3as):

Yellow solid (28.5 mg, yield 55%, petroleum ether/ethyl acetate = 15:1), m.p. 235–237 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.04 (s, 1H), 9.73 (s, 1H), 8.36 (d, *J* = 7.8 Hz, 1H), 7.79 (d, *J* = 7.2 Hz, 1H), 7.71–7.61 (m, 2H), 7.52–7.37 (m, 4H), 2.53 (s, 3H). ¹³C NMR (101 MHz,

DMSO- d_6) δ 156.2, 140.2, 137.5, 131.7, 131.0, 130.7, 129.5, 126.5, 122.9, 122.8, 119.4, 113.1, 112.4, 20.8. HRMS (ESI): m/z calcd. for $C_{17}H_{14}N_3$ $[M + H]^+$ 260.1182, found 260.1181.

2-(2-Fluorophenyl)-9H-pyrimido[4,5-*b*]indole (3at):

Yellow solid (30.0 mg, yield 57%, petroleum ether/ethyl acetate = 15:1), m.p. 271–273 °C. 1H NMR (400 MHz, DMSO- d_6) δ 12.46 (s, 1H), 9.57 (s, 1H), 8.27 (d, J = 7.8 Hz, 1H), 8.11 (t, J = 7.9 Hz, 1H), 7.61–7.49 (m, 3H), 7.40–7.31 (m, 3H). ^{13}C NMR (101 MHz, DMSO- d_6) δ 162.1, 159.6, 158.7 (d, J = 4.3 Hz), 149.3, 139.4, 132.5 (d, J = 2.1 Hz), 131.9 (d, J = 8.5 Hz), 128.3, 127.7 (d, J = 9.7 Hz), 124.8 (d, J = 3.7 Hz), 122.3, 121.5, 119.2, 117.2 (d, J = 22.3 Hz), 112.6, 112.4. ^{19}F NMR (376 MHz, DMSO) δ –114.4. HRMS (ESI): m/z calcd. for $C_{16}H_{11}FN_3$ $[M + H]^+$ 264.0932, found 264.0929.

2-(2-Chlorophenyl)-9H-pyrimido[4,5-*b*]indole (3au):

Yellow solid (30.0 mg, yield 43%, petroleum ether/ethyl acetate = 15:1), m.p. 261–263 °C. 1H NMR (400 MHz, DMSO- d_6) δ 12.43 (s, 1H), 9.57 (s, 1H), 8.28 (d, J = 7.5 Hz, 1H), 7.82–7.76 (m, 1H), 7.63–7.45 (m, 5H), 7.37 (t, J = 6.1 Hz, 1H). ^{13}C NMR (101 MHz, DMSO) δ 161.0, 155.9, 149.1, 139.4, 139.2, 132.4, 132.1, 130.7, 130.6, 128.3, 127.5, 122.3, 121.6, 119.2, 112.6, 112.4. HRMS (ESI): m/z calcd. for $C_{16}H_{11}ClN_3$ $[M + H]^+$ 280.0636, found 280.0632.

2-(Naphthalen-2-yl)-9H-pyrimido[4,5-*b*]indole (3av):

Yellow solid (41.2 mg, yield 70%, petroleum ether/ethyl acetate = 15:1), m.p. 304–306 °C. 1H NMR (400 MHz, DMSO- d_6) δ 12.35 (s, 1H), 9.56 (s, 1H), 9.06 (s, 1H), 8.63 (d, J = 8.6 Hz, 1H), 8.25 (d, J = 7.8 Hz, 1H), 8.12–8.07 (m, 1H), 8.05 (d, J = 8.7 Hz, 1H), 8.00–7.94 (m, 1H), 7.62–7.51 (m, 4H), 7.33 (t, J = 6.9 Hz, 1H). ^{13}C NMR (101 MHz, DMSO) δ 160.1, 156.6, 149.4, 139.5, 136.1, 134.4, 133.4, 129.4, 128.5, 128.1, 128.1, 127.6, 127.0, 125.5, 122.1, 121.6, 119.5, 112.9, 112.4. HRMS (ESI): m/z calcd. for $C_{20}H_{14}N_3$ $[M + H]^+$ 296.1182, found 296.1170.

2-(2,4-Dichlorophenyl)-9H-pyrimido[4,5-*b*]indole (3aw):

Yellow solid (29.5 mg, yield 47%, petroleum ether/ethyl acetate = 15:1), m.p. 212–214 °C. 1H NMR (400 MHz, DMSO- d_6) δ 12.49 (s, 1H), 9.58 (s, 1H), 8.29 (d, J = 7.8 Hz, 1H), 7.84 (d, J = 8.4 Hz, 1H), 7.77 (s, 1H), 7.63–7.55 (m, 3H), 7.37 (t, J = 6.4 Hz, 1H). ^{13}C NMR (101 MHz, DMSO) δ 160.0, 155.8, 149.1, 139.5, 138.0, 134.5, 133.8, 133.2, 130.1, 128.5, 127.8, 122.4, 121.7, 119.1, 112.8, 112.4. HRMS (ESI): m/z calcd. for $C_{16}H_{10}Cl_2N_3$ $[M + H]^+$ 314.0246, found 314.0273.

2-(3,4-Dimethylphenyl)-9H-pyrimido[4,5-*b*]indole (3ax):

Yellow solid (35.0 mg, yield 64%, petroleum ether/ethyl acetate = 15:1), m.p. 299–301 °C. 1H NMR (400 MHz, DMSO- d_6) δ 12.30 (s, 1H), 9.49 (s, 1H), 8.30 (s, 1H), 8.22 (d, J = 7.9 Hz, 2H), 7.58–7.47 (m, 2H), 7.35–7.25 (m, 2H), 2.34 (s, 3H), 2.30 (s, 3H). ^{13}C NMR (101 MHz, DMSO) δ 160.4, 156.6, 149.3, 139.4, 139.1, 136.7, 136.4, 130.2, 129.3, 127.9, 125.8, 121.9, 121.4, 119.6, 112.5, 112.3, 20.1, 19.9. HRMS (ESI): m/z calcd. for $C_{18}H_{16}N_3$ $[M + H]^+$ 274.1339, found 274.1338.

2-(Thiophen-3-yl)-9H-pyrimido[4,5-*b*]indole (3ay):

Yellow solid (22.6 mg, yield 45%, petroleum ether/ethyl acetate = 15:1), m.p. 283–285 °C. 1H NMR (400 MHz, DMSO- d_6) δ 12.33 (s, 1H), 9.46 (s, 1H), 8.39 (s, 1H), 8.22 (d, J = 7.8 Hz, 1H), 7.91 (d, J = 5.0 Hz, 1H), 7.70–7.63 (m, 1H), 7.58–7.47 (m, 2H), 7.32 (t, J = 6.4 Hz, 1H). ^{13}C NMR (101 MHz, DMSO) δ 157.8, 156.4, 149.5, 142.7, 139.3, 127.9, 127.7, 127.7, 127.4, 121.9, 121.5, 119.6, 112.4, 112.3. HRMS (ESI): m/z calcd. for $C_{14}H_{10}N_3S$ $[M + H]^+$ 252.0590, found 252.0588.

2-(Thiophen-2-yl)-9H-pyrimido[4,5-*b*]indole (3az):

Yellow solid (13.6 mg, yield 27%, petroleum ether/ethyl acetate = 15:1), m.p. 272–274 °C. 1H NMR (400 MHz, DMSO- d_6) δ 12.38 (s, 1H), 9.42 (s, 1H), 8.21 (d, J = 7.8 Hz, 1H), 8.02–7.96 (m, 1H), 7.73 (d, J = 4.1 Hz, 1H), 7.57–7.47 (m, 2H), 7.36–7.28 (m, 1H), 7.25–7.18 (m, 1H). ^{13}C NMR (101 MHz, DMSO) δ 157.2, 156.2, 149.4, 144.7, 139.3, 130.3, 128.9, 128.4, 128.0, 121.9, 121.6, 119.6, 112.5, 112.3. HRMS (ESI): m/z calcd. for $C_{14}H_{10}N_3S$ $[M + H]^+$ 252.0590, found 252.0588.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/catal13030623/s1>, ^1H NMR, ^{13}C NMR and ^{19}F NMR spectra of products **3aa–3az**.

Author Contributions: Conceptualization, Y.C., F.X. and G.-J.D.; Methodology, Y.C.; data curation, Y.C., R.Y. and T.L.; validation, Y.C., R.Y., F.X., T.L. and G.-J.D.; investigation, Y.C.; writing—original draft preparation, Y.C.; writing—review and editing, Y.C., F.X. and G.-J.D.; supervision, F.X. and G.-J.D.; funding acquisition, G.M. and G.-J.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Natural Science Foundation of China (22271244, 21871226), the Undergraduate Investigated Study and Innovated Experiment Plan from Ministry of Education of China and Hunan Province, and the Open Research Fund of School of Chemistry and Chemical Engineering, Henan Normal University (2022C02).

Data Availability Statement: The authors declare that the raw data supporting the findings and conclusions of this research are available from the corresponding author upon request.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Okasha, R.M.; Albalawi, F.F.; Afifi, T.H.; Fouda, A.M.; Al-Dies, A.-A.M.; El-Agrody, A.M. Structural Characterization and Antimicrobial Activities of 7H-Benzo[h]chromeno[2,3-d]pyrimidine and 14H-Benzo[h]chromeno[3,2-e][1,2,4]triazolo[1,5-c] pyrimidine Derivatives. *Molecules* **2016**, *21*, 1450. [[CrossRef](#)] [[PubMed](#)]
2. De Coen, L.M.; Heugebaert, T.S.A.; Garcia, D.; Stevens, C.V. Synthetic entries to and biological activity of pyrrolopyrimidines. *Chem. Rev.* **2016**, *116*, 80–139. [[CrossRef](#)] [[PubMed](#)]
3. Chauhan, R.; Siddiqi, A.A.; Dwivedi, J. Synthesis of pyrimidino derivatives: Potent antimicrobial candidates. *Pharm. Chem. J.* **2017**, *50*, 678–683. [[CrossRef](#)]
4. Guillena, G.; Ramon, D.J.; Yus, M. Hydrogen autotransfer in the N-alkylation of amines and related compounds using alcohols and amines as electrophiles. *Chem. Rev.* **2010**, *110*, 1611–1641. [[CrossRef](#)]
5. Walker, S.R.; Carter, E.J.; Huff, B.C.; Morris, J.C. Variolins and related alkaloids. *Chem. Rev.* **2009**, *109*, 3080–3098. [[CrossRef](#)]
6. Parker, W.B. Enzymology of purine and pyrimidine antimetabolites used in the treatment of cancer. *Chem. Rev.* **2009**, *109*, 2880–2893. [[CrossRef](#)]
7. Feng, Y.; Xie, X.Y.; Yang, Y.Q.; Sun, Y.T.; Ma, W.H.; Zhou, P.J.; Li, Z.Y.; Liu, H.Q.; Wang, Y.F.; Huang, Y.S. Synthesis and evaluation of pyrimidindole analogs in umbilical cord blood ex vivo expansion. *Eur. J. Med. Chem.* **2019**, *174*, 181–197. [[CrossRef](#)]
8. Akaki, T.; Bessho, Y.; Ito, T.; Fujioka, S.; Ubukata, M.; Mori, G.; Yamanaka, K.; Orita, T.; Doi, S.; Iwanaga, T.; et al. Fragment-based lead discovery to identify novel inhibitors that target the ATP binding site of pyruvate dehydrogenase kinases. *Bioorg. Med. Chem.* **2021**, *44*, 116283. [[CrossRef](#)]
9. Hollis Showalter, H.D.; Bridges, A.J.; Zhou, H.; Sercel, A.D.; McMichael, A.; Fry, D.W. Tyrosine Kinase Inhibitors. 16. 6,5,6-Tricyclic Benzothieno[3,2-d]pyrimidines and Pyrimido[5,4-b]- and -[4,5-b]indoles as Potent Inhibitors of the Epidermal Growth Factor Receptor Tyrosine Kinase. *J. Med. Chem.* **1999**, *42*, 5464–5474. [[CrossRef](#)]
10. Eschke, D.; Brand, A.; Scheibler, P.; Hess, S.; Eger, K.; Allgaier, C.; Nieber, K. Effect of an adenosine A1 receptor agonist and a novel pyrimidoindole on membrane properties and neurotransmitter release in rat cortical and hippocampal neurons, *Neurochemistry International*. *Neurochem. Int.* **2001**, *38*, 391–398. [[CrossRef](#)]
11. Hurter, J.; Ramp, T.; Patrian, B.; Städler, E.; Roessingh, P.; Baur, R.; De Jong, R.; Nielsen, J.K.; Winkler, T.; Richter, W.J.; et al. Oviposition stimulants for the cabbage root fly: Isolation from cabbage leaves. *Phytochemistry* **1999**, *51*, 377–382. [[CrossRef](#)]
12. Portonov, Y.N.; Bulaga, S.N.; Zabrodnyaya, V.G.; Smirnov, L.D. Synthesis of 9H-pyrimido[4,5-b]indoles from 2-amino-3-cyanoindoles. *Chem. Heterocycl. Compd.* **1991**, *27*, 325–327. [[CrossRef](#)]
13. Li, B.; Guo, S.; Zhang, J.; Zhang, X.; Fan, X. Selective Access to 3-Cyano-1H-indoles, 9H-Pyrimido[4,5-b]indoles, or 9H-Pyrido[2,3-b]indoles through Copper-Catalyzed One-Pot Multicomponent Cascade Reactions. *J. Org. Chem.* **2015**, *80*, 5444–5456. [[CrossRef](#)] [[PubMed](#)]
14. Adib, M.; Mohammadi, B.; Bijanzadeh, H.R. A Novel and Simple Synthesis of 9H-Pyrimido[4,5-b]indoles under Microwave Irradiation and Solvent-Free Conditions. *Synlett* **2008**, *2008*, 177–180. [[CrossRef](#)]
15. Jafari, B.; Safarov, S.; Khalikova, M.; Ehlers, P.; Langer, P. Synthesis of 1,3,4-Thiadiazolo[2',3':2,3]imidazo[4,5-b]indoles. *Synlett* **2019**, *30*, 1791–1794. [[CrossRef](#)]
16. Hyatt, J.A.; Swenton, J.S. A facile synthesis of 9H-pyrimido [4,5-b]indole via photolysis of 4-azido-5-phenylpyrimidine. *J. Heterocycl. Chem.* **1972**, *9*, 409–410. [[CrossRef](#)]
17. Xu, G.; Zheng, L.; Wang, S.; Wang, Q.; Bai, X. Regiospecific Syntheses of 3-Aza- α -carboline via Inverse Electron-Demand Diels-Alder Reactions of 2-Aminoindoles with 1,3,5-Triazines. *Synlett* **2009**, *2009*, 3206–3210. [[CrossRef](#)]

18. Gao, W.; Wang, D.; Li, Y. A simple and facile synthesis of tricyclic-fused pyrimido[4,5-b]indol-2-amines. *Mol. Divers.* **2017**, *21*, 831–840. [[CrossRef](#)]
19. Qu, Z.; Wang, P.; Chen, X.; Deng, G.-J.; Huang, H. Visible-light-driven Cadogan reaction. *Chin. Chem. Lett.* **2021**, *32*, 2582–2586. [[CrossRef](#)]
20. Borovik, V.P.; Gatilov, Y.V.; Shkurko, O.P. Reaction of 3-(R-methylidene)-2-ethoxylindolenines with N,N'-binucleophiles. *Russ. J. Org. Chem.* **2016**, *52*, 228–234. [[CrossRef](#)]
21. Burange, A.S.; Gopinath, C.S. Catalytic applications of hydrotalcite and related materials in multi-component reactions: Concepts, challenges and future scope. *Sustain. Chem. Pharm.* **2021**, *22*, 100458. [[CrossRef](#)]
22. Vachan, B.S.; Karuppasamy, M.; Vinoth, P.; Vivek Kumar, S.; Perumal, S.; Sridharan, V.; Menéndez, J.C. Proline and its Derivatives as Organocatalysts for Multi-Component Reactions in Aqueous Media: Synergic Pathways to the Green Synthesis of Heterocycles. *Adv. Synth. Catal.* **2020**, *362*, 87. [[CrossRef](#)]
23. Zhang, L.-L.; Zhang, J.-W.; Xiang, S.-H.; Guo, Z.; Tan, B. Stereoselective Construction of Complex Spirooxindoles via Bisthiourea Catalyzed Three-Component Reactions. *Chin. J. Chem.* **2018**, *36*, 1182–1186. [[CrossRef](#)]
24. Ren, C.; Wang, Z.; Wang, T.; Guo, J.; Dai, Y.; Yuan, H.; Tan, Y. Ultralong Organic Phosphorescence Modulation of Aromatic Carbonyls and Multi-Component Systems. *Chin. J. Chem.* **2022**, *40*, 1987–2000. [[CrossRef](#)]
25. Khan, I.; Zaib, S.; Ibrar, A. New frontiers in the transition-metal-free synthesis of heterocycles from alkynoates: An overview and current status. *Org. Chem. Front.* **2020**, *7*, 3734–3791. [[CrossRef](#)]
26. Guo, H.; Tian, L.; Liu, Y.; Wan, J.-P. DMSO as a C1 Source for [2 + 2 + 1] Pyrazole Ring Construction via Metal-Free Annulation with Enaminones and Hydrazines. *Org. Lett.* **2022**, *24*, 228–233. [[CrossRef](#)]
27. Liu, H.; Cai, Z.-J.; Ji, S.-J. Access to 3-alkylselenindoles by multicomponent reaction of indoles, selenium powder and unactivated alkyl halides under transition-metal-free conditions. *Chin. Chem. Lett.* **2022**, *33*, 4303–4305. [[CrossRef](#)]
28. Xu, Z.; Fu, L.; Fang, X.; Huang, B.; Zhou, L.; Wan, J.-P. Tunable Trifunctionalization of Tertiary Enaminones for the Regioselective and Metal-Free Synthesis of Discrete and Proximal Phosphoryl Nitriles. *Org. Lett.* **2021**, *23*, 5049–5053. [[CrossRef](#)]
29. Han, C.; Nie, L.; Han, X.; Zhang, Y.; Sun, K.; Shi, L.; Cui, G.; Meng, W. One-Pot Three-Component Synthesis of Novel 1,5-Benzodiazepine Derivatives and Their anti-BVDV (Bovine Viral Diarrhea Virus) Activity. *Chin. J. Org. Chem.* **2021**, *41*, 819–825. [[CrossRef](#)]
30. Khalili, B.; Jajarmi, P.; Eftekhari-Sis, B.; Hashemi, M.M. Novel One-Pot, Three-Component Synthesis of New 2-Alkyl-5-aryl-(1H)-pyrrole-4-ol in Water. *J. Org. Chem.* **2008**, *73*, 2090–2095. [[CrossRef](#)]
31. Kumar, D.; Kommi, D.N.; Bollineni, N.; Patel, A.R.; Chakraborti, A.K. Catalytic procedures for multicomponent synthesis of imidazoles: Selectivity control during the competitive formation of tri- and tetrasubstituted imidazoles. *Green Chem.* **2012**, *14*, 2038–2049. [[CrossRef](#)]
32. Xiang, S.; Fan, W.; Zhang, W.; Li, Y.; Guo, S.; Huang, D. Aqueous CO₂ fixation: Construction of pyridine skeletons in cooperation with ammonium cations. *Green Chem.* **2021**, *23*, 7950–7955. [[CrossRef](#)]
33. Li, J.; Li, J.; He, R.; Liu, J.; Liu, Y.; Chen, L.; Huang, Y.; Li, Y. Selective Synthesis of Substituted Pyridines and Pyrimidines through Cascade Annulation of Isopropene Derivatives. *Org. Lett.* **2022**, *24*, 1620–1625. [[CrossRef](#)] [[PubMed](#)]
34. Shukla, G.; Srivastava, A.; Singh, M.S. Metal- and Catalyst-Free, Formal [4 + 1] Annulation via Tandem C=O/C=S Functionalization: One-Pot Access to 3,5-Disubstituted/Annulated Isothiazoles. *Org. Lett.* **2016**, *18*, 2451–2454. [[CrossRef](#)] [[PubMed](#)]
35. Kumar, D.; Sharma, H.; Saha, N.; Chakraborti, A.K. Domino synthesis of functionalized pyridine carboxylates under gallium catalysis: Unravelling the reaction pathway and the role of the nitrogen source counter anion. *Chem. Asian J.* **2022**, *17*, e202200304. [[CrossRef](#)]
36. Yaremenko, I.A.; Belyakova, Y.Y.; Radulov, P.S.; Novikov, R.A.; Medvedev, M.G.; Krivoschapov, N.V.; Korlyukov, A.A.; Alabugin, I.V.; Terent'ev, A.O. Inverse α -Effect as the Ariadne's Thread on the Way to Tricyclic Aminoperoxides: Avoiding Thermodynamic Traps in the Labyrinth of Possibilities. *J. Am. Chem. Soc.* **2022**, *144*, 7264–7282. [[CrossRef](#)]
37. Sasada, T.; Kobayashi, F.; Sakai, N.; Konakahara, T. An Unprecedented Approach to 4,5-Disubstituted Pyrimidine Derivatives by a ZnCl₂-Catalyzed Three-Component Coupling Reaction. *Org. Lett.* **2009**, *11*, 2161–2164. [[CrossRef](#)]
38. Chen, J.; Meng, H.; Zhang, F.; Xiao, F.; Deng, G.-J. Transition-metal-free selective pyrimidines and pyridines formation from aromatic ketones, aldehydes and ammonium salts. *Green Chem.* **2019**, *21*, 5201–5206. [[CrossRef](#)]
39. Jiang, J.; Huang, H.; Deng, G.J. Four-component thiazole formation from simple chemicals under metal-free conditions. *Green Chem.* **2019**, *21*, 986–990. [[CrossRef](#)]
40. Bai, Y.; Tang, L.; Huang, H.; Deng, G.-J. Synthesis of 2,4-diarylsubstituted-pyridines through a Ru-catalyzed four component reaction. *Org. Biomol. Chem.* **2015**, *13*, 4404–4407. [[CrossRef](#)]
41. Chen, J.; Chang, D.; Xiao, F.; Deng, G.-J. Four-component quinazoline synthesis from simple anilines, aromatic aldehydes and ammonium iodide under metal-free conditions. *Green Chem.* **2018**, *20*, 5459–5463. [[CrossRef](#)]
42. Chen, J.; Sun, Z.; Xiao, F.; Deng, G.-J. Base-promoted aerobic oxidative synthesis of fused 1,3,5-triazines under metal-free conditions. *Green Chem.* **2020**, *22*, 6778–6782. [[CrossRef](#)]
43. Wang, P.; Zhang, X.; Liu, Y.; Chen, B. Synthesis of Pyrimidines with Ammonium Acetate as Nitrogen Source Under Solvent-Free Conditions. *Asian J. Org. Chem.* **2019**, *8*, 1122. [[CrossRef](#)]

44. Duan, T.T.; Zhai, T.R.; Liu, H.H.; Yan, Z.L.; Zhao, Y.; Feng, L.; Ma, C. One-pot three-component synthesis of quinazolines via a copper-catalysed oxidative amination reaction. *Org. Biomol. Chem.* **2016**, *14*, 6561–6567. [[CrossRef](#)]
45. Chen, Z.Y.; Chen, J.X.; Liu, M.C.; Ding, J.C.; Gao, W.X.; Huang, X.B.; Wu, H.Y. Unexpected Copper-Catalyzed Cascade Synthesis of Quinazoline Derivatives. *J. Org. Chem.* **2013**, *78*, 11342–11348. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.