

Supplementary material

Synthesis of Optically and Redox Active Polyenaminones from Diamines and α,α' -Bis[(dimethylamino)methyldene]cyclohexanediones.

Urša Štanfel ¹, Tomaž Kotnik ¹, Sebastijan Ričko ¹, Uroš Grošelj ¹, Bogdan Štefane ¹, Klemen Pirnat ³, Ema Žagar ², Boštjan Genorio ^{1*} and Jurij Svete ^{1,*}

¹ University of Ljubljana, Faculty of Chemistry and Chemical Technology, Večna pot 113, 1000 Ljubljana, Slovenia


² National Institute of Chemistry, Department of Materials Chemistry, Hajdrihova 19, 1000 Ljubljana, Slovenia



³ National Institute of Chemistry, Department of Polymer Chemistry and Technology, Hajdrihova 19, 1000 Ljubljana, Slovenia



Table of contents


1. Copies of ¹ H NMR and ¹³ C NMR Spectra of Compounds 2 and 4 .	2–4
2. Copies of IR spectra of compounds 2 and 4 .	5–13
3. Copies of UV-VIS spectra of compounds 4aa–4af and 4ba–4bf .	9–14





1. Copies of ^1H NMR and ^{13}C NMR Spectra of Compounds 2 and 4.





















3. Copies of UV-VIS spectra of compounds 4aa–4af and 4ba–4bf.

