

Correction



Correction: Almeida-Dalmet, S.; et al. Differential Gene Expression in Response to Salinity and Temperature in a *Haloarcula* Strain from Great Salt Lake, Utah. *Genes* 2017, 9, 52

Swati Almeida-Dalmet¹, Carol D. Litchfield¹, Patrick Gillevet² and Bonnie K. Baxter^{3,*}

- ¹ Department of Environmental Science and Policy, George Mason University, 10900 University Blvd, Manassas, VA 20110, USA; swati_almeida2002@yahoo.com
- ² Department of Biology, George Mason University, 10900 University Blvd, Manassas, VA 20110, USA; pgilleve@gmu.edu
- ³ Great Salt Lake Institute, Westminster College, 1840 South 1300 East, Salt Lake City, UT 84105, USA
- * Correspondence: bbaxter@westminstercollege.edu; Tel.: +1-801-832-2345

Received: 23 February 2018; Accepted: 2 March 2018; Published: 7 March 2018

The authors wish to make the following changes to their paper [1]. In the original published version of Figure 2 ("Confirmation of differential gene expression ... "), many of the photographic panels were missing. This figure has been replaced with a complete version (Figure 1). In addition, the position and numbering of Figures 2 and 3 have been exchanged to reflect the order in which they are mentioned in the text, and the references to these figures have been updated in the text.

We apologize for any inconvenience caused to the readers by this change. The manuscript will be updated and the original will remain online on the article webpage.

(A) Salinity 15% 20% 27%		(B) Temperature 12°C 37°C 42°C
-	Transcriptional regulator	46 - 16E
-	<i>hEF-Tu</i> , haloarchaea EF	No Data
	<i>hcp</i> , halocyanin	No Data
	ATPase in partitioning	100 100 000
	ABC transporter	Re 18.10
1000	ftsZ, cell division protein	
	Phenyl acetyl CoA ligase	
	Phosphoserine phosphatase	
10 (C.)	Ser/Thr protein Kinase	18 102 102
60 100	Amino acid permease	

Figure 1. Confirmation of differential gene expression of RAP-PCR fragments in NA6-27 grown under the following conditions. (**A**) Salinity 15, 20, and 27% NaCl (w/v) and (**B**) Temperature 12, 37 and 42 °C. Quantitative PCR was performed using gene specific primers as described by Benson et al. [2–4]. Gel documentation photographs are aligned with data on genes for each growth condition.

References

- Almeida-Dalmet, S.; Litchfield, C.D.; Gillevet, P.; Baxter, B.K. Differential Gene Expression in Response to Salinity and Temperature in a Haloarcula Strain from Great Salt Lake, Utah. *Genes* 2018, 9, 52. [CrossRef] [PubMed]
- 2. Bidle, K.A. Differential expression of genes influenced by changing salinity using RNA arbitrarily primed PCR in the archaeal halophile Haloferax volcanii. *Extremophiles* **2003**, *7*, 1–7. [PubMed]
- 3. Bidle, K.A.; Bartlett, D.H. RNA arbitrarily primed PCR survey of genes regulated by ToxR in the deep-sea bacterium Photobacterium profundum str. SS9. *J. Bacteriol.* **2001**, *183*, 1688–1693. [CrossRef] [PubMed]
- Benson, N.R.; Wong, R.M.-Y.; McClelland, M. Analysis of the SOS response in *Salmonella enterica* serovar typhimurium using RNA fingerprinting by arbitrarily primed PCR. *J. Bacteriol.* 2000, 182, 3490–3497. [CrossRef] [PubMed]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).