



## Article

# Initial Coefficient Estimates and Fekete–Szegő Inequalities for New Families of Bi-Univalent Functions Governed by $(p - q)$ -Wanas Operator

Abbas Kareem Wanas <sup>1,†</sup>  and Luminița-Ioana Cotîrlă <sup>2,\*,†</sup> 

<sup>1</sup> Department of Mathematics, College of Science, University of Al-Qadisiyah, Al-Qadisiyah 58001, Iraq; abbas.kareem.w@qu.edu.iq

<sup>2</sup> Department of Mathematics, Technical University of Cluj-Napoca, 400114 Cluj-Napoca, Romania

\* Correspondence: luminita.cotirla@math.utcluj.ro or luminita.cotirla@yahoo.com

† These authors contributed equally to this work.

**Abstract:** The motivation of the present article is to define the  $(p - q)$ -Wanas operator in geometric function theory by the symmetric nature of quantum calculus. We also initiate and explore certain new families of holomorphic and bi-univalent functions  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  and  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  which are defined in the unit disk  $U$  associated with the  $(p - q)$ -Wanas operator. The upper bounds for the initial Taylor–Maclaurin coefficients and Fekete–Szegő-type inequalities for the functions in these families are obtained. Furthermore, several consequences of our results are pointed out based on the various special choices of the involved parameters.

**Keywords:** holomorphic function; bi-univalent function; upper bounds; Fekete–Szegő functional;  $(p - q)$ -Wanas operator

**MSC:** 30C45; 30C20



**Citation:** Wanas, A.K.; Cotîrlă, L.-I. Initial Coefficient Estimates and Fekete–Szegő Inequalities for New Families of Bi-Univalent Functions Governed by  $(p - q)$ -Wanas Operator. *Symmetry* **2021**, *13*, 2118. <https://doi.org/10.3390/sym13112118>

Academic Editor: Alina Alb Lupas

Received: 1 October 2021

Accepted: 1 November 2021

Published: 8 November 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Indicate by  $\mathcal{A}$  the family of all holomorphic functions of the form

$$f(z) = z + \sum_{k=2}^{\infty} b_k z^k \quad (1)$$

in the open unit disk  $U = \{z \in \mathbb{C} : |z| < 1\}$ .

We also denote by  $T$  the subfamily of  $\mathcal{A}$  consisting of functions which are also univalent in  $U$ .

The famous Koebe one-quarter theorem [1] ensures that the image of  $U$  under each univalent function  $f \in \mathcal{A}$  contains a disk of radius  $\frac{1}{4}$ . Furthermore, each function  $f \in T$  has an inverse  $f^{-1}$  defined by  $f^{-1}(f(z)) = z$  and

$$f(f^{-1}(w)) = w, \quad \left(|w| < r_0(f), r_0(f) \geq \frac{1}{4}\right)$$

where

$$g(w) = f^{-1}(w) = w - b_2 w^2 + (2b_2^2 - b_3)w^3 - (5b_2^3 - 5b_2 b_3 + b_4)w^4 + \dots$$

A function  $f \in \mathcal{A}$  is named bi-univalent in  $U$  if both  $f$  and  $f^{-1}$  are univalent in  $U$ . The family of all bi-univalent functions in  $U$  is denoted by  $E$ .

From the work of Srivastava et al. [2], we choose to recall the following examples of functions in the family  $E$ :

$$\frac{z}{1-z}, \quad -\log(1-z) \quad \text{and} \quad \frac{1}{2} \log\left(\frac{1+z}{1-z}\right).$$

In fact, Srivastava et al. [2] have actually revived the study of analytic and bi-univalent functions in the recent years. This was followed by works such as those by Frasin and Aouf [3], Ali et al. [4], Bulut et al. [5], Srivastava and et al. [6] and others (see, for example, [7–15]).

We notice that the family  $E$  is not empty. However, the Koebe function is not a member of  $E$ .

The problem to obtain the general coefficient bounds on the Taylor–Maclaurin coefficients

$$|b_n| \quad (n \in \mathbb{N}; n \geq 3)$$

for functions  $f \in E$  is still not completely addressed for many of the subfamilies of  $E$ . The origin of the Fekete-Szegő functional  $|b_3 - \eta b_2^2|$  for  $f \in T$  was in the disproof [16] by Fekete and Szegő of the Littlewood-Paley conjecture that the coefficients of odd univalent functions are bounded by unity. Researchers in the Theory of Geometric Function have recently obtained remarkable results on this topic (see, for example, [17–21]).

With a view to recalling the principle of subordination between holomorphic functions, let the functions  $f$  and  $g$  be holomorphic in  $U$ . The function  $f$  is subordinate to  $g$ , if there exists a Schwarz function  $\omega$ , which is analytic in  $U$  with

$$\omega(0) = 0 \quad \text{and} \quad |\omega(z)| < 1 \quad (z \in U),$$

such that

$$f(z) = g(\omega(z)).$$

The subordination is denoted by

$$f \prec g \quad \text{or} \quad f(z) \prec g(z) \quad (z \in U).$$

It is well known that (see [22]), if the function  $g$  is univalent in  $U$ , then

$$f \prec g \quad (z \in U) \iff f(0) = g(0) \quad \text{and} \quad f(U) \subseteq g(U).$$

For  $0 < q < p \leq 1$ , the  $(p, q)$ -derivative operator or  $(p, q)$ -difference operator for a function  $f$  is defined by

$$D_{p,q}f(z) = \frac{f(pz) - f(qz)}{(p-q)z} \quad (z \in U^* = U \setminus \{0\}), \quad (2)$$

and

$$D_{p,q}f(0) = f'(0).$$

For more details on the concepts of  $(p, q)$ -calculus see [20,23–27].

For function  $f \in \mathcal{A}$ , we deduce that

$$D_{p,q}f(z) = 1 + \sum_{k=2}^{\infty} [k]_{p,q} b_k z^{k-1}, \quad (3)$$

where the  $(p, q)$ -bracket number or twin-basic  $[k]_{p,q}$  is given by

$$[k]_{p,q} = \frac{p^k - q^k}{p - q} = p^{k-1} + p^{k-2}q + p^{k-3}q^2 + \cdots + pq^{k-2} + q^{k-1} \quad (p \neq q),$$

which is a natural generalization of the  $q$ -number, that is, we have (see [28–30])

$$\lim_{p \rightarrow 1^-} [k]_{p,q} = [k]_q = \frac{1 - q^k}{1 - q}.$$

It is clear that the notation  $[k]_{p,q}$  is symmetric, that is,

$$[k]_{p,q} = [k]_{q,p}.$$

In 2019, Wanas [31] introduced the following operator, which can also be called Wanas operator  $W_{s,t}^{\sigma,\delta} : \mathcal{A} \rightarrow \mathcal{A}$  defined by

$$W_{s,t}^{\sigma,\delta} f = z + \sum_{k=2}^{\infty} \left[ \sum_{\tau=1}^{\sigma} \binom{\sigma}{\tau} (-1)^{\tau+1} \left( \frac{s^{\tau} + kt^{\tau}}{s^{\tau} + t^{\tau}} \right) \right]^{\delta} b_k z^k, \quad (4)$$

where  $s \in \mathbb{R}$ ,  $t \in \mathbb{R}_0^+$  with  $s + t > 0$ ,  $k - 1, \sigma \in \mathbb{N}$  and  $\delta \in \mathbb{N}_0$ .

Now, for  $f \in \mathcal{A}$ , we define the  $(p - q)$ -difference Wanas operator as given below

$$\begin{aligned} W_{0,1,p,q}^{1,0} f(z) &= f(z) \\ W_{0,1,p,q}^{1,1} f(z) &= z W_{p,q} f(z) \\ W_{0,1,p,q}^{1,\delta} f(z) &= z W_{p,q} (W_{p,q}^{\delta-1} f(z)) \\ W_{s,t,p,q}^{\sigma,\delta} f(z) &= z + \sum_{k=2}^{\infty} \left( \frac{[\Psi_k(\sigma, s, t)]_{p,q}}{[\Psi_1(\sigma, s, t)]_{p,q}} \right)^{\delta} b_k z^k \end{aligned}$$

where

$$\begin{aligned} W_{p,q} f(z) &= z + \sum_{k=2}^{\infty} \frac{p^k - q^k}{p - q} b_k z^k, \quad W_{p,q}^{\delta-1} f(z) = z + \sum_{k=2}^{\infty} \left( \frac{p^k - q^k}{p - q} \right)^{\delta-1} b_k z^k, \\ \Psi_k(\sigma, s, t) &= \sum_{\tau=1}^{\sigma} \binom{\sigma}{\tau} (-1)^{\tau+1} (s^{\tau} + kt^{\tau}), \quad \Psi_1(\sigma, s, t) = \sum_{\tau=1}^{\sigma} \binom{\sigma}{\tau} (-1)^{\tau+1} (s^{\tau} + t^{\tau}), \\ s &\in \mathbb{R}, t \in \mathbb{R}_0^+ \text{ with } s + t > 0, k - 1, \sigma \in \mathbb{N}, \delta \in \mathbb{N}_0, 0 < q < p \leq 1 \text{ and } z \in U. \end{aligned}$$

**Remark 1.** The operator  $W_{s,t,p,q}^{\sigma,\delta}$  is a generalization of several known operators studied in earlier investigations which are being recalled below.

1. For  $p = \sigma = t = 1$ ,  $\delta = -\nu$ ,  $\Re(\nu) > 1$  and  $s \in \mathbb{C} \setminus \mathbb{Z}_0^-$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the  $q$ -Srivastava–Attiya operator  $J_{q,s}^{\nu}$  [32];
2. For  $p = \sigma = t = 1$ ,  $\delta = -1$  and  $s > -1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the  $q$ -Bernardi operator [33];
3. For  $p = \sigma = s = t = 1$  and  $\delta = -1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the  $q$ -Libera operator [33];
4. For  $s = 0$  and  $p = \sigma = t = 1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the  $q$ -Sălăgean operator [34];
5. For  $q \rightarrow 1^-$  and  $p = \sigma = 1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the operator  $I_{s,t}^{\delta}$  that was introduced and studied by Swamy [35];
6. For  $q \rightarrow 1^-$ ,  $p = \sigma = t = 1$ ,  $\delta = -\nu$ ,  $\Re(\nu) > 1$  and  $s \in \mathbb{C} \setminus \mathbb{Z}_0^-$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the operator  $J_s^{\nu}$  that was investigated by Srivastava and Attiya [36]. The operator  $J_s^{\nu}$  is now popularly known in the literature as the Srivastava–Attiya operator;
7. For  $q \rightarrow 1^-$ ,  $p = \sigma = t = 1$  and  $s > -1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the operator  $I_s^{\delta}$  that was investigated by Cho and Srivastava [37];

8. For  $q \rightarrow 1^-$ ,  $p = \sigma = s = t = 1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the operator  $I^\delta$  that was considered by Uralegaddi and Somanatha [38];
9. For  $q \rightarrow 1^-$ ,  $p = \sigma = s = t = 1$ ,  $\delta = -\xi$  and  $\xi > 0$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the operator  $I^\xi$  that was introduced by Jung et al. [39]. The operator  $I^\xi$  is the Jung–Kim–Srivastava integral operator;
10. For  $q \rightarrow 1^-$ ,  $p = \sigma = t = 1$ ,  $\delta = -1$  and  $s > -1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the Bernardi operator [40];
11. For  $q \rightarrow 1^-$ ,  $s = 0$ ,  $p = \sigma = t = 1$  and  $\delta = -1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the Alexander operator [41];
12. For  $q \rightarrow 1^-$ ,  $p = \sigma = 1$ ,  $s = 1 - t$  and  $t \geq 0$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the operator  $D_t^\delta$  that was given by Al-Oboudi [42];
13. For  $q \rightarrow 1^-$ ,  $p = \sigma = 1$ ,  $s = 0$  and  $t = 1$ , the operator  $W_{s,t,p,q}^{\sigma,\delta}$  reduces to the operator  $S^\delta$  that was considered by Sălăgean [43].

We shall need the following Lemma in our investigation.

**Lemma 1** ([44], p. 41 and [45], p. 41). Let the function  $x \in P$  be given by the following series:

$$x(z) = 1 + x_1 z + x_2 z^2 + \cdots \quad (z \in U).$$

The sharp estimate given by

$$|x_n| \leq 2 \quad (n \in \mathbb{N})$$

holds true.

## 2. A Set of Main Results

Indicate by  $\vartheta(z)$  the holomorphic function with positive real part in  $U$  such that

$$\vartheta(0) = 1, \quad \vartheta'(0) > 0$$

and  $\vartheta(U)$  is symmetric with respect to the real axis, which is of the type:

$$\vartheta(z) = 1 + B_1 z + B_2 z^2 + B_3 z^3 + \cdots \quad (5)$$

where  $B_1 > 0$ .

Using the  $(p - q)$ -Wanas operator, we now provide the following subfamilies of holomorphic and bi-univalent functions.

**Definition 1.** For  $0 \leq \lambda \leq 1$ , a function  $f \in E$  is said to be in the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  if it fulfills the subordinations:

$$(1 - \lambda) \frac{z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'}{W_{s,t,p,q}^{\sigma,\delta} f(z)} + \lambda \left( 1 + \frac{z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)''}{\left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'} \right) \prec \vartheta(z)$$

and

$$(1 - \lambda) \frac{w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'}{W_{s,t,p,q}^{\sigma,\delta} g(w)} + \lambda \left( 1 + \frac{w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)''}{\left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'} \right) \prec \vartheta(w),$$

where  $g(w) = f^{-1}(w)$ .

**Definition 2.** For  $\mu \geq 1$  and  $\gamma \geq 0$ , a function  $f \in E$  is said to be in the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  if it fulfills the subordinations:

$$(1 - \mu) \frac{W_{s,t,p,q}^{\sigma,\delta} f(z)}{z} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)' + \gamma z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'' \prec \vartheta(z)$$

and

$$(1 - \mu) \frac{W_{s,t,p,q}^{\sigma,\delta} g(w)}{w} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)' + \gamma w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'' \prec \vartheta(w),$$

where  $g(w) = f^{-1}(w)$ .

In particular, if we choose

$$\vartheta(z) = \left( \frac{1+z}{1-z} \right)^\alpha \quad (0 < \alpha \leq 1) \quad \text{and} \quad \vartheta(z) = \frac{1+(1-2\beta)z}{1-z} \quad (0 \leq \beta < 1),$$

the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the families  $H_E(\lambda, \sigma, \delta, s, t, p, q; \alpha)$  and  $H_E(\lambda, \sigma, \delta, s, t, p, q; \beta)$  which are families of the functions  $f \in E$  satisfying

$$\left| \arg \left( (1 - \lambda) \frac{z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'}{W_{s,t,p,q}^{\sigma,\delta} f(z)} + \lambda \left( 1 + \frac{z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)''}{\left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'} \right) \right) \right| < \frac{\alpha\pi}{2},$$

$$\left| \arg \left( (1 - \lambda) \frac{w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'}{W_{s,t,p,q}^{\sigma,\delta} g(w)} + \lambda \left( 1 + \frac{w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)''}{\left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'} \right) \right) \right| < \frac{\alpha\pi}{2}$$

and

$$\Re \left( (1 - \lambda) \frac{z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'}{W_{s,t,p,q}^{\sigma,\delta} f(z)} + \lambda \left( 1 + \frac{z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)''}{\left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'} \right) \right) > \beta,$$

$$\Re \left( (1 - \lambda) \frac{w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'}{W_{s,t,p,q}^{\sigma,\delta} g(w)} + \lambda \left( 1 + \frac{w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)''}{\left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'} \right) \right) > \beta,$$

respectively.

In addition, the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the families  $T_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \alpha)$  and  $T_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \beta)$  which are families of the functions  $f \in E$  satisfying

$$\left| \arg \left( (1 - \mu) \frac{W_{s,t,p,q}^{\sigma,\delta} f(z)}{z} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)' + \gamma z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'' \right) \right| < \frac{\alpha\pi}{2},$$

$$\left| \arg \left( (1 - \mu) \frac{W_{s,t,p,q}^{\sigma,\delta} g(w)}{w} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)' + \gamma w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'' \right) \right| < \frac{\alpha\pi}{2}$$

and

$$\Re \left( (1 - \mu) \frac{W_{s,t,p,q}^{\sigma,\delta} f(z)}{z} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)' + \gamma z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'' \right) > \beta,$$

$$\Re \left( (1 - \mu) \frac{W_{s,t,p,q}^{\sigma,\delta} g(w)}{w} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)' + \gamma w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'' \right) > \beta,$$

respectively.

**Remark 2.** The families  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  and  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  are a generalization of several known families studied in earlier investigations which are being recalled below.

1. For  $\delta = 0$  and  $\vartheta(z) = \frac{a+(b-ap_1)rz}{1-p_1rz-q_1z^2} + 1 - a$ ,  $r \in \mathbb{R}$ ,  $a, b, p_1$  and  $q_1$  are real constants, the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{M}_E(\lambda, r)$  which was studied by Abirami et al. [1];
2. For  $\delta = 0$ ,  $\lambda = 1$  and  $\vartheta(z) = \frac{a+(b-ap_1)rz}{1-p_1rz-q_1z^2} + 1 - a$ ,  $r \in \mathbb{R}$ ,  $a, b, p_1$  and  $q_1$  are real constants, the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{K}_E(r)$  which was introduced by Abirami et al. [1];
3. For  $\delta = \gamma = 0$ ,  $\mu = 1$  and  $\vartheta(z) = \left(\frac{1+z}{1-z}\right)^\alpha$ ,  $0 < \alpha \leq 1$ , the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{H}_E(\alpha)$  which was investigated by Srivastava et al. [2];
4. For  $\delta = \gamma = 0$ ,  $\mu = 1$  and  $\vartheta(z) = \frac{1+(1-2\beta)z}{1-z}$ ,  $0 \leq \beta < 1$ , the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{H}_E(\beta)$  which was defined by Srivastava et al. [2].
5. For  $\delta = \gamma = 0$  and  $\vartheta(z) = \left(\frac{1+z}{1-z}\right)^\alpha$ ,  $0 < \alpha \leq 1$ , the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{B}_E(\alpha, \mu)$  which was considered by Frasin and Aouf [3];
6. For  $\delta = \gamma = 0$  and  $\vartheta(z) = \frac{1+(1-2\beta)z}{1-z}$ ,  $0 \leq \beta < 1$ , the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{B}_E(\beta, \mu)$  which was studied by Frasin and Aouf [3];
7. For  $\delta = 0$ , the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{M}_E(\lambda; \vartheta)$  which was introduced by Ali et al. [4];
8. For  $\delta = \lambda = 0$  and  $\vartheta(z) = \frac{1}{1-2tz+z^2}$ ,  $t \in (\frac{1}{2}, 1]$ , the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{S}_E^*(t)$  which was introduced by Bulut et al. [5];
9. For  $\delta = \lambda = 0$  and  $\vartheta(z) = \frac{a+(b-ap_1)rz}{1-p_1rz-q_1z^2} + 1 - a$ ,  $r \in \mathbb{R}$ ,  $a, b, p_1$  and  $q_1$  are real constants, the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{W}_E(r)$  which was defined by Srivastava et al. [10];
10. For  $\delta = 0$  and  $\vartheta(z) = \left(\frac{1+z}{1-z}\right)^\alpha$ ,  $0 < \alpha \leq 1$ , the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{M}_E(\alpha, \lambda)$  which was considered by Liu and Wang [46];
11. For  $\delta = 0$  and  $\vartheta(z) = \frac{1+(1-2\beta)z}{1-z}$ ,  $0 \leq \beta < 1$ , the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{M}_E(\lambda)$  which was studied by Liu and Wang [46];
12. For  $\delta = \lambda = 0$  and  $\vartheta(z) = \left(\frac{1+z}{1-z}\right)^\alpha$ ,  $0 < \alpha \leq 1$ , the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{S}_E^*(\alpha)$  which was considered by Brannan and Taha [47];
13. For  $\delta = \lambda = 0$  and  $\vartheta(z) = \frac{1+(1-2\beta)z}{1-z}$ ,  $0 \leq \beta < 1$ , the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{S}_E^*(\beta)$  which was investigated by Brannan and Taha [47];
14. For  $\delta = 0$  and  $\vartheta(z) = \frac{1}{1-2tz+z^2}$ ,  $t \in (\frac{\sqrt{2}}{2}, 1]$ , the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{H}_E(\lambda, t)$  which was studied by Altinkaya and Yalçın [48];
15. For  $\delta = 0$ ,  $\mu = 1$  and  $\vartheta(z) = \left(\frac{1+z}{1-z}\right)^\alpha$ ,  $0 < \alpha \leq 1$ , the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{H}_E(\alpha, \gamma)$  which was considered by Frasin [49];
16. For  $\delta = 0$ ,  $\mu = 1$  and  $\vartheta(z) = \frac{1+(1-2\beta)z}{1-z}$ ,  $0 \leq \beta < 1$ , the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{H}_E(\gamma, \beta)$  which was studied by Frasin [49];
17. For  $\delta = 0$  and  $\vartheta(z) = \frac{1+(1-2\beta)z}{1-z}$ ,  $0 \leq \beta < 1$ , the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  reduces to the family  $\mathcal{N}_E(\beta, \mu, \gamma)$  which was defined by Bulut [50];

**Theorem 1.** Let  $f$  given by (1) be in the family  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$ . Then,

$$|b_2| \leq \min \left\{ \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(\lambda+1)[\Psi_2(\sigma, s, t)]_{p,q}^\delta}, \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta B_1^{\frac{3}{2}}}{\sqrt{2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} \left( (\lambda+1)^2(B_1 - B_2) - (3\lambda+1)B_1^2 \right)}} \right\}$$

and

$$|b_3| \leq \min \left\{ \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{B_2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{2(2\lambda + 1)[1 + t]_q^\delta [\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta - (3\lambda + 1)[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}, \right. \\ \left. \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{(\lambda + 1)^2[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}} \right\},$$

where the coefficients  $B_1$  and  $B_2$  are defined as in (5).

**Proof.** Let  $f \in \mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  and  $g = f^{-1}$ . Then, there are holomorphic functions  $S, T : U \rightarrow U$  with  $S(0) = T(0) = 0$ , which fulfill the following conditions:

$$(1 - \lambda) \frac{z(W_{s,t,p,q}^{\sigma,\delta} f(z))'}{W_{s,t,p,q}^{\sigma,\delta} f(z)} + \lambda \left( 1 + \frac{z(W_{s,t,p,q}^{\sigma,\delta} f(z))''}{(W_{s,t,p,q}^{\sigma,\delta} f(z))'} \right) = \vartheta(S(z)), \quad z \in U \quad (6)$$

and

$$(1 - \lambda) \frac{w(W_{s,t,p,q}^{\sigma,\delta} g(w))'}{W_{s,t,p,q}^{\sigma,\delta} g(w)} + \lambda \left( 1 + \frac{w(W_{s,t,p,q}^{\sigma,\delta} g(w))''}{(W_{s,t,p,q}^{\sigma,\delta} g(w))'} \right) = \vartheta(T(w)), \quad w \in U. \quad (7)$$

Define the functions  $x$  and  $y$  by

$$x(z) = \frac{1 + S(z)}{1 - S(z)} = 1 + x_1 z + x_2 z^2 + \dots$$

and

$$y(z) = \frac{1 + T(z)}{1 - T(z)} = 1 + y_1 z + y_2 z^2 + \dots$$

Then,  $x$  and  $y$  are analytic in  $U$  with  $x(0) = y(0) = 1$ . Since we have  $S, T : U \rightarrow U$ , each of the functions  $x$  and  $y$  has a positive real part in  $U$ .

Solving for  $S(z)$  and  $T(z)$ , we have

$$S(z) = \frac{x(z) - 1}{x(z) + 1} = \frac{1}{2} \left[ x_1 z + \left( x_2 - \frac{x_1^2}{2} \right) z^2 \right] + \dots \quad (z \in U) \quad (8)$$

and

$$T(z) = \frac{y(z) - 1}{y(z) + 1} = \frac{1}{2} \left[ y_1 z + \left( y_2 - \frac{y_1^2}{2} \right) z^2 \right] + \dots \quad (z \in U). \quad (9)$$

By substituting (8) and (9) into (6) and (7) and applying (5), we obtain

$$(1 - \lambda) \frac{z(W_{s,t,p,q}^{\sigma,\delta} f(z))'}{W_{s,t,p,q}^{\sigma,\delta} f(z)} + \lambda \left( 1 + \frac{z(W_{s,t,p,q}^{\sigma,\delta} f(z))''}{(W_{s,t,p,q}^{\sigma,\delta} f(z))'} \right) \\ = \vartheta \left( \frac{x(z) - 1}{x(z) + 1} \right) = 1 + \frac{1}{2} B_1 x_1 z + \left[ \frac{1}{2} B_1 \left( x_2 - \frac{x_1^2}{2} \right) + \frac{1}{4} B_2 x_1^2 \right] z^2 + \dots \quad (10)$$

and

$$\begin{aligned} (1-\lambda) \frac{w \left( W_{s,t,p,q}^{\sigma,\delta} \mathcal{G}(w) \right)'}{W_{s,t,p,q}^{\sigma,\delta} \mathcal{G}(w)} + \lambda \left( 1 + \frac{w \left( W_{s,t,p,q}^{\sigma,\delta} \mathcal{G}(w) \right)''}{\left( W_{s,t,p,q}^{\sigma,\delta} \mathcal{G}(w) \right)'} \right) \\ = \vartheta \left( \frac{y(w)-1}{y(w)+1} \right) = 1 + \frac{1}{2} B_1 y_1 w + \left[ \frac{1}{2} B_1 \left( y_2 - \frac{y_1^2}{2} \right) + \frac{1}{4} B_2 y_1^2 \right] w^2 + \dots \quad (11) \end{aligned}$$

Equating the coefficients in (10) and (11), yields

$$\frac{(\lambda+1)[\Psi_2(\sigma, s, t)]_{p,q}^{\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{\delta}} b_2 = \frac{1}{2} B_1 x_1, \quad (12)$$

$$\frac{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^{\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{\delta}} b_3 - \frac{(3\lambda+1)[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}} b_2^2 = \frac{1}{2} B_1 \left( x_2 - \frac{x_1^2}{2} \right) + \frac{1}{4} B_2 x_1^2, \quad (13)$$

$$- \frac{(\lambda+1)[\Psi_2(\sigma, s, t)]_{p,q}^{\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{\delta}} b_2 = \frac{1}{2} B_1 y_1 \quad (14)$$

and

$$\frac{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^{\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{\delta}} (2b_2^2 - b_3) - \frac{(3\lambda+1)[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}} b_2^2 = \frac{1}{2} B_1 \left( y_2 - \frac{y_1^2}{2} \right) + \frac{1}{4} B_2 y_1^2. \quad (15)$$

From (12) and (14), we have

$$x_1 = -y_1 \quad (16)$$

and

$$\frac{2(\lambda+1)^2 [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}} b_2^2 = \frac{1}{4} B_1^2 (x_1^2 + y_1^2). \quad (17)$$

If we add (13) to (15), we obtain

$$\begin{aligned} \frac{4(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^{\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{\delta}} b_2^2 - \frac{2(3\lambda+1)[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}} b_2^2 \\ = \frac{1}{2} B_1 \left[ x_2 + y_2 - \left( \frac{x_1^2 + y_1^2}{2} \right) \right] + \frac{1}{4} B_2 [x_1^2 + y_1^2] \quad (18) \end{aligned}$$

Substituting the value of  $x_1^2 + y_1^2$  from (17) in the right-hand side of (18), we deduce that

$$b_2^2 = \frac{B_1^3 [\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} (x_2 + y_2)}{4 \left[ 2B_1^2 (2\lambda+1) [\Psi_1(\sigma, s, t)]_{p,q}^{\delta} [\Psi_3(\sigma, s, t)]_{p,q}^{\delta} + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} \left( (\lambda+1)^2 (B_1 - B_2) - (3\lambda+1) B_1^2 \right) \right]}. \quad (19)$$

Applying Lemma 1 for the coefficients  $x_1, x_2, y_1, y_2$  in (17) and (19), we obtain

$$\begin{aligned} |b_2| &\leq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^{\delta} B_1^{\frac{3}{2}}}{\sqrt{\left| 2B_1^2 (2\lambda+1) [\Psi_1(\sigma, s, t)]_{p,q}^{\delta} [\Psi_3(\sigma, s, t)]_{p,q}^{\delta} + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} \left( (\lambda+1)^2 (B_1 - B_2) - (3\lambda+1) B_1^2 \right) \right|}}, \\ |b_2| &\leq \frac{B_1 [\Psi_1(\sigma, s, t)]_{p,q}^{\delta}}{(\lambda+1) [\Psi_2(\sigma, s, t)]_{p,q}^{\delta}}, \end{aligned}$$

which gives the estimates of  $|b_2|$ .



Furthermore, in order to find the bound on  $|b_3|$ , we subtract (15) from (13) and also apply (16). We obtain  $x_1^2 = y_1^2$ , hence,

$$\frac{4(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta} (b_3 - b_2^2) = \frac{1}{2} B_1(x_2 - y_2), \quad (20)$$

then, by substituting the value of  $b_2^2$  from (17) into (20), gives

$$b_3 = \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta (x_2 - y_2)}{8(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} (x_1^2 + y_1^2)}{8(\lambda + 1)^2 [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}.$$

So, we have

$$|b_3| \leq \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{(\lambda + 1)^2 [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}.$$

In addition, substituting the value of  $b_2^2$  from (18) into (20), we obtain

$$b_3 = \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta (x_2 - y_2)}{8(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{(B_1(x_2 + y_2) + \frac{1}{2}(x_1^2 + y_1^2)(B_2 - B_1))[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{8(2\lambda + 1)[1 + t]_q^\delta [\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta - 4(3\lambda + 1)[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}$$

and we have

$$|b_3| \leq \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{B_2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{2(2\lambda + 1)[1 + t]_q^\delta [\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta - (3\lambda + 1)[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}},$$

which gives us the desired estimates of the coefficient  $|b_3|$ .  $\square$

Taking  $\vartheta(z) = \left(\frac{1+z}{1-z}\right)^\alpha = 1 + 2\alpha z + 2\alpha^2 z^2 + \dots$  ( $0 < \alpha \leq 1$ ) in Theorem 1, we obtain the next corollary.

**Corollary 1.** Let  $f$  given by (1) be in the family  $H_E(\lambda, \sigma, \delta, s, t, p, q; \alpha)$ , where ( $0 < \alpha \leq 1$ ). Then,

$$|b_2| \leq \min \left\{ \frac{2\alpha[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(\lambda + 1)[\Psi_2(\sigma, s, t)]_{p,q}^\delta}, \frac{2\alpha\sqrt{2\alpha}[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{\sqrt{|4\alpha^2(2\lambda + 1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (2\alpha(1 - \alpha)(\lambda + 1)^2 - 4\alpha^2(3\lambda + 1))|}} \right\}$$

and

$$|b_3| \leq \min \left\{ \frac{\alpha[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{2\alpha^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{2(2\lambda + 1)[1 + t]_q^\delta [\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta - (3\lambda + 1)[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}, \frac{\alpha[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(2\lambda + 1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{4\alpha^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{(\lambda + 1)^2 [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}} \right\}.$$

Taking  $\vartheta(z) = \frac{1+(1-2\beta)z}{1-z} = 1 + 2(1 - \beta)z + 2(1 - \beta)z^2 + \dots$  ( $0 \leq \beta < 1$ ) in Theorem 1, we obtain the next corollary.

**Corollary 2.** Let the function  $f$  given by (1) be in the function family  $H_E(\lambda, \sigma, \delta, s, t, p, q; \beta)$ , where ( $0 \leq \beta < 1$ ). Then,

$$|b_2| \leq \min \left\{ \frac{2(1 - \beta)[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(\lambda + 1)[\Psi_2(\sigma, s, t)]_{p,q}^\delta}, \frac{\sqrt{2(1 - \beta)}[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{\sqrt{|2(2\lambda + 1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta - [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (3\lambda + 1)|}} \right\}$$

and

$$|b_3| \leq \min \left\{ \frac{(1-\beta)[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{2(1-\beta)[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{2(2\lambda+1)[1+t]_q^\delta [\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta - (3\lambda+1)[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}, \right. \\ \left. \frac{(1-\beta)[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{4(1-\beta)^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{(\lambda+1)^2[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}} \right\}.$$

**Theorem 2.** Let  $f$  given by (1) be in the family  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$ . Then,

$$|b_2| \leq \min \left\{ \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+\mu+2\gamma)[\Psi_2(\sigma, s, t)]_{p,q}^\delta}, \right. \\ \left. \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta B_1^{\frac{3}{2}}}{\sqrt{B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}(1+\mu+2\gamma)^2(B_1-B_2)}} \right\}$$

and

$$|b_3| \leq \min \left\{ \frac{B_2[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}, \right. \\ \left. \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{(1+\mu+2\gamma)^2[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}} \right\},$$

where the coefficients  $B_1$  and  $B_2$  are defined as in (5).

**Proof.** Let  $f \in \mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  and  $g = f^{-1}$ . Then, there are holomorphic functions  $S, T : U \rightarrow U$  such that

$$(1-\mu) \frac{W_{s,t,p,q}^{\sigma,\delta} f(z)}{z} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)' + \gamma z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'' = \vartheta(S(z)), \quad z \in U \quad (21)$$

and

$$(1-\mu) \frac{W_{s,t,p,q}^{\sigma,\delta} g(w)}{w} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)' + \gamma w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'' = \vartheta(T(w)), \quad w \in U, \quad (22)$$

where  $S(z)$  and  $T(z)$  have the forms (8) and (9). From (21), (22) and (5), we deduce that

$$(1-\mu) \frac{W_{s,t,p,q}^{\sigma,\delta} f(z)}{z} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)' + \gamma z \left( W_{s,t,p,q}^{\sigma,\delta} f(z) \right)'' \\ = \vartheta \left( \frac{x(z)-1}{x(z)+1} \right) = 1 + \frac{1}{2} B_1 x_1 z + \left[ \frac{1}{2} B_1 \left( x_2 - \frac{x_1^2}{2} \right) + \frac{1}{4} B_2 x_1^2 \right] z^2 + \dots \quad (23)$$

and

$$(1-\mu) \frac{W_{s,t,p,q}^{\sigma,\delta} g(w)}{w} + \mu \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)' + \gamma w \left( W_{s,t,p,q}^{\sigma,\delta} g(w) \right)'' \\ = \vartheta \left( \frac{y(w)-1}{y(w)+1} \right) = 1 + \frac{1}{2} B_1 y_1 w + \left[ \frac{1}{2} B_1 \left( y_2 - \frac{y_1^2}{2} \right) + \frac{1}{4} B_2 y_1^2 \right] w^2 + \dots \quad (24)$$

Equating the coefficients in (23) and (24), yields

$$\frac{(1+\mu+2\gamma)[\Psi_2(\sigma, s, t)]_{p,q}^\delta}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta} b_2 = \frac{1}{2} B_1 x_1, \quad (25)$$

$$\frac{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta} b_3 = \frac{1}{2} B_1 \left( x_2 - \frac{x_1^2}{2} \right) + \frac{1}{4} B_2 x_1^2, \quad (26)$$

$$- \frac{(1+\mu+2\gamma)[\Psi_2(\sigma, s, t)]_{p,q}^\delta}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta} b_2 = \frac{1}{2} B_1 y_1 \quad (27)$$

and

$$\frac{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta} (2b_2^2 - b_3) = \frac{1}{2} B_1 \left( y_2 - \frac{y_1^2}{2} \right) + \frac{1}{4} B_2 y_1^2. \quad (28)$$

From (25) and (27), we have

$$x_1 = -y_1 \quad (29)$$

and

$$\frac{2(1+\mu+2\gamma)^2[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}{[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}} b_2^2 = \frac{1}{4} B_1^2 (x_1^2 + y_1^2). \quad (30)$$

If we add (26) to (28), we obtain

$$\frac{2(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta} b_2^2 = \frac{1}{2} B_1 \left[ x_2 + y_2 - \left( \frac{x_1^2 + y_1^2}{2} \right) \right] + \frac{1}{4} B_2 [x_1^2 + y_1^2] \quad (31)$$

Substituting the value of  $x_1^2 + y_1^2$  from (30) in the right-hand side of (31), we deduce that

$$b_2^2 = \frac{B_1^3 [\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} (x_2 + y_2)}{4 \left[ B_1^2 (1+2\mu+6\gamma) [\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2 (B_1 - B_2) \right]}. \quad (32)$$

Applying Lemma 1 for the coefficients  $x_1, x_2, y_1, y_2$  in (30) and (32), we obtain

$$|b_2| \leq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta B_1^{\frac{3}{2}}}{\sqrt{\left| B_1^2 (1+2\mu+6\gamma) [\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2 (B_1 - B_2) \right|}},$$

$$|b_2| \leq \frac{B_1 [\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+\mu+2\gamma) [\Psi_2(\sigma, s, t)]_{p,q}^\delta},$$

which gives the estimates of  $|b_2|$ .

Furthermore, in order to find the bound of  $|b_3|$ , we subtract (28) from (26) and also apply (29). Then, we obtain  $x_1^2 = y_1^2$ , and hence,

$$\frac{2(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta} (b_3 - b_2^2) = \frac{1}{2} B_1 (x_2 - y_2), \quad (33)$$

then, by substituting the value of  $b_2^2$  from (30) into (33), gives

$$b_3 = \frac{B_1 [\Psi_1(\sigma, s, t)]_{p,q}^\delta (x_2 - y_2)}{4(1+2\mu+6\gamma) [\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{B_1^2 [\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} (x_1^2 + y_1^2)}{8(1+\mu+2\gamma)^2 [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}.$$

So, we have

$$|b_3| \leq \frac{B_1 [\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma) [\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{B_1^2 [\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{(1+\mu+2\gamma)^2 [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}}$$

In addition, substituting the value of  $b_2^2$  from (31) into (33), we obtain

$$b_3 = \frac{2B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta x_2 + \frac{1}{2}(B_2 - B_1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta (x_1^2 + y_1^2)}{4(1 + 2\mu + 6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}$$

and we have

$$|b_3| \leq \frac{B_2[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1 + 2\mu + 6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta},$$

which gives us the desired estimates of the coefficient  $|b_3|$ .  $\square$

Taking  $\vartheta(z) = \left(\frac{1+z}{1-z}\right)^\alpha = 1 + 2\alpha z + 2\alpha^2 z^2 + \dots$  ( $0 < \alpha \leq 1$ ) in Theorem 2, we obtain the next corollary.

**Corollary 3.** Let  $f$  given by (1) be in the family  $T_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \alpha)$ , where ( $0 < \alpha \leq 1$ ). Then,

$$|b_2| \leq \min \left\{ \frac{2\alpha[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1 + \mu + 2\gamma)[\Psi_2(\sigma, s, t)]_{p,q}^\delta}, \frac{2\alpha\sqrt{2\alpha}[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{\sqrt{|4\alpha^2(1 + 2\mu + 6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta[\Psi_3(\sigma, s, t)]_{p,q}^\delta + 2\alpha(1 - \alpha)(1 + \mu + 2\gamma)^2[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}|}} \right\}$$

and

$$|b_3| \leq \min \left\{ \frac{2\alpha^2[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1 + 2\mu + 6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}, \frac{2\alpha[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1 + 2\mu + 6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{4\alpha^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{(1 + \mu + 2\gamma)^2[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}} \right\}.$$

Taking  $\vartheta(z) = \frac{1+(1-2\beta)z}{1-z} = 1 + 2(1 - \beta)z + 2(1 - \beta)z^2 + \dots$  ( $0 \leq \beta < 1$ ) in Theorem 2, we obtain the next corollary.

**Corollary 4.** Let the function  $f$  given by (1) be in the function family  $T_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \beta)$ , where ( $0 \leq \beta < 1$ ). Then,

$$|b_2| \leq \min \left\{ \frac{2(1 - \beta)[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1 + \mu + 2\gamma)[\Psi_2(\sigma, s, t)]_{p,q}^\delta}, \frac{\sqrt{2(1 - \beta)}[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{\sqrt{|(1 + 2\mu + 6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta[\Psi_3(\sigma, s, t)]_{p,q}^\delta|}} \right\}$$

and

$$|b_3| \leq \min \left\{ \frac{2(1 - \beta)[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1 + 2\mu + 6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}, \frac{2(1 - \beta)[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1 + 2\mu + 6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + \frac{4(1 - \beta)^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta}}{(1 + \mu + 2\gamma)^2[\Psi_2(\sigma, s, t)]_{p,q}^{2\delta}} \right\}.$$

**Remark 3.** The problem of maximizing the absolute value of the functional  $|b_3 - \eta b_2^2|$  is called the Fekete–Szegő problem. Many authors obtained Fekete–Szegő inequalities for different classes of functions. Obtaining Fekete–Szegő inequalities for different classes of functions defined by operators, the study of bi-univalent functions using operators and the study on coefficients of the functions is a topic of interest at this time (see [1–5,7,10,11,14,46–51]).

1. In [1], the authors obtained Fekete–Szegő inequalities and coefficient inequalities for certain classes of bi-univalent functions defined by Horadam Polynomials;
2. In [8], the authors obtained Fekete–Szegő inequalities for classes of analytic and bi-univalent functions defined by  $(p, q)$ -derivative operator;
3. In [23], the authors obtained Fekete–Szegő inequalities for subclasses of analytic and bi-univalent functions defined by subordinations using the Sălăgean operator;
4. In [52], the author obtained Fekete–Szegő inequalities for analytic and bi-univalent functions subordinate to  $(p, q)$ -Lucas Polynomials;
5. In [53], the authors obtained Fekete–Szegő inequalities for analytic and bi-univalent functions subordinate to Gegenbauer polynomials;
6. In [54], the authors obtained Fekete–Szegő inequalities for analytic and bi-univalent functions subordinate to Chebyshev polynomials;
7. In [55], the authors obtained Fekete–Szegő inequalities and coefficients bounds for new classes of bi-univalent functions defined by the Sălăgean integro-differential operator;
8. In [56], the author obtained Fekete–Szegő inequalities for classes of bi-univalent functions defined in terms of subordinations.

In the next theorems, we provide the Fekete–Szegő type inequalities for the functions of the families  $\mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  and  $\mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$ .

**Theorem 3.** For  $\eta \in \mathbb{R}$ , let  $f \in \mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  be of the form (1). Then,

$$|b_3 - \eta b_2^2| \leq \begin{cases} \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}; \\ |\eta - 1| \leq \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta [2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)}{2B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} [\Psi_3(\sigma, s, t)]_{p,q}^\delta (2\lambda+1)}, \\ B_1^3[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} |\eta - 1| \\ \frac{[2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)}{[2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)}; \\ |\eta - 1| \geq \frac{[2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)}{2B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} [\Psi_3(\sigma, s, t)]_{p,q}^\delta (2\lambda+1)}. \end{cases}$$

**Proof.** It follows from (19) and (20) that

$$\begin{aligned} b_3 - \eta b_2^2 &= \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta (x_2 - y_2)}{8(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + (1-\eta)b_2^2 \\ &= \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta (x_2 - y_2)}{8(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} \\ &\quad + \frac{B_1^3[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} (x_2 + y_2)(1-\eta)}{4[2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)} \\ &= \frac{B_1}{4} \left[ \left( Y(\eta) + \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} \right) x_2 + \left( Y(\eta) - \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} \right) y_2 \right], \end{aligned}$$

where

$$Y(\eta) = \frac{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} (1-\eta)}{2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)}.$$

According to Lemma 1 and (5), we find that

$$|b_3 - \eta b_2^2| \leq \begin{cases} \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}, & 0 \leq |Y(\eta)| \leq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}, \\ B_1|Y(\eta)|, & |Y(\eta)| \geq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}. \end{cases}$$

After some computations, we obtain

$$|b_3 - \eta b_2^2| \leq \begin{cases} \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}; \\ |\eta - 1| \leq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta |2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)|}{2B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} [\Psi_3(\sigma, s, t)]_{p,q}^\delta (2\lambda+1)}; \\ B_1^3[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} |\eta - 1| \\ |\eta - 1| \geq \frac{|2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)|}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta |2B_1^2(2\lambda+1)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} ((\lambda+1)^2(B_1-B_2) - (3\lambda+1)B_1^2)|}. \end{cases}$$

□

Putting  $\eta = 1$  in Theorem 3, we obtain the following result.

**Corollary 5.** If  $f \in \mathcal{A}_E(\lambda, \sigma, \delta, s, t, p, q; \vartheta)$  is of the form (1), then

$$|b_3 - b_2^2| \leq \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{2(2\lambda+1)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}.$$

**Theorem 4.** For  $\eta \in \mathbb{R}$ , let  $f \in \mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  be of the form (1). Then,

$$|b_3 - \eta b_2^2| \leq \begin{cases} \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}; \\ |\eta - 1| \leq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta |B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2(B_1-B_2)|}{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} [\Psi_3(\sigma, s, t)]_{p,q}^\delta (1+2\mu+6\gamma)}; \\ B_1^3[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} |\eta - 1| \\ |\eta - 1| \geq \frac{|B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2(B_1-B_2)|}{[\Psi_1(\sigma, s, t)]_{p,q}^\delta |B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2(B_1-B_2)|}. \end{cases}$$

**Proof.** It follows from (32) and (33) that

$$\begin{aligned} b_3 - \eta b_2^2 &= \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta (x_2 - y_2)}{4(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} + (1-\eta)b_2^2 \\ &= \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta (x_2 - y_2)}{4(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} \\ &\quad + \frac{B_1^3[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} (x_2 + y_2)(1-\eta)}{4[B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2(B_1-B_2)]} \\ &= \frac{B_1}{4} \left[ \left( \Omega(\eta) + \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} \right) x_2 + \left( \Omega(\eta) - \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta} \right) y_2 \right], \end{aligned}$$

where

$$\Omega(\eta) = \frac{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} (1-\eta)}{B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2(B_1-B_2)}.$$

According to Lemma 1 and (5), we find that

$$|b_3 - \eta b_2^2| \leq \begin{cases} \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}, & 0 \leq |\Omega(\eta)| \leq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}, \\ B_1|\Omega(\eta)|, & |\Omega(\eta)| \geq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}. \end{cases}$$

After some computations, we obtain

$$|b_3 - \eta b_2^2| \leq \begin{cases} \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}; \\ |\eta - 1| \leq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta |B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2 (B_1-B_2)|}{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} [\Psi_3(\sigma, s, t)]_{p,q}^\delta (1+2\mu+6\gamma)}, \\ \frac{B_1^3[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} |\eta - 1|}{|B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2 (B_1-B_2)|}; \\ |\eta - 1| \geq \frac{[\Psi_1(\sigma, s, t)]_{p,q}^\delta |B_1^2(1+2\mu+6\gamma)[\Psi_1(\sigma, s, t)]_{p,q}^\delta [\Psi_3(\sigma, s, t)]_{p,q}^\delta + [\Psi_2(\sigma, s, t)]_{p,q}^{2\delta} (1+\mu+2\gamma)^2 (B_1-B_2)|}{B_1^2[\Psi_1(\sigma, s, t)]_{p,q}^{2\delta} [\Psi_3(\sigma, s, t)]_{p,q}^\delta (1+2\mu+6\gamma)}. \end{cases}$$

□

Putting  $\eta = 1$  in Theorem 4, we obtain the following result.

**Corollary 6.** If  $f \in \mathcal{S}_E(\mu, \gamma, \sigma, \delta, s, t, p, q; \vartheta)$  is of the form (1), then

$$|b_3 - b_2^2| \leq \frac{B_1[\Psi_1(\sigma, s, t)]_{p,q}^\delta}{(1+2\mu+6\gamma)[\Psi_3(\sigma, s, t)]_{p,q}^\delta}.$$

### 3. Conclusions

As future research directions, the symmetry properties of this newly introduced operator can be studied.

**Author Contributions:** These authors contributed equally to this work. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** The authors would like to thank the referees for their careful reading and helpful comments.

**Conflicts of Interest:** The authors declare no conflict of interest in this paper.

### References

- Abirami, C.; Magesh, N.; Yamini, J. Initial bounds for certain classes of bi-univalent functions defined by Horadam Polynomials. *Abstr. Appl. Anal.* **2020**, *2020*, 7391058. [\[CrossRef\]](#)
- Srivastava, H.M.; Mishra, A.K.; Gochhayat, P. Certain subclasses of analytic and bi-univalent functions. *Appl. Math. Lett.* **2010**, *23*, 1188–1192. [\[CrossRef\]](#)
- Frasin, B.A.; Aouf, M.K. New subclasses of bi-univalent functions. *Appl. Math. Lett.* **2011**, *24*, 1569–1573. [\[CrossRef\]](#)
- Ali, R.M.; Lee, S.K.; Ravichandran, V.; Supramaniam, S. Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions. *Appl. Math. Lett.* **2012**, *25*, 344–351. [\[CrossRef\]](#)
- Bulut, S.; Magesh, N.; Abirami, C. A comprehensive class of analytic bi-univalent functions by means of Chebyshev polynomials. *J. Fract. Calc. Appl.* **2017**, *8*, 32–39.
- Srivastava, H.M.; Wanas, A.K.; Murugusundaramoorthy, G. Certain family of bi-univalent functions associated with Pascal distribution series based on Horadam polynomials. *Surv. Math. Appl.* **2021**, *16*, 193–205.
- Al-Amoush, A.G. Coefficient estimates for a new subclasses of  $\lambda$ -pseudo biunivalent functions with respect to symmetrical points associated with the Horadam Polynomials. *Turk. J. Math.* **2019**, *43*, 2865–2875. [\[CrossRef\]](#)
- Coțîrlă, L.I. New classes of analytic and bi-univalent functions. *AIMS Math.* **2021**, *6*, 10642–10651. [\[CrossRef\]](#)
- Güney, H.Ö.; Murugusundaramoorthy, G.; Sokół, J. Subclasses of bi-univalent functions related to shell-like curves connected with Fibonacci numbers. *Acta Univ. Sapient. Math.* **2018**, *10*, 70–84. [\[CrossRef\]](#)
- Srivastava, H.M.; Altınkaya, Ş.; Yalçın, S. Certain subclasses of bi-univalent functions associated with the Horadam polynomials. *Iran. J. Sci. Technol. Trans. A Sci.* **2019**, *43*, 1873–1879. [\[CrossRef\]](#)



11. Srivastava, H.M.; Gaboury, S.; Ghanim, F. Coefficient estimates for some general subclasses of analytic and bi-univalent functions. *Afr. Mat.* **2017**, *28*, 693–706. [\[CrossRef\]](#)
12. Srivastava, H.M.; Wanas, A.K.; Güney, H.Ö. New families of bi-univalent functions associated with the Bazilevič functions and the  $\lambda$ -Pseudo-starlike functions. *Iran. J. Sci. Technol. Trans. A Sci.* **2021**, *45*, 1799–1804. [\[CrossRef\]](#)
13. Srivastava, H.M.; Wanas, A.K.; Srivastava, R. Applications of the  $q$ -Srivastava-Attiya operator involving a certain family of bi-univalent functions associated with the Horadam polynomials. *Symmetry* **2021**, *13*, 1230. [\[CrossRef\]](#)
14. Wanas, A.K. Applications of (M,N)-Lucas polynomials for holomorphic and bi-univalent functions. *Filomat* **2020**, *34*, 3361–3368. [\[CrossRef\]](#)
15. Wanas, A.K. Horadam polynomials for a new family of  $\lambda$ -pseudo bi-univalent functions associated with Sakaguchi type functions. *Afr. Mat.* **2021**, *32*, 879–889. [\[CrossRef\]](#)
16. Fekete, M.; Szegő, G. Eine bemerkung uber ungerade schlichte funktionen. *J. Lond. Math. Soc.* **1933**, *2*, 85–89. [\[CrossRef\]](#)
17. Cataş, A. A note on subclasses of univalent functions defined by a generalized Sălăgean operator. *Acta Univ. Apulensis* **2006**, *12*, 73–78.
18. Magesh, N.; Yamini, J. Fekete-Szegő problem and second Hankel determinant for a class of bi-univalent functions. *Tbilisi Math. J.* **2018**, *11*, 141–157. [\[CrossRef\]](#)
19. Raina, R.K.; Sokol, J. Fekete-Szegő problem for some starlike functions related to shell-like curves. *Math. Slovaca* **2016**, *66*, 135–140. [\[CrossRef\]](#)
20. Srivastava, H.M.; Raza, N.; AbuJarad, E.S.A.; Srivastava, G.; AbuJarad, M.H. Fekete-Szegő inequality for classes of  $(p, q)$ -starlike and  $(p, q)$ -convex functions. *Revista de la Real Academia de Ciencias Exactas Físicas y Naturales Serie A Matemáticas* **2019**, *113*, 3563–3584. Available online: <https://arxiv.org/abs/1912.05298> (accessed on 1 November 2021).
21. Wanas, A.K.; Lupas, A.A. Applications of Horadam Polynomials on Bazilevic Bi- Univalent Function Satisfying Subordinate Conditions. *J. Phys. Conf. Ser.* **2019**, *1294*, 032003. [\[CrossRef\]](#)
22. Miller, S.S.; Mocanu, P.T. *Differential Subordinations: Theory and Applications*; Series on Monographs and Textbooks in Pure and Applied Mathematics; Marcel Dekker Incorporated: New York, NY, USA; Basel, Switzerland, 2000; Volume 225.
23. Çağlar, M.; Aslan, S. Fekete-Szegő inequalities for subclasses of bi-univalent functions satisfying subordinate conditions. *AIP Conf. Proc.* **2016**, *1726*, 020078. [\[CrossRef\]](#)
24. Jagannathan, R.; Rao, K.S. Two-parameter quantum algebras, twin-basic numbers, and associated generalized hypergeometric series. *arXiv* **2006**, arXiv:math/0602613.
25. Srivastava, H.M. Some generalizations and basic (or  $q$ -) extensions of the Bernoulli, Euler and Genocchi polynomials. *Appl. Math. Inf. Sci.* **2011**, *5*, 390–444.
26. Srivastava, H.M.; Choi, J. *Zeta and  $q$ -Zeta Functions and Associated Series and Integrals*; Elsevier: Amsterdam, The Netherlands, 2012.
27. Victor, K.; Pokman, C. *Quantum Calculus*; Springer: New York, NY, USA, 2002.
28. Chakrabarti, R.; Jagannathan, R. A  $(p, q)$ -oscillator realization of two-parameter quantum algebras. *J. Phys. A Math. Gen.* **1991**, *24*, L711. [\[CrossRef\]](#)
29. Corcino, R.B. On  $p, q$ -binomial coefficients. *Integers* **2008**, *8*, A29.
30. Sadjang, P.N. On the fundamental theorem of  $(p, q)$ -calculus and some  $(p, q)$ -Taylor formulas. *arXiv* **2013**, arXiv:1309.3934.
31. Wanas, A.K. New differential operator for holomorphic functions. *Earthline J. Math. Sci.* **2019**, *2*, 527–537. [\[CrossRef\]](#)
32. Shah, S.A.; Noor, K.I. Study on the  $q$ -analogue of a certain family of linear operators. *Turk J. Math.* **2019**, *43*, 2707–2714. [\[CrossRef\]](#)
33. Noor, K.I.; Riaz, S.; Noor, M.A. On  $q$ -Bernardi integral operator. *TWMS J. Pure Appl. Math.* **2017**, *8*, 3–11.
34. Govindaraj, M.; Sivasubramanian, S. On a class of analytic functions related to conic domains involving  $q$ -calculus. *Anal. Math.* **2017**, *43*, 475–487. [\[CrossRef\]](#)
35. Swamy, S.R. Inclusion properties of certain subclasses of analytic functions. *Int. Math. Forum* **2012**, *7*, 1751–1760.
36. Srivastava, H.M.; Attiya, A.A. An integral operator associated with the Hurwitz-Lerch Zeta function and differential subordination. *Integral Transform. Spec. Funct.* **2007**, *18*, 207–216. [\[CrossRef\]](#)
37. Cho, N.E.; Srivastava, H.M. Argument estimates of certain analytic functions defined by a class of multiplier transformations. *Math. Comput. Model.* **2003**, *37*, 39–49. [\[CrossRef\]](#)
38. Uralegaddi, B.A.; Somanatha, C. Certain classes of univalent functions. In *Current Topics in Analytic Function Theory*; Srivastava, H.M., Own, S., Eds.; World Scientific: Singapore, 1992; pp. 371–374.
39. Jung, I.B.; Kim, Y.C.; Srivastava, H.M. The Hardy space of analytic functions associated with certain one-parameter families of integral operators. *J. Math. Anal. Appl.* **1993**, *176*, 138–147. [\[CrossRef\]](#)
40. Bernardi, S.D. Convex and starlike univalent functions. *Trans. Am. Math. Soc.* **1969**, *135*, 429–446. [\[CrossRef\]](#)
41. Alexander, J.W. Functions which map the interior of the unit circle upon simple region. *Ann. Math.* **1915**, *17*, 12–22. [\[CrossRef\]](#)
42. Al-Oboudi, F.M. On univalent functions defined by a generalized Sălăgean operator. *Int. J. Math. Math. Sci.* **2004**, *27*, 1429–1436. [\[CrossRef\]](#)
43. Sălăgean, G.S. Subclasses of univalent functions. In *Lecture Notes in Mathematics*; Springer: Berlin/Heidelberg, Germany, 1983; Volume 1013, pp. 362–372.
44. Duren, P.L. Grundlehren der Mathematischen Wissenschaften, Band 259. In *Univalent Functions*; Springer: New York, NY, USA; Berlin/Heidelberg, Germany; Tokyo, Japan, 1983.
45. Pommerenke, C. *Univalent Functions*; Vandenhoeck and Ruprecht: Göttingen, Germany, 1975.



- 
46. Li, X.-F.; Wang, A.-P. Two new subclasses of bi-univalent functions. *Int. Math. Forum* **2012**, *7*, 1495–1504.
  47. Brannan, D.A.; Taha, T.S. On some classes of bi-univalent functions. *Stud. Univ. Babeş-Bolyai Math.* **1988**, *31*, 53–60.
  48. Altinkaya, Ş.; Yalçın, S. Chebyshev polynomial coefficient bounds for a subclass of bi-univalent functions. *arXiv* **2017**, arXiv:1605.08224v2.
  49. Frasin, B.A. Coefficient bounds for certain classes of bi-univalent functions. *Hacet. J. Math. Stat.* **2014**, *43*, 383–389. [[CrossRef](#)]
  50. Bulut, S. Faber polynomial coefficient estimates for a subclass of analytic bi-univalent functions. *Filomat* **2016**, *30*, 1567–1575. [[CrossRef](#)]
  51. Altinkaya, Ş.; Yalçın, S. Fekete-Szegő inequalities for certain classes of biunivalent functions. *Int. Sch. Res. Not.* **2014**, 1726. [[CrossRef](#)] [[PubMed](#)]
  52. Amourah, A. Fekete-Szegő inequalities for analytic and bi-univalent functions subordinate to  $(p, q)$ -Lucas Polynomials. *arXiv* **2020**, arXiv:2004.00409.
  53. Amourah, A.; Frasin, B.A.; Abdeljawad, T. Fekete-Szegő inequality for analytic and bi-univalent functions subordinate to Gegenbauer polynomials. *J. Funct. Spaces* **2021**, *2021*, 5574673.
  54. Yousef, F.; Frasin, B.A.; Al-Hawary, T. Fekete-Szegő inequality for analytic and bi-univalent functions subordinate to Chebyshev polynomials. *arXiv* **2018**, arXiv:1801.09531.
  55. Páll-Szabó, A.O.; Oros, G.I. Coefficient Related Studies for New Classes of Bi-Univalent Functions. *Mathematics* **2020**, *8*, 1110. [[CrossRef](#)]
  56. Zaprawa, P. On the Fekete-Szegő problem for classes of bi-univalent functions. *Bull. Belg. Math. Soc. Simon Stevin* **2014**, *21*, 169–178. [[CrossRef](#)]