

Article

Determinant Formulae of Matrices with Certain Symmetry and Its Applications

Yongju Bae [†] and In Sook Lee ^{*,†}

Department of Mathematics, Kyungpook National University, Daegu 702-701, Korea; ybae@knu.ac.kr

* Correspondence: insooki1109@knu.ac.kr; Tel.: +82-53-950-5306

† These authors contributed equally to this work.

Received: 13 August 2017; Accepted: 29 November 2017; Published: 6 December 2017

Abstract: In this paper, we introduce formulae for the determinants of matrices with certain symmetry. As applications, we will study the Alexander polynomial and the determinant of a periodic link which is presented as the closure of an oriented 4-tangle.

Keywords: determinant of a matrix; Seifert matrix of a link; periodic links; Alexander polynomial of a link

MSC: 15A15; 57M25

1. Introduction

A *block matrix* is a matrix which is partitioned into submatrices, called *blocks*, such that the subscripts of the blocks are defined in the same fashion as those for the elements of a matrix [1].

$$\left(\begin{array}{ccc|ccc} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} \\ \hline a_{41} & a_{42} & a_{43} & a_{44} & a_{45} \\ a_{51} & a_{52} & a_{51} & a_{54} & a_{55} \end{array} \right) = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}.$$

Let us consider periodic subject that consists of finite number of units with the same properties, see Figure 1. There are 5 subjects with the same properties which are arranged periodically. The brown color is used to depict their periodical placement, while the green color in the second and third figure presents an extra relationship which is acting between neighbouring two subjects.




Figure 1. Type of periodic subject.

The matrix encoding the properties of the whole subject can be presented by block matrices. In the first picture in Figure 1, there are a finite number of units which are placed periodically but there are no interaction between units as seen. The matrix for the periodic subject will be of the form

$$\begin{pmatrix} A & O & \dots & O \\ O & A & \dots & O \\ \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & A \end{pmatrix}. \tag{1}$$

Indeed, the determinant of the matrix is $(\det A)^n$.

In the second picture in Figure 1, there are a finite number of units which are placed periodically and each unit is affected by neighbouring units as seen. The matrix for such a periodic subject is of the form

$$\begin{pmatrix} A & O & \dots & O & B \\ O & A & \dots & O & B \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \dots & A & B \\ C & C & \dots & C & nD \end{pmatrix}. \tag{2}$$

In [2], the authors showed that the determinant of such a matrix is given by $n^r (\det(A))^{n-1} \det \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where D is an $r \times r$ matrix.

On the other hand, in the last picture in Figure 1, there are a finite number of units which are placed periodically and each unit is affected by the periodic structure itself (rather than neighbouring units). The matrix for such a periodic object can be presented by a matrix of the form

$$\begin{pmatrix} A & O & O & \dots & O & -B & -B & \dots & -B \\ O & A & O & \dots & O & B & O & \dots & O \\ O & O & A & \dots & O & O & B & \dots & O \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \dots & A & O & O & \dots & B \\ -C & C & O & \dots & O & 2D & D & \dots & D \\ -C & O & C & \dots & O & D & 2D & \dots & D \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ -C & O & O & \dots & C & D & D & \dots & 2D \end{pmatrix} \tag{3}$$

or a matrix of the form

$$\begin{pmatrix} A & O & O & \dots & O & -B & -B & \dots & -B \\ O & A & O & \dots & O & B & O & \dots & O \\ O & O & A & \dots & O & O & B & \dots & O \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \dots & A & O & O & \dots & B \\ -C & C & O & \dots & O & D+E & D & \dots & D \\ -C & O & C & \dots & O & E & D+E & \dots & D \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ -C & O & O & \dots & C & E & E & \dots & D+E \end{pmatrix}. \tag{4}$$

The applications in the last section will be helpful to understand the difference between (3) and (4).

In this paper, we will show that the determinant of the matrix (3) is

$$n^r \det(A) \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-1},$$

while the determinant of the matrix (4) is

$$\det(A) \sum_{k=1}^n \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-k} \left\{ \det \begin{pmatrix} A & B \\ C & E \end{pmatrix} \right\}^{k-1}.$$

As an application, we will find the Alexander polynomial and the determinant of a periodic link (Theorems 4–7). Notice that, if a matrix M is singular, then we define $(\det(M))^0 = 1$.

2. Determinants

In this section, we introduce formulae for the determinants of block matrices (3) and (4).

Theorem 1. Let $A, B, C,$ and D be matrices of size $m \times m, m \times r, r \times m$ and $r \times r,$ respectively and O the zero-matrix. Then

$$\det \begin{pmatrix} A & O & O & \cdots & O & -B & -B & \cdots & -B \\ O & A & O & \cdots & O & B & O & \cdots & O \\ O & O & A & \cdots & O & O & B & \cdots & O \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \cdots & A & O & O & \cdots & B \\ -C & C & O & \cdots & O & 2D & D & \cdots & D \\ -C & O & C & \cdots & O & D & 2D & \cdots & D \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ -C & O & O & \cdots & C & D & D & \cdots & 2D \end{pmatrix} = n^r \det(A) \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-1},$$

where the number of A 's in the diagonal is n and the number of $(2D)$'s in the diagonal is $n - 1$ ($n \geq 2$).

Proof. The identity can be obtained by elementary determinant calculation. We put the detailed proof at Appendix A for the convenience of readers. □

Theorem 2. Let A, B, C, D and E be matrices of size $m \times m, m \times 1, 1 \times m, 1 \times 1$ and $1 \times 1,$ respectively and O the zero-matrix. Then

$$\det \begin{pmatrix} A & O & O & \cdots & O & -B & -B & \cdots & -B \\ O & A & O & \cdots & O & B & O & \cdots & O \\ O & O & A & \cdots & O & O & B & \cdots & O \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \cdots & A & O & O & \cdots & B \\ -C & C & O & \cdots & O & D+E & D & \cdots & D \\ -C & O & C & \cdots & O & E & D+E & \cdots & D \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ -C & O & O & \cdots & C & E & E & \cdots & D+E \end{pmatrix} = \det(A) \sum_{k=1}^n \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-k} \left\{ \det \begin{pmatrix} A & B \\ C & E \end{pmatrix} \right\}^{k-1},$$

where the number of A 's in the diagonal is n and the number of $D + E$'s in the diagonal is $n - 1$ ($n \geq 2$).

Proof.

$$\begin{aligned}
 \text{LHS} &= \det \begin{pmatrix} A & O & O & \cdots & O & -B & -B & \cdots & -B \\ O & A & O & \cdots & O & 2B & B & \cdots & B \\ O & O & A & \cdots & O & B & 2B & \cdots & B \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \cdots & A & B & B & \cdots & 2B \\ O & C & O & \cdots & O & D+E & D & \cdots & D \\ O & O & C & \cdots & O & E & D+E & \cdots & D \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \cdots & C & E & E & \cdots & D+E \end{pmatrix} \\
 &= \det(A) \sum_{k=1}^n \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-k} \left\{ \det \begin{pmatrix} A & B \\ C & E \end{pmatrix} \right\}^{k-1}. \tag{2}
 \end{aligned}$$

The identity can be obtained by elementary determinant calculation. The identity (1) comes by adding the k th column to the first column and then, adding the (-1) (the first row) to the k th row for any $k = 2, 3, \dots, n$, while the identity (2) can be found in Appendix A. \square

Remark 1. If A is invertible, then Theorems 1 and 2 can be proved by using the Schur complement. We put those proofs at Appendix B. The Authors appreciate such valuable comments given by our reviewer.

3. Application: Alexander Polynomials and the Determinants of Periodic Links

We start this section by reviewing results of knot theory which are related with the calculation of the Alexander polynomial and the determinant of a link, see [2–4] in detail.

A knot K is an embedding $K : S^1 \hookrightarrow \mathbb{R}^3$ of S^1 into the 3-space \mathbb{R}^3 . A link is a finite disjoint union of knots: $L = K_1 \cup \dots \cup K_n$. Each knot K_i is called a *component* of the link L . Two links L and L' are *equivalent* (or *ambient isotopic*) if one can be transformed into the other via a deformation of \mathbb{R}^3 upon itself. A *diagram* of a link L is a regular projection image $p(L)$ from the link L into \mathbb{R}^2 such that the over-path and the under-path at each double points of $p(L)$ are distinguished. There are examples of a knot and a link in Figure 2. Two link diagrams are *equivalent* if one can be transformed into another by a finite sequence of Reidemeister moves in Figure 3.




Figure 2. A trefoil knot diagram and a Hopf link diagram.




Figure 3. Reidemeister moves.

A *Seifert surface* for an oriented link L in S^3 is a connected compact oriented surface contained in S^3 which has L as its boundary. The following *Seifert algorithm* is one way to get a Seifert surface from a diagram D of L .

Let D be a diagram of an oriented link L . In a small neighborhood of each crossing, make the following local change to the diagram;

Delete the crossing and reconnect the loose ends in the only way compatible with the orientation.

When this has been done at every crossing, the diagram becomes a set of disjoint simple loops in the plane. It is a diagram with no crossings. These loops are called *Seifert circles*. By attaching a disc to each Seifert circle and by connecting a half-twisted band at the place of each crossing of D according to the crossing sign, we get a Seifert surface F for L .

The *Seifert graph* Γ of F is constructed as follows;

Associate a vertex with each Seifert circle and connect two vertices with an edge if their Seifert circles are connected by a twisted band.

Note that the Seifert graph Γ is planar, and that if D is connected, so does Γ . Since Γ is a deformation retract of a Seifert surface F , their homology groups are isomorphic: $H_1(F) \cong H_1(\Gamma)$. Let T be a spanning tree for Γ . For each edge $e \in E(\Gamma) \setminus E(T)$, $T \cup \{e\}$ contains the unique simple closed circuit T_e which represents an 1-cycle in $H_1(F)$. The set $\{T_e \mid e \in E(\Gamma) \setminus E(T)\}$ of these 1-cycles is a homology basis for F . For such a circuit T_e , let T_e^+ denote the circuit in S^3 obtained by lifting slightly along the positive normal direction of F . For $E(\Gamma) \setminus E(T) = \{e_1, \dots, e_n\}$, the *linking number* between T_{e_i} and $T_{e_j}^+$ is defined by

$$lk(T_{e_i}, T_{e_j}^+) = \frac{1}{2} \sum_{\text{crossing } c \in T_{e_i} \cap T_{e_j}^+} \text{sign}(c).$$

A *Seifert matrix* of L associated to F is the $n \times n$ matrix $M = (m_{ij})$ defined by

$$m_{ij} = lk(T_{e_i}, T_{e_j}^+),$$

where $E(\Gamma) \setminus E(T) = \{e_1, \dots, e_n\}$. A Seifert matrix of L depends on the Seifert surface F and the choice of generators of $H_1(F)$.

Let M be any Seifert matrix for an oriented link L . The *Alexander polynomial* $\Delta_L(t) \in \mathbb{Z}[t, t^{-1}]$ and *determinant* of L are defined by

$$\begin{aligned} \Delta_L(t) &\doteq \det(t^{\frac{1}{2}}M - t^{-\frac{1}{2}}M^T) \\ \det(L) &= |\det(M + M^T)| \end{aligned}$$

For details, see [4,5].

For $e, f \in E(\Gamma) \setminus E(T)$, $T_e \cap T_f$ is either an empty set, one vertex or a simple path in the spanning tree T . If $T_e \cap T_f$ is a simple path, v_0 and v_1 are two ends of $T_e \cap T_f$, we may assume that the neighborhood of v_0 looks like Figure 4. In other words, the cyclic order of edges incident to v_0 is given by $T_e \cap T_f, T_e, T_f$ with respect to the positive normal direction of the Seifert surface. Also we may assume that the directions of T_e and T_f are given so that v_0 is the starting point of $T_e \cap T_f$. For, if the direction is reversed, one can change the direction to adapt to our setting so that the resulting linking number changes its sign.




Figure 4. $T_e \cap T_f$.

In [6], the authors showed the following proposition which is the key tool to calculate the linking numbers for Seifert matrix of a link.

The Alexander polynomial of a knot or a link is the first polynomial of the knot theory [7]. The polynomial was reformulate and derived in several different ways over the next 50 years. Perhaps the most satisfying of these is from the homology of the branched cyclic covering space of the knot complement. This reveals the underlying geometry and generalizes to higher dimensions and to a multi-variable version for links. See [4]. Many researchers reformulate the Alexander polynomial as a state sum, Kauffman [8] and Conway [9], etc. Recently, many authors interested in the twisted Alexander polynomial and Knot Floer homology, it provides geometric information of a knot or a link, see [10–13]. The Alexander polynomial is categorified by Knot Floer homology, see [14,15]. Furthermore, since Alexander polynomial is a topological content of quantum invariants, Alexander polynomial is one of the most important invarinat of knot theory, see [16].

Proposition 1 ([6]). For $e, f \in E(\Gamma) \setminus E(T)$, let p and q denote the numbers of edges in $T_e \cap T_f$ corresponding to positive crossings and negative crossings, respectively. Suppose that the local shape of $T_e \cap T_f$ in F looks like Figure 4. Then,

$$lk(T_e, T_f^+) = \begin{cases} -\frac{1}{2}(p - q), & \text{if } p + q \text{ is even;} \\ -\frac{1}{2}(p - q + 1), & \text{if } p + q \text{ is odd, and} \end{cases}$$

$$lk(T_f, T_e^+) = \begin{cases} -\frac{1}{2}(p - q), & \text{if } p + q \text{ is even;} \\ -\frac{1}{2}(p - q - 1), & \text{if } p + q \text{ is odd.} \end{cases}$$

A link L in the 3-sphere S^3 is called a *periodic link* of order n ($n \geq 2$) if there is an orientation preserving auto-homeomorphism ψ of S^3 which satisfies the following conditions; $\psi(L) = L$, $\text{Fix}(\psi)$, the fixed-point set of ψ , is a 1-sphere disjoint from L and ψ is of period n . The link $L/\psi = L_*$ is called the *factor link* of the periodic link L . We denote $\text{Fix}(\psi)/\psi$ by F_* . One of an important concern of knot theory is to find the relationship between periodic links and their factor links, see [17,18]. In 2011, the authors expressed the Seifert matrix of a periodic link which is presented as the closure of a 4-tangle with some extra restrictions, in terms of the Seifert matrix of quotient link in [2].

Let I be a closed interval $[a, b]$ and k a positive integer. Fix k points in the upper plane $I^2 \times \{b\}$ of the cube I^3 and the corresponding k points in the lower plane $I^2 \times \{a\}$. A (k, k) -tangle is obtained by embedding oriented curves and some oriented circles in I^3 so that the end points of the curves are the fixed $2k$ -points. By a $2k$ -tangle, we mean a (k, k) -tangle. Let T be a $2k$ -tangle. For an integer $n \geq 2$, let T^n denote the $2k$ -tangle obtained by stacking T n -times. The *denominator* $D(T)$ of T is defined by connecting the top ends of T to the bottom ends by parallel lines, see Figure 5. In particular, if T is a 4-tangle, then the *numerator* $N(T)$ of T is defined as the last picture in Figure 5. Clearly, $D(T^n)$ is a periodic link of order n whose factor link is $D(T)$. If an orientation is given on $D(T)$, then it induces an orientation of $D(T^n)$. Notice that every (oriented) periodic link can be constructed in this way.

Let T be a 4-tangle and let $D(T)$ denote the denominator of T which is obtained by connecting the top ends and the bottom ends of T by parallel curves α and β . Suppose that $D(T)$ is oriented. Note that the induced orientation at α and β are either opposite or parallel, see Figure 6.

If the induced orientation at α and β are parallel, then α and β are contained in different Seifert circles of $D(T)$. Hence we have the following three cases.

- Case I: The orientations at the end points of the curves in T are opposite and the two outer arcs α and β of $D(T)$ are contained in the same Seifert circle, see the first picture in Figure 7.
- Case II: The orientations at the end points of the curves in T are opposite and the two outer arcs of $D(T)$ are contained in different Seifert circles, see the second picture in Figure 7.
- Case III: The orientations at the end points of the curves in T are parallel, see the last picture in Figure 7.




Figure 5. (k, k) -tangle, denominator and numerator.




Figure 6. Orientation of a tangle.




Figure 7. Three types of denominator $D(T)$.

3.1. Periodic Links with Periodicity in Case I

In Case I and Case II, the numerator $N(T)$ of T is well-defined as an oriented link. In particular, in Case I, the linking number between any Seifert circle of $D(T)$ and the periodic axis F_* is always 0, which is equivalent to that $lk(C, F_*) = 0$ for any Seifert circle C of $D(T)$. For Case I, the authors gave the following criteria for Alexander polynomial of the periodic link $D(T^n)$ by using those of the denominator $D(T)$ and the numerator $N(T)$ of a 4-tangle T .

Proposition 2 ([2]). *Let L be a periodic link of order n with the factor link L_* . Suppose that $L = D(T^n)$ and $L_* = D(T)$ for a 4-tangle T . If $lk(C, F_*) = 0$ for any Seifert circle C of $D(T)$, then the Alexander polynomials of L, L_* and $N(T)$ are related as follows;*

$$\Delta_L(t) = n\Delta_{L_*}(t)(\Delta_{N(T)}(t))^{n-1}.$$

Indeed, to get the result, the authors showed the following proposition and then applied the determinant formula for the matrix (1).

Proposition 3. *If T is a 4-tangle in Case I, then there exist Seifert matrices $M_{D(T)}, M_{N(T)}$, and $M_{D(T^n)}$ of $D(T), N(T)$, and $D(T^n)$, respectively, such that*

$$M_{D(T)} = \begin{pmatrix} M_{N(T)} & B \\ C & D \end{pmatrix} \text{ and}$$

$$M_{D(T^n)} = \begin{pmatrix} M_{N(T)} & O & \cdots & O & B \\ O & M_{N(T)} & \cdots & O & B \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \cdots & M_{N(T)} & B \\ C & C & \cdots & C & nD \end{pmatrix},$$

where B is a column vector, C is a row vector, O is the zero-matrix and the number of block $M_{N(T)}$ is n .

3.2. Periodic Links with Periodicity in Case II

In Case II, there is a Seifert circle C of $D(T)$ such that $lk(C, F_*) \neq 0$. In fact $lk(C_1, F_*) = -lk(C_2, F_*) (= \pm 1)$ where C_1 and C_2 denote Seifert circles in $D(T)$ containing α and β , respectively.

Lemma 1. *If T is a 4-tangle in Case II, then there exist Seifert matrices $M_{D(T)}$, $M_{N(T)}$ and $M_{D(T^n)}$ of $D(T)$, $N(T)$ and $D(T^n)$, respectively, such that*

$$M_{N(T)} = \begin{pmatrix} M_{D(T)} & B \\ C & D \end{pmatrix} \text{ and}$$

$$M_{D(T^n)} = \left(\begin{array}{ccccc|cccc} M_{D(T)} & O & O & \cdots & O & -B & -B & \cdots & -B \\ O & M_{D(T)} & O & \cdots & O & B & O & \cdots & O \\ O & O & M_{D(T)} & \cdots & O & O & B & \cdots & O \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ O & O & O & \cdots & M_{D(T)} & O & O & \cdots & B \\ \hline -C & C & O & \cdots & O & 2D & D & \cdots & D \\ -C & O & C & \cdots & O & D & 2D & \cdots & D \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ -C & O & O & \cdots & C & D & D & \cdots & 2D \end{array} \right),$$

where B is a row vector, C is a column vector, O is the zero-matrix, the number of block $M_{D(T)}$ is n , and the number of block $2D$ is $n - 1$ ($n \geq 2$).

Proof. Suppose that the orientations at the end points of the curves in T are opposite and the two outer arcs of $D(T)$ are contained in different Seifert circles. Without loss of generality, we may assume that T looks like T' as seen in Figure 8 that obtained from T by applying the Reidemeister move II between the left two outer curves and the right two outer curves of T , respectively. Note that $D(T')$, $N(T')$ and $D(T'^n)$ are ambient isotopic to $D(T)$, $N(T)$ and $D(T^n)$, respectively.




Figure 8. Applying Reidemeister Move II.

The Seifert graphs $\Gamma_{D(T')}$ and $\Gamma_{N(T')}$ of $D(T')$ and $N(T')$ are of the form in Figure 9, in which spanning trees $\tau_{D(T')}$ and $\tau_{N(T')}$ of $\Gamma_{D(T')}$ and $\Gamma_{N(T')}$ are given by dotted edges in Figure 9. Notice that

$\Gamma_{N(T')}$ is obtained from $\Gamma_{D(T')}$ by identifying the left vertex u to the right vertex v in $\Gamma_{D(T')}$ as shown in the right picture in Figure 9. If $E(\Gamma_{D(T')}) \setminus E(\tau_{D(T')}) = \{e_1, \dots, e_k\}$, then $E(\Gamma_{N(T')}) \setminus E(\tau_{N(T')}) = \{e_1, \dots, e_k\} \cup \{d\}$, where d is the new edge of $\Gamma_{D(T')}$ as shown in Figure 9.




Figure 9. Seifert graphs for $D(T)$ and $N(T)$.

The corresponding Seifert matrix $M_{D(T')} = [m_{ij}(D(T'))]$ of $D(T')$ is a $k \times k$ matrix, while the Seifert matrix $M_{N(T')} = [m_{ij}(N(T'))]$ of $N(T')$ is a $(k + 1) \times (k + 1)$ matrix. Furthermore, the linking number between T_{e_i} and $T_{e_j}^+$ in $N(T')$ is equal to the linking number between T_{e_i} and $T_{e_j}^+$ in $D(T')$ for all $i, j = 1, 2, \dots, k$, by Proposition 1. Indeed, $m_{ij}(D(T')) = m_{ij}(N(T'))$ for all $i, j = 1, 2, \dots, k$. Hence the Seifert matrix $M_{N(T')}$ of $N(T')$ is given by

$$M_{N(T')} = \begin{pmatrix} M_{D(T')} & B \\ C & D \end{pmatrix},$$

where $B = (lk(T_{e_1}, T_d^+), \dots, lk(T_{e_k}, T_d^+))^T, C = (lk(T_d, T_{e_1}^+), \dots, lk(T_d, T_{e_k}^+))$ and $D = lk(T_d, T_d^+)$.

From now on, we will try to find a Seifert matrix $M_{D(T^m)}$ of $D(T^m)$. The Seifert graph $\Gamma_{D(T^m)}$ of $D(T^m)$ consists of n copies of $\Gamma_{D(T')}$ whose final vertices u and v are used to connect the copies of $\Gamma_{D(T')}$ as shown in Figure 10. Let d_p be the corresponding p th copy of d for all $p = 1, 2, \dots, n$ in $\Gamma_{D(T^m)}$. By removing $(n - 1)$ -copies of the edge d , e.g., d_2, d_3, \dots, d_n in Figure 10, we get a spanning tree $\tau_{D(T^m)}$ of $\Gamma_{D(T^m)}$. Indeed,

$$E(\Gamma_{D(T^m)}) \setminus E(\tau_{D(T^m)}) = \cup_{p=1}^n \{e_1^p, \dots, e_k^p\} \cup \{d_2, d_3, \dots, d_n\},$$

where $\{e_1^p, \dots, e_k^p\}$ is the corresponding p th copy of $\{e_1, \dots, e_k\}$.




Figure 10. Seifert graph for $D(T^n)$.

Since the linking number between $T_{e_i^p}$ and $T_{e_j^+}$ in $D(T^n)$ is equal to the linking number between T_{e_i} and $T_{e_j^+}$ in $D(T')$ for all $i, j = 1, 2, \dots, k$ and $p = 1, \dots, n$, we have $m_{ipjq}(D(T^n)) = m_{ij}(D(T'))$, where $m_{ipjq}(D(T^n)) = lk(T_{e_i^p}, T_{e_j^+})$. If $p \neq q$, since $T_{e_i^p}$ and $T_{e_j^+}$ do not intersect, $m_{ipjq}(D(T^n)) = 0$ for all $i, j = 1, 2, \dots, k$, by Proposition 1. Hence,

$$m_{ipjq}(D(T^n)) = \begin{cases} m_{ij}(D(T')), & \text{if } p = q; \\ 0, & \text{if } p \neq q. \end{cases}$$

On the other hand, since T_{d_p} lies in the 1st copy and p th copy of $T_{e_i} \cap T_{d_p}$, we have $lk(T_{e_i^1}, T_{d_p}^+) = -lk(T_{e_i}, T_d^+) = -B$, $lk(T_{d_p}, T_{e_i^1}^+) = -lk(T_d, T_{e_i}^+) = -C$, $lk(T_{e_i^p}, T_{d_p}^+) = lk(T_{e_i}, T_d^+) = B$ and $lk(T_{d_p}, T_{e_i^p}^+) = lk(T_d, T_{e_i}^+) = C$ for all $p = 2, 3, \dots, n$. For $p \neq q$ and $p, q = 2, 3, \dots, n$, $lk(T_{e_i^p}, T_{d_q}^+) = 0$ and $lk(T_{d_q}, T_{e_i^p}^+) = 0$ since $T_{e_i^p}$ and T_{d_q} do not intersect in $D(T^n)$.

Since T is connected, the generator T_{d_p} runs through 2 copies, in each of which the self linking number of T_d is equal to D for all $p = 2, 3, \dots, n$. Furthermore, since the orientations at the end points of the curves in T are opposite, D is even. Hence, $lk(T_{d_p}, T_{d_p}^+) = 2 \cdot lk(T_d, T_d^+) = 2D$ for all $p = 2, 3, \dots, n$, by Proposition 1. For all $p, q = 2, 3, \dots, n$ and $p \neq q$, $lk(T_{d_p}, T_{d_q}^+) = lk(T_d, T_d^+) = D$. Because generators T_{d_p} and T_{d_q} meet in the just 1st copy. \square

Hence by using Theorem 1, we get the following result.

Theorem 3. Let L be a periodic link of order n ($n \geq 2$) with the factor link L_* . Suppose that $L = D(T^n)$ and $L_* = D(T)$ for a 4-tangle T in Case II. Then the Alexander polynomials of $D(T^n)$, $D(T)$ and $N(T)$ are related as follows;

$$\Delta_L(t) = n\Delta_{L_*}(t)(\Delta_{N(T)}(t))^{n-1}.$$

Proof. By the definition of the Alexander polynomial of a link and by Lemma 1, we have

$$\begin{aligned} \Delta_{N(T)}(t) &= \det \begin{pmatrix} t^{\frac{1}{2}}M_{D(T)} - t^{-\frac{1}{2}}M_{D(T)}^T & t^{\frac{1}{2}}B - t^{-\frac{1}{2}}C^T \\ t^{\frac{1}{2}}C - t^{-\frac{1}{2}}B^T & (t^{\frac{1}{2}} - t^{-\frac{1}{2}})D \end{pmatrix}, \\ \Delta_{D(T^n)}(t) &= n \det(t^{\frac{1}{2}}M_{D(T)} - t^{-\frac{1}{2}}M_{D(T)}^T) \\ &\quad \times \det \begin{pmatrix} t^{\frac{1}{2}}M_{D(T)} - t^{-\frac{1}{2}}M_{D(T)}^T & t^{\frac{1}{2}}B - t^{-\frac{1}{2}}C^T \\ t^{\frac{1}{2}}C - t^{-\frac{1}{2}}B^T & (t^{\frac{1}{2}} - t^{-\frac{1}{2}})D \end{pmatrix}^{n-1} \\ &= n\Delta_{D(T)}(t)(\Delta_{N(T)}(t))^{n-1}. \end{aligned}$$

\square

Since the result in Theorem 3 (Case II) equals that in Proposition 2 (Case I), we can summarize them as

Theorem 4. Let L be a periodic link of order n ($n \geq 2$) with the factor link L_* . Suppose that $L = D(T^n)$ and $L_* = D(T)$ for a 4-tangle T whose numerator $N(T)$ is defined. Then the Alexander polynomials of L , L_* and $N(T)$ are related as follows;

$$\Delta_L(t) = n\Delta_{L_*}(t)(\Delta_{N(T)}(t))^{n-1}.$$

Theorem 5. Let L be a periodic link of order n ($n \geq 2$) with the factor link L_* . Suppose that $L = D(T^n)$ and $L_* = D(T)$ for a 4-tangle T whose numerator $N(T)$ is defined. Then the determinants of L , L_* and $N(T)$ are related as follows;

$$\det(L) = n \det(L_*)\{\det(N(T))\}^{n-1}.$$

Proof. Note that $\det(L) = |\Delta_L(-1)|$ for any oriented link L . By Theorem 4, we have

$$\begin{aligned} \det D(T^n) &= |\Delta_{D(T^n)}(-1)| \\ &= |n\Delta_{D(T)}(-1)\{\Delta_{N(T)}(-1)\}^{n-1}| \\ &= n|\Delta_{D(T)}(-1)||\{\Delta_{N(T)}(-1)\}^{n-1}| \\ &= n \det(D(T))\{\det N(T)\}^{n-1}. \end{aligned}$$

□

Example 1. Consider the oriented 4-tangle T in Figure 11, which is a 4-tangle in Case II.




Figure 11. Example for Case II.

The Seifert matrices of $N(T)$ and $D(T)$ are given by

$$M_{D(T)} = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & -1 \\ 0 & 0 & -2 \end{pmatrix} \text{ and } M_{N(T)} = \left(\begin{array}{ccc|ccc} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & -2 \\ \hline -1 & -1 & -1 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 1 & 1 & 1 & -1 & -1 & -1 \\ \hline -1 & -1 & -1 & -2 & -2 & -2 \end{array} \right),$$

while

$$M_{D(T^3)} = \left(\begin{array}{ccc|ccc|ccc|cc} -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ \hline 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 & 0 & 0 & 0 & -1 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -2 & 0 & -1 \\ \hline 1 & 1 & 1 & -1 & -1 & -1 & 0 & 0 & 0 & -4 & -2 \\ \hline 1 & 1 & 1 & 0 & 0 & 0 & -1 & -1 & -1 & -2 & -4 \end{array} \right).$$

By the direct calculation, one can see that the Alexander polynomials of $D(T)$ and $N(T)$ are

$$\begin{aligned} \Delta_{D(T)}(t) &= -2t^{\frac{3}{2}} + 5t^{\frac{1}{2}} - 5t^{-\frac{1}{2}} + 2t^{-\frac{3}{2}} \text{ and} \\ \Delta_{N(T)}(t) &= 4t^2 - 8t + 9 - 8t^{-1} + 4t^{-2}. \end{aligned}$$

By using Theorem 3, we get the Alexander polynomial of $D(T^3)$;

$$\begin{aligned} \Delta_{D(T^3)}(t) &= 3(-2t^{\frac{3}{2}} + 5t^{\frac{1}{2}} - 5t^{-\frac{1}{2}} + 2t^{-\frac{3}{2}})(4t^2 - 8t + 9 - 8t^{-1} + 4t^{-2})^2 \\ &= -96t^{\frac{11}{2}} + 624t^{\frac{9}{2}} - 2016t^{\frac{7}{2}} + 4344t^{\frac{5}{2}} - 6990t^{\frac{3}{2}} + 8799t^{\frac{1}{2}} - 8799t^{-\frac{1}{2}} \\ &\quad + 6990t^{-\frac{3}{2}} - 4344t^{\frac{5}{2}} + 2016t^{-\frac{7}{2}} - 624t^{-\frac{9}{2}} + 96t^{-\frac{11}{2}}. \end{aligned}$$

Finally, one can get $\det(D(T^3)) = 3(14)(33)^2 = 45738$ by Theorem 5 because $\det(D(T)) = 14$ and $\det(N(T)) = 33$.

3.3. Periodic Links with Periodicity in Case III

In Case III, recall that the orientation of T is given as the left of Figure 12 so that there exist exactly two Seifert circles C_1 and C_2 in $D(T)$ such that $lk(C_1, F_*) = lk(C_2, F_*) (= \pm 1)$. Note that the orientation of T cannot be extended to an orientation of $N(T)$. Define T_+ and T_- by adding a positive crossing and a negative crossing at the bottom of T respectively, as shown in Figure 12.




Figure 12. T_+ and T_- .

Lemma 2. *If T is a 4-tangle in Case III, then there exist Seifert matrices $M_{D(T)}, M_{D(T_+)}, M_{D(T_-)}$ and $M_{D(T^n)}$ of $D(T), D(T_+), D(T_-)$ and $D(T^n)$, respectively, such that*

$$\begin{aligned}
 M_{D(T_+)} &= \begin{pmatrix} M_{D(T)} & B \\ C & D \end{pmatrix}, \\
 M_{D(T_-)} &= \begin{pmatrix} M_{D(T)} & B \\ C & E \end{pmatrix} \text{ and} \\
 M_{D(T^n)} &= \left(\begin{array}{ccccc|cccc}
 M_{D(T)} & O & O & \cdots & O & -B & -B & \cdots & -B \\
 O & M_{D(T)} & O & \cdots & O & B & O & \cdots & O \\
 O & O & M_{D(T)} & \cdots & O & O & B & \cdots & O \\
 \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\
 O & O & O & \cdots & M_{D(T)} & O & O & \cdots & B \\
 \hline
 -C & C & O & \cdots & O & D+E & D & \cdots & D \\
 -C & O & C & \cdots & O & E & D+E & \cdots & D \\
 \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\
 -C & O & O & \cdots & C & E & E & \cdots & D+E
 \end{array} \right),
 \end{aligned}$$

where B is a row vector, C is a column vector, O is the zero-matrix, $E = D + 1$, the number of block $M_{D(T)}$ is n and the number of block $(D + E)$ is $n - 1$.

Proof. Since the process of the proof is similar to that of Theorem 3, we will give briefly sketch of the proof.

The Seifert graphs $\Gamma_{D(T)}, \Gamma_{D(T_+)}$ and $\Gamma_{D(T_-)}$ of $D(T), D(T_+)$ and $D(T_-)$ are of the form in Figure 13, in which spanning trees $\tau_{D(T)}, \tau_{D(T_+)}$ and $\tau_{D(T_-)}$ of $\Gamma_{D(T)}, \Gamma_{D(T_+)}$ and $\Gamma_{D(T_-)}$ are given by dotted edges in Figure 13.

The Seifert graph $\Gamma_{D(T^n)}$ of $D(T^n)$ consists of n copies of $\Gamma_{D(T)}$ whose the end vertices u and v are used to connect the copies of $\Gamma_{D(T)}$ as shown in Figure 14. Let d_p be the corresponding p th copy of d for all $p = 1, 2, \dots, n$ in $D(T^n)$. Notice that by the construction of $D(T^n)$, d and d_* correspond to the same edge in $D(T^n)$, where d and d_* were new edges in Figure 13. By removing $(n - 1)$ -copies of the edge d (or d_*) in $\Gamma_{D(T)}$, e.g., d_2, d_3, \dots, d_n in Figure 14, we get a spanning tree $\tau_{D(T^n)}$ of $D(T^n)$. \square




Figure 13. Seifert graphs of $D(T')$, $D(T'_+)$ and $D(T'_-)$.




Figure 14. Seifert graph of $D(T^m)$.

By using the determinant formula in Theorem 2, we get the following theorem.

Theorem 6. Let L be a periodic link of order n ($n \geq 2$) with the factor link L_* . Suppose that $L = D(T^n)$ and $L_* = D(T)$ for a 4-tangle T in Case III. Then the Alexander polynomials of L , L_* , $D(T_+)$ and $D(T_-)$ are related as follows;

$$\Delta_L(t) = \Delta_{L_*}(t) \sum_{k=1}^n (\Delta_{D(T_+)}(t))^{n-k} (\Delta_{D(T_-)}(t))^{k-1}.$$

Proof. By the definition of the Alexander polynomial of a link and by Lemma 2, we have

$$\Delta_{D(T_+)}(t) = \det \begin{pmatrix} t^{\frac{1}{2}} M_{D(T)} - t^{-\frac{1}{2}} M_{D(T)}^T & t^{\frac{1}{2}} B - t^{-\frac{1}{2}} C^T \\ t^{\frac{1}{2}} C - t^{-\frac{1}{2}} B^T & (t^{\frac{1}{2}} - t^{-\frac{1}{2}}) D \end{pmatrix},$$

$$\Delta_{D(T_-)}(t) = \det \begin{pmatrix} t^{\frac{1}{2}} M_{D(T)} - t^{-\frac{1}{2}} M_{D(T)}^T & t^{\frac{1}{2}} B - t^{-\frac{1}{2}} C^T \\ t^{\frac{1}{2}} C - t^{-\frac{1}{2}} B^T & (t^{\frac{1}{2}} - t^{-\frac{1}{2}}) E \end{pmatrix},$$

$$\begin{aligned} \Delta_{D(T^n)}(t) &= \det(t^{\frac{1}{2}}M_{D(T)} - t^{-\frac{1}{2}}M_{D(T)}^T) \\ &\times \sum_{k=1}^n \left[\left\{ \det \begin{pmatrix} t^{\frac{1}{2}}M_{D(T)} - t^{-\frac{1}{2}}M_{D(T)}^T & t^{\frac{1}{2}}B - t^{-\frac{1}{2}}C^T \\ t^{\frac{1}{2}}C - t^{-\frac{1}{2}}B^T & (t^{\frac{1}{2}} - t^{-\frac{1}{2}})D \end{pmatrix} \right\}^{n-k} \right. \\ &\times \left. \left\{ \det \begin{pmatrix} t^{\frac{1}{2}}M_{D(T)} - t^{-\frac{1}{2}}M_{D(T)}^T & t^{\frac{1}{2}}B - t^{-\frac{1}{2}}C^T \\ t^{\frac{1}{2}}C - t^{-\frac{1}{2}}B^T & (t^{\frac{1}{2}} - t^{-\frac{1}{2}})E \end{pmatrix} \right\}^{k-1} \right] \\ &= \Delta_{D(T)}(t) \sum_{k=1}^n \{\Delta_{D(T_+)}(t)\}^{n-k} \{\Delta_{D(T_-)}(t)\}^{k-1}. \end{aligned}$$

□

In general, the determinant of $D(T^n)$ cannot be calculated by using Theorem 6 because $|\det(A + B)| \neq |\det A| + |\det B|$. But we can calculate the determinant of $D(T^n)$ under certain conditions.

Theorem 7. Let L be a periodic link of order n ($n \geq 2$) with the factor link L_* . Suppose that $L = D(T^n)$ and $L_* = D(T)$ for a 4-tangle T in Case III. If $(-1)^{m+1}\Delta_{D(T_+)}(-1) \cdot \Delta_{D(T_-)}(-1) \geq 0$, then

$$\det(D(T^n)) = \det(D(T)) \sum_{k=1}^n \{\det(D(T_+))\}^{n-k} \{\det(D(T_-))\}^{k-1},$$

where m is the size of a Seifert matrix of $D(T)$.

Proof. Notice that, $\Delta_L(-1) = i^m \det(M + M^T)$ for a Seifert matrix $M_{m \times m}$ of a link L .

From the definition of the determinant of a link, Theorem 2 and Lemma 2, if the Seifert matrix $M_{D(T)}$ of $D(T)$ is an $m \times m$ matrix and $(-1)^{m+1}\Delta_{D(T_+)}(-1) \cdot \Delta_{D(T_-)}(-1) \geq 0$, then the determinant of $D(T^n)$ is

$$\begin{aligned} &|\det((-1)^{\frac{1}{2}}M_{D(T)} - (-1)^{-\frac{1}{2}}M_{D(T)}^T) \\ &\times \sum_{k=1}^n \left[\left\{ \det \begin{pmatrix} (-1)^{\frac{1}{2}}M_{D(T)} - (-1)^{-\frac{1}{2}}M_{D(T)}^T & (-1)^{\frac{1}{2}}B - (-1)^{-\frac{1}{2}}C^T \\ (-1)^{\frac{1}{2}}C - (-1)^{-\frac{1}{2}}B^T & ((-1)^{\frac{1}{2}} - (-1)^{-\frac{1}{2}})D \end{pmatrix} \right\}^{n-k} \right. \\ &\times \left. \left\{ \left(\begin{pmatrix} (-1)^{\frac{1}{2}}M_{D(T)} - (-1)^{-\frac{1}{2}}M_{D(T)}^T & (-1)^{\frac{1}{2}}B - (-1)^{-\frac{1}{2}}C^T \\ (-1)^{\frac{1}{2}}C - (-1)^{-\frac{1}{2}}B^T & ((-1)^{\frac{1}{2}} - (-1)^{-\frac{1}{2}})E \end{pmatrix} \right)^{k-1} \right\} \right] \\ &= |\Delta_{D(T)}(-1)| \times \left| \sum_{k=1}^n \{\Delta_{D(T_+)}(-1)\}^{n-k} \{\Delta_{D(T_-)}(-1)\}^{k-1} \right| \\ &= |\Delta_{D(T)}(-1)| \times \left| \sum_{k=1}^n \{i^{m+1} \det(M_{D(T_+)} + M_{D(T_+)}^T)\}^{n-k} \{i^{m+1} \det(M_{D(T_-)} + M_{D(T_-)}^T)\}^{k-1} \right| \\ &= |\Delta_{D(T)}(-1)| \times \left| \sum_{k=1}^n \{\det(M_{D(T_+)} + M_{D(T_+)}^T)\}^{n-k} \{\det(M_{D(T_-)} + M_{D(T_-)}^T)\}^{k-1} \right| \\ &= |\Delta_{D(T)}(-1)| \sum_{k=1}^n |\det(M_{D(T_+)} + M_{D(T_+)}^T)|^{n-k} |\det(M_{D(T_-)} + M_{D(T_-)}^T)|^{k-1} \\ (1) \quad &= \det(D(T)) \sum_{k=1}^n \{\det(D(T_+))\}^{n-k} \{\det(D(T_-))\}^{k-1}. \end{aligned}$$

The identity (1) comes by the condition $(-1)^{m+1} \Delta_{D(T_+)}(-1) \cdot \Delta_{D(T_-)}(-1) \geq 0$ because

$$\begin{aligned} & \det(M_{D(T_+)} + M_{D(T_+)}^T) \det(M_{D(T_-)} + M_{D(T_-)}^T) \geq 0 \\ \Leftrightarrow & (i)^{-(m+1)} \Delta_{D(T_+)}(-1) \cdot (i)^{-(m+1)} \Delta_{D(T_-)}(-1) \geq 0 \\ \Leftrightarrow & (-1)^{m+1} \Delta_{D(T_+)}(-1) \cdot \Delta_{D(T_-)}(-1) \geq 0. \end{aligned}$$

□

Example 2. Consider the oriented 4-tangle T in Figure 15, which is a 4-tangle in Case III.




Figure 15. Example for Case III.

The Seifert matrices of $D(T)$, $D(T_+)$ and $D(T_-)$ are given by

$$M_{D(T)} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, M_{D(T_+)} = \begin{pmatrix} 1 & 0 & 0 & 0 & | & 0 \\ -1 & 1 & 0 & 0 & | & 1 \\ -1 & 0 & 1 & 0 & | & 1 \\ 0 & 0 & 1 & 1 & | & 1 \\ \hline 0 & 0 & 0 & 0 & | & 0 \end{pmatrix} \text{ and } M_{D(T_-)} = \begin{pmatrix} 1 & 0 & 0 & 0 & | & 0 \\ -1 & 1 & 0 & 0 & | & 1 \\ -1 & 0 & 1 & 0 & | & 1 \\ 0 & 0 & 1 & 1 & | & 1 \\ \hline 0 & 0 & 0 & 0 & | & 1 \end{pmatrix},$$

while

$$M_{D(T^3)} = \begin{pmatrix} 1 & 0 & 0 & 0 & | & 0 & 0 & 0 & 0 & | & 0 & | & 0 \\ -1 & 1 & 0 & 0 & | & 0 & 0 & 0 & 0 & | & -1 & | & -1 \\ -1 & 0 & 1 & 0 & | & 0 & 0 & 0 & 0 & | & -1 & | & -1 \\ 0 & 0 & 1 & 1 & | & 0 & 0 & 0 & 0 & | & -1 & | & -1 \\ \hline 0 & 0 & 0 & 0 & | & 1 & 0 & 0 & 0 & | & 0 & | & 0 \\ 0 & 0 & 0 & 0 & | & -1 & 1 & 0 & 0 & | & 1 & | & 0 \\ 0 & 0 & 0 & 0 & | & -1 & 0 & 1 & 0 & | & 1 & | & 0 \\ 0 & 0 & 0 & 0 & | & 0 & 0 & 1 & 1 & | & 1 & | & 0 \\ \hline 0 & 0 & 0 & 0 & | & 0 & 0 & 0 & 0 & | & 1 & | & 0 \\ 0 & 0 & 0 & 0 & | & 0 & 0 & 0 & 0 & | & 0 & | & 0 \\ 0 & 0 & 0 & 0 & | & 0 & 0 & 0 & 0 & | & -1 & | & 1 \\ 0 & 0 & 0 & 0 & | & 0 & 0 & 1 & 0 & | & 0 & | & 1 \\ 0 & 0 & 0 & 0 & | & 0 & 0 & 1 & 1 & | & 0 & | & 1 \\ \hline 0 & 0 & 0 & 0 & | & 0 & 0 & 0 & 0 & | & 1 & | & 0 \\ 0 & 0 & 0 & 0 & | & 0 & 0 & 0 & 0 & | & 1 & | & 1 \end{pmatrix}.$$

By direct calculation, one can see that the Alexander polynomials of $D(T)$, $D(T_+)$ and $D(T_-)$ are

$$\begin{aligned} \Delta_{D(T)}(t) &= t^2 - t + 1 - t^{-1} + t^{-2}, \\ \Delta_{D(T_+)}(t) &= 2t^{\frac{3}{2}} - 3t^{\frac{1}{2}} + 3t^{-\frac{1}{2}} - 2t^{-\frac{3}{2}} \text{ and} \\ \Delta_{D(T_-)}(t) &= t^{\frac{5}{2}} - t^{\frac{1}{2}} + t^{-\frac{1}{2}} - t^{-\frac{5}{2}}. \end{aligned}$$

By using Theorem 6, we get the Alexander polynomial of $D(T^3)$;

$$\begin{aligned} &(t^2 - t + 1 - t^{-1} + t^{-2})\{(2t^{\frac{3}{2}} - 3t^{\frac{1}{2}} + 3t^{-\frac{1}{2}} - 2t^{-\frac{3}{2}})^2 \\ &+ (2t^{\frac{3}{2}} - 3t^{\frac{1}{2}} + 3t^{-\frac{1}{2}} - 2t^{-\frac{3}{2}})(t^{\frac{5}{2}} - t^{\frac{1}{2}} + t^{-\frac{1}{2}} - t^{-\frac{5}{2}}) + (t^{\frac{5}{2}} - t^{\frac{1}{2}} + t^{-\frac{1}{2}} - t^{-\frac{5}{2}})^2\} \\ &= t^7 + t^6 + 4t^5 - 19t^4 + 38t^3 - 61t^2 + 82t - 92 + 82t^{-1} - 61t^{-2} + 38t^{-3} \\ &\quad - 19t^{-4} + 4t^{-5} + t^{-6} + t^{-7}. \end{aligned}$$

Remark 2. In Theorem 7, the condition $(-1)^{m+1}\Delta_{D(T_+)}(-1) \cdot \Delta_{D(T_-)}(-1) \geq 0$ is essential. Consider the oriented 4-tangle T in Figure 15, which is a 4-tangle in Case III. By direct calculation, one can see that the determinant of $D(T^3) = 500$. The result is the same with $\det(D(T^3)) = 5\{(-10)^2 + (-10)(0) + (0)^2\}$ in Theorem 7. We can easily check this example satisfies the condition. Consider the oriented 4-tangle T in Figure 16, which is a 4-tangle in Case III.




Figure 16. $(-1)^3\Delta_{D(T_+)}(-1)\Delta_{D(T_-)}(-1) = -8$.

The Seifert matrices of $D(T)$, $D(T_+)$ and $D(T_-)$ are given by

$$M_{D(T)} = \begin{pmatrix} -1 & -1 \\ 0 & -1 \end{pmatrix}, M_{D(T_+)} = \begin{pmatrix} -1 & -1 & -1 \\ 0 & -1 & -1 \\ 0 & 0 & -1 \end{pmatrix} \text{ and } M_{D(T_-)} = \begin{pmatrix} -1 & -1 & -1 \\ 0 & -1 & -1 \\ 0 & 0 & 0 \end{pmatrix},$$

while

$$M_{D(T^3)} = \begin{pmatrix} -1 & -1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ \hline 0 & 0 & -1 & -1 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 & 0 \\ \hline 0 & 0 & 0 & 0 & -1 & -1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & -1 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

By direct calculation, one can see that the Alexander polynomials of $D(T)$, $D(T_+)$ and $D(T_-)$ are

$$\begin{aligned} \Delta_{D(T)}(t) &= t^{-1} - 1 + t, \\ \Delta_{D(T_+)}(t) &= t^{-\frac{3}{2}} - t^{-\frac{1}{2}} + t^{\frac{1}{2}} - t^{\frac{3}{2}} \text{ and} \\ \Delta_{D(T_-)}(t) &= t^{-\frac{1}{2}} - t^{\frac{1}{2}}. \end{aligned}$$

By using Theorem 6, we get the Alexander polynomial of $D(T^3)$;

$$\begin{aligned} &(t^{-1} - 1 + t)\{(t^{-\frac{3}{2}} - t^{-\frac{1}{2}} + t^{\frac{1}{2}} - t^{\frac{3}{2}})^2 + (t^{-\frac{3}{2}} - t^{-\frac{1}{2}} + t^{\frac{1}{2}} - t^{\frac{3}{2}})(t^{-\frac{1}{2}} - t^{\frac{1}{2}}) + (t^{-\frac{1}{2}} - t^{\frac{1}{2}})^2\} \\ &= t^4 - 2t^3 + 4t^2 - 7t + 8 - 7t^{-1} + 4t^{-2} - 2t^{-3} + t^{-4}. \end{aligned}$$

Finally, one can see that $\det(D(T^3)) = 9$ by direct calculation. The result is not equal to $\det(D(T))\{\det(D(T_+))^2 + \det(D(T_+)\det(D(T_-)) + \det(D(T_-))^2\} = 3\{(4)^2 + (4)2 + 2^2\}$ since $\det(D(T)) = 3$, $\det(D(T_+)) = 4$ and $\det(D(T_-)) = 2$. We can check that this example doesn't satisfy the condition since $(-1)^3\Delta_{D(T_+)}(-1)\Delta_{D(T_-)}(-1) = -(4i)(-2i) = -8 < 0$.

Acknowledgments: This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2016R1D1A3B01007669).

Author Contributions: These authors contributed equally to this work.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Proof of Theorem 1.

$$\begin{aligned} LHS &= \det \begin{pmatrix} A & O & O & \dots & O & -B & -B & \dots & -B \\ O & A & O & \dots & O & 2B & B & \dots & B \\ O & O & A & \dots & O & B & 2B & \dots & B \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \dots & A & B & B & \dots & 2B \\ O & C & O & \dots & O & 2D & D & \dots & D \\ O & O & C & \dots & O & D & 2D & \dots & D \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \dots & C & D & D & \dots & 2D \end{pmatrix} \\ &= \det(A) \det \begin{pmatrix} A & O & \dots & O & 2B & B & \dots & nB \\ O & A & \dots & O & B & 2B & \dots & nB \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & A & B & B & \dots & nB \\ C & O & \dots & O & 2D & D & \dots & nD \\ O & C & \dots & O & D & 2D & \dots & nD \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & C & D & D & \dots & nD \end{pmatrix} \\ &= n^r \det(A) \det \begin{pmatrix} A & O & \dots & O & 2B & B & \dots & B \\ O & A & \dots & O & B & 2B & \dots & B \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & A & B & B & \dots & B \\ C & O & \dots & O & 2D & D & \dots & D \\ O & C & \dots & O & D & 2D & \dots & D \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & C & D & D & \dots & D \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
 &= n^r \det(A) \det \begin{pmatrix} A & O & \cdots & O & B & O & \cdots & B \\ O & A & \cdots & O & O & B & \cdots & B \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \cdots & A & O & O & \cdots & B \\ C & O & \cdots & O & D & O & \cdots & D \\ O & C & \cdots & O & O & D & \cdots & D \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \cdots & C & O & O & \cdots & D \end{pmatrix} \\
 (4) & \\
 &= n^r \det(A) \det \begin{pmatrix} A & O & \cdots & O & B & O & \cdots & O \\ O & A & \cdots & O & O & B & \cdots & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \cdots & A & O & O & \cdots & B \\ C & O & \cdots & O & D & O & \cdots & O \\ O & C & \cdots & O & O & D & \cdots & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \cdots & C & O & O & \cdots & D \end{pmatrix} \\
 (5) & \\
 &= n^r \det(A) \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-1}.
 \end{aligned}$$

The reasons for the identities (1)–(5) are the following;

- (1) Add the k th column to the first column and then, add (-1) (the first row) to the k th row for any $k = 2, 3, \dots, n$.
- (2) Add the k th column to the last column for any $k = n, n + 1, \dots, 2n - 3$.
- (3) B and D are an $m \times r$ and an $r \times r$ matrices, respectively.
- (4) Add (-1) (the last column) to the k th column for any $k = n, n + 1, \dots, 2n - 3$.
- (5) Add (-1) (the k th column) to the last column for any $k = n, n + 1, \dots, 2n - 3$.

It is the end of the proof of Theorem 1. \square

Proof of Theorem 2. To prove

$$\begin{aligned}
 &\det \begin{pmatrix} A & O & \cdots & O & 2B & B & \cdots & B \\ O & A & \cdots & O & B & 2B & \cdots & B \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \cdots & A & B & B & \cdots & 2B \\ C & O & \cdots & O & D+E & D & \cdots & D \\ O & C & \cdots & O & E & D+E & \cdots & D \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \cdots & C & E & E & \cdots & D+E \end{pmatrix} \\
 &= \sum_{k=1}^n \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-k} \left\{ \det \begin{pmatrix} A & B \\ C & E \end{pmatrix} \right\}^{k-1},
 \end{aligned}$$

where the number of A 's and the number of $(D + E)$'s in the diagonal are $n - 1$ ($n \geq 2$), we proceed by the mathematical induction on n , the number of block A . It is clear that

$$\det \begin{pmatrix} A & 2B \\ C & D+E \end{pmatrix} = \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} + \det \begin{pmatrix} A & B \\ C & E \end{pmatrix}.$$

Assume that the formula is true for $n - 2$ ($n \geq 3$). It is well-known that the determinant can be obtained by adding the determinants of the following two matrices.

$$\begin{pmatrix} A & O & \cdots & O & O & 2B & B & \cdots & B & B \\ O & A & \cdots & O & O & B & 2B & \cdots & B & B \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \cdots & A & O & B & B & \cdots & 2B & B \\ O & O & \cdots & O & A & B & B & \cdots & B & B \\ C & O & \cdots & O & O & D+E & D & \cdots & D & D \\ O & C & \cdots & O & O & E & D+E & \cdots & D & D \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \cdots & C & O & E & E & \cdots & D+E & D \\ O & O & \cdots & O & C & E & E & \cdots & E & D \end{pmatrix}$$

and

$$\begin{pmatrix} A & O & \cdots & O & O & 2B & B & \cdots & B & O \\ O & A & \cdots & O & O & B & 2B & \cdots & B & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \cdots & A & O & B & B & \cdots & 2B & O \\ O & O & \cdots & O & A & B & B & \cdots & B & B \\ C & O & \cdots & O & O & D+E & D & \cdots & D & O \\ O & C & \cdots & O & O & E & D+E & \cdots & D & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \cdots & C & O & E & E & \cdots & D+E & O \\ O & O & \cdots & O & C & E & E & \cdots & E & E \end{pmatrix}$$

By simple calculation, one can calculate their determinants. Indeed, the determinant of the first matrix (I) is

$$\begin{aligned} \text{(I)} &= \det \begin{pmatrix} A & O & \cdots & O & -A & B & O & \cdots & O & O \\ O & A & \cdots & O & -A & O & B & \cdots & O & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \cdots & A & -A & O & O & \cdots & B & O \\ O & O & \cdots & O & A & B & B & \cdots & B & B \\ C & O & \cdots & O & -C & D & D-E & \cdots & D-E & O \\ O & C & \cdots & O & -C & O & D & \cdots & D-E & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \cdots & C & -C & O & O & \cdots & D & O \\ O & O & \cdots & O & C & E & E & \cdots & E & D \end{pmatrix} \\ &= \det \begin{pmatrix} A & O & \cdots & O & | & O & B & O & \cdots & O & | & O \\ O & A & \cdots & O & | & O & O & B & \cdots & O & | & O \\ \vdots & \vdots & \ddots & \vdots & | & \vdots & \vdots & \vdots & \ddots & \vdots & | & \vdots \\ O & O & \cdots & A & | & O & O & O & \cdots & B & | & O \\ \hline O & O & \cdots & O & | & A & O & O & \cdots & O & | & B \\ C & O & \cdots & O & | & O & D & D-E & \cdots & D-E & | & O \\ O & C & \cdots & O & | & O & O & D & \cdots & D-E & | & O \\ \vdots & \vdots & \ddots & \vdots & | & \vdots & \vdots & \vdots & \ddots & \vdots & | & \vdots \\ O & O & \cdots & C & | & O & O & O & \cdots & D & | & O \\ O & O & \cdots & O & | & C & E-D & E-D & \cdots & E-D & | & D \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
 &= \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \det \begin{pmatrix} A & O & \dots & O & B & O & \dots & O \\ O & A & \dots & O & O & B & \dots & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & A & O & O & \dots & B \\ C & O & \dots & O & D & D-E & \dots & D-E \\ O & C & \dots & O & O & D & \dots & D-E \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & C & O & O & \dots & D \end{pmatrix} \\
 &= \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-1}, \tag{3}
 \end{aligned}$$

while the determinant of the second matrix (II) is

$$\begin{aligned}
 \text{(II)} &= \det \begin{pmatrix} A & O & \dots & O & O & 2B & B & \dots & B & O \\ O & A & \dots & O & O & B & 2B & \dots & B & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \dots & A & O & B & B & \dots & 2B & O \\ \hline O & O & \dots & O & A & O & O & \dots & O & B \\ C & O & \dots & O & O & D+E & D & \dots & D & O \\ O & C & \dots & O & O & E & D+E & \dots & D & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ O & O & \dots & C & O & E & E & \dots & D+E & O \\ \hline O & O & \dots & O & C & O & O & \dots & O & E \end{pmatrix} \\
 &= \det \begin{pmatrix} A & B \\ C & E \end{pmatrix} \det \begin{pmatrix} A & O & \dots & O & 2B & B & \dots & B \\ O & A & \dots & O & B & 2B & \dots & B \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & A & B & B & \dots & 2B \\ C & O & \dots & O & D+E & D & \dots & D \\ O & C & \dots & O & E & D+E & \dots & D \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & C & E & E & \dots & D+E \end{pmatrix} \\
 &= \det \begin{pmatrix} A & B \\ C & E \end{pmatrix} \sum_{k=1}^{n-1} \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-1-k} \left\{ \det \begin{pmatrix} A & B \\ C & E \end{pmatrix} \right\}^{k-1}.
 \end{aligned}$$

The reasons for the identities (1)–(4) are the following;

- (1) Add (-1) {the $(n - 1)$ th row} to the k th row for any $k = 1, 2, \dots, n - 2$, and then, add (-1) {the $(2n - 2)$ th row} to the k th row for any $k = n, n + 1, \dots, 2n - 3$.
- (2) Add the k th column to the $(n - 1)$ th column for any $k = 1, 2, \dots, n - 2$, and then, add (-1) {the $(2n - 2)$ th column} to the k th column for any $k = n, n + 1, \dots, 2n - 3$.
- (3) Apply the following identity repeatedly.

$$\det \begin{pmatrix} A & O & \dots & O & B & O & \dots & O \\ O & A & \dots & O & O & B & \dots & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \hline O & O & \dots & A & O & O & \dots & B \\ C & O & \dots & O & D & D-E & \dots & D-E \\ O & C & \dots & O & O & D & \dots & D-E \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \hline O & O & \dots & C & O & O & \dots & D \end{pmatrix}$$

$$= \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \det \begin{pmatrix} A & O & \cdots & O & B & O & \cdots & O \\ O & A & \cdots & O & O & B & \cdots & O \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \cdots & A & O & O & \cdots & B \\ C & O & \cdots & O & D & D-E & \cdots & D-E \\ O & C & \cdots & O & O & D & \cdots & D-E \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ O & O & \cdots & C & O & O & \cdots & D \end{pmatrix}.$$

(4) Add (-1) {the $(2n - 2)$ th column} to the k th column for any $k = n, n + 1, \dots, 2n - 3$.

Therefore the determinant of our matrix is given by

$$\sum_{k=1}^n \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-k} \left\{ \det \begin{pmatrix} A & B \\ C & E \end{pmatrix} \right\}^{k-1}.$$

By using this result, we can prove Theorem 2. \square

Appendix B

The following proofs use more sophisticated matrix theory tools and henceforth are more compact. We assume that A is invertible so that the Schur complement can be defined. However this is only a technical assumption: in fact, looking at the final formula in Theorem 1, this depends with continuity with respect to the entries of A (and in particular with respect to $\det(A)$) and therefore if the statement holds for $\det(A)$ different from zero, then it must be true also when $\det(A) = 0$ and the reason is that the set of invertible matrices is dense in the set of all matrices (see [19]).

Two preliminary things are needed. The first is the Schur complement (see [20] and references therein) of a block matrix $X = \begin{pmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{pmatrix}$ which corresponds at a single block step of the Gaussian Elimination so that the determinant of X is equal to

$$\det(A_{1,1}) \det(S), \quad S = A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2}. \tag{A1}$$

The second is the tensor product [19] of square matrices Y, Z so that

$$\det(Y \otimes Z) = \{\det(Y)\}^v \{\det(Z)\}^u, \tag{A2}$$

where Y is square of size u and Z is square of size v .

Now we are ready to prove Theorem 1. First we observe that $A_{1,1} = I_n \times A$ so that by (A2) we have $\det(A_{1,1}) = \{\det(A)\}^n$. Second we compute the Schur complement S according to (A1) and we find

$$S = (I_{n-1} + ee^T) \otimes (D - CA^{-1}B)$$

with $e^T = (1, \dots, 1)$ vector of all ones of size $n - 1$. Now $D - CA^{-1}B$ is the Schur complement of $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ and hence by the first part of (A1) we have $\det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(A) \det(D - CA^{-1}B)$. The matrix ee^T has rank 1 and hence it has $n - 2$ eigenvalues equal to zero and one coinciding with the trace, that is $n - 1$: hence the matrix $I_{n-1} + ee^T$ has eigenvalue 1 with multiplicity $n - 2$ and eigenvalue n with multiplicity 1 so that its determinant is n

$$\begin{aligned} \det(S) &= \left\{ \det(I_{n-1} + ee^T) \right\}^r \left\{ \det(D - CA^{-1}B) \right\}^{n-1} \\ &= n^r \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-1} \{\det(A)\}^{1-n} \end{aligned}$$

so that, by putting together the latter relation and $\det(A_{1,1}) = \{\det(A)\}^n$, we obtain

$$\det(X) = \det \begin{pmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{pmatrix} = n^r \det(A) \left\{ \det \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right\}^{n-1}$$

and Theorem 1 is proved.

With the very same tools and by following the same steps, Theorem 2 can be proven as well.

References

- Howard, A. *Elementary Linear Algebra*; John Wiley & Sons: New York, NY, USA, 1994.
- Bae, Y.; Lee, I.S. On Alexander polynomial of periodic links. *J. Knot Theory Ramif.* **2011**, *20*, 749–761.
- Banks, J.E. Homogeneous links, Seifert surfaces, digraphs and the reduced Alexander polynomial. *Geom. Dedicata* **2013**, *166*, 67–98.
- Cromwell, P. *Knots and Links*; Cambridge University Press: Cambridge, UK, 2004.
- Kawauchi, A. *A Survey of Knot Theory*; Birkhäuser Verlag: Basel, Switzerland; Boston, MA, USA, 1996.
- Bae, Y.; Lee, I.S. On Seifert matrices of symmetric links. *Kyungpook Math. J.* **2011**, *51*, 261–281.
- Alexander, J. Topological invariants for knots and links. *Trans. Am. Math. Soc.* **1928**, *30*, 275–306.
- Kauffman, L.H. *Formal Knot Theory*; Mathematical Notes, 30; Princeton University Press: Princeton, NJ, USA, 1983.
- Conway, J. An enumeration of knots and links and some of their algebraic properties. In Proceedings of the Conference on Computational Problems in Abstract Algebra, Oxford, UK, 29 August–2 September 1967.
- Hillman, J.A.; Livingston, C.; Naik, S. Twisted Alexander polynomials of periodic knots. *Algebr. Geom. Topol.* **2006**, *6*, 145–169.
- Kitayama, T. Twisted Alexander polynomials and ideal points giving Seifert surfaces. *Acta Math. Vietnam.* **2014**, *39*, 567–574.
- Ozsváth, P.; Szabó, Z. Holomorphic disks and knot invariants. *Adv. Math.* **2004**, *186*, 58–116.
- Rasmussen, J.A. Flouer Homology and Knot Complements. Ph.D. Thesis, Harvard University, Cambridge, MA, USA, March 2003.
- Kauffman, L.H.; Silvero, M. Alexander-Conway polynomial state model and link homology. *J. Knot Theory Ramif.* **2015**, *25*, doi:10.1142/S0218216516400058.
- Ozsváth, P.; Szabó, Z. An introduction to Heegaard Flouer homology. Available online: <http://math.mit.edu/~petero/Introduction.pdf> (accessed on 12 August 2017).
- Sartori, A. The Alexander polynomial as quantum invariant of links. *Ark. Mat.* **2015**, *53*, 1–26.
- Murasugi, K. On periodic knots. *Comment. Math. Helv.* **1971**, *46*, 162–177.
- Sakuma, M.A. On the polynomials of periodic links. *Math. Ann.* **1981**, *257*, 487–494.
- Bhatia, R. *Matrix Analysis*; Springer Verlag: New York, NY, USA, 1997.
- Dorostkar, A.; Neytcheva, M.; Serra-Capizzano, S. Spectral analysis of coupled PDEs and of their Schur complements via the notion of Generalized Locally Toeplitz sequences. *Comput. Methods Appl. Mech. Eng.* **2016**, *309*, 74–105.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).