

**Table S1.** Antifungal susceptibility of *Candida* isolates to amphotericin B and caspofungin. %resistant: percentage of isolates resistant to antifungal.

<i>Candida</i> species	n	Amphotericin B				Caspofungin			
		Sensitive	intermediate	resistant	% resistant	n	sensitive	resistant	%resistant
<i>C. albicans</i>	20	20	0	0	0.0%	5	5	0	0.0%
<i>C. glabrata</i>	20	19	0	1	5.0%	5	5	0	0.0%
<i>C. tropicalis</i>	8	8	0	0	0.0%	2	2	0	0.0%
<i>C. parapsilosis</i>	5	5	0	0	0.0%	1	1	0	0.0%
<i>P. kudriavzevii</i> ( <i>C. krusei</i> )	2	2	0	0	0.0%	0	0	0	0.0%
<i>K. marxianus</i> ( <i>C. kefyr</i> )	2	2	0	0	0.0%	1	1	0	0.0%
<i>C. lusitaniae</i>	1	0	1	0	0.0%	0	0	0	0.0%
Total	8	56	1	1	1.7%	4	14	0	0.0%

**Table S2.** Univariate analysis of possible risk factors associated with 30-day mortality. Data are presented as percentages unless otherwise specified.  
CLABSI: Central Line-Associated Bloodstream Infection. \*:p-values<0.05

Characteristic	Alive at 30 days	Dead at 30 days	p-value
Age (years, mean ± SD)	61.4 ± 18.6	70 ± 14.9	<b>0.002*</b>
Female	47.90	49.50	0.848
Year of admission			
2004-2009	19.2	15.4	
2010-2014	45.2	30.8	0.062
2015-2019	35.6	53.8	
Critical care unit	27.4	64.8	<b>&lt;0.001*</b>
Diabetes mellitus	27.4	30.8	0.637
Hemodialysis patient	13.7	20.9	0.231
End-stage renal disease	6.8	5.5	0.753
Acute kidney injury	6.8	15.4	0.09
Solid organ transplant recipient	0.0	2.2	0.503
Stem cell transplant recipient	4.1	4.40	1.00
Auto-transplant	33.3	50.0	
Allo-transplant	33.3	50.0	1.00
Both	33.3	0.0	
Neutropenic	6.8	17.6	<b>0.041*</b>
1000-1500/mm <sup>3</sup>	20.0	12.5	
500-1000/mm <sup>3</sup>	0.0	37.5	0.351
<500/mm <sup>3</sup>	80.0	50.0	
Immunocompromised	56.2	70.3	0.060
Current steroids	16.4	25.3	0.170

Congenital syndromes	2.7	1.1	0.586
Other immunosuppressants	2.7	6.6	0.301
Malignancy	46.6	53.8	0.355
Non-hematologic	23.5	24.5	
Hematologic	73.5	69.4	0.926
Both	2.9	6.1	
Chemotherapy	24.7	22.0	0.686
<1 week from infection	33.3	20.0	0.468
Immunotherapy			
Abdominal surgery within 30 days	31.5	23.1	0.226
Antibiotics within 30 days	95.9	93.4	0.732
Antifungal history	23.3	28.6	0.445
Fluconazole	19.2	18.7	0.936
Voriconazole	1.4	1.1	1.00
Amphotericin B	1.4	3.3	0.63
Caspofungin	1.4	5.5	0.227
Anidulafungin	0.0	2.2	0.503
Duration of previous antifungal (days, mean ± SD)	9.13 ± 12.2	16.7 ± 17.9	0.152
Parenteral nutrition	27.4	25.3	0.759
Central venous catheter	63.0	75.8	0.075
Source			
CLABSI	27.4	19.8	0.251
Gastrointestinal translocation	35.6	33.0	0.722
Urinary tract infection	13.7	6.6	0.128
Unknown	19.2	38.5	<b>0.007*</b>
Other	1.4	1.1	1.00
Candida non-albicans	61.6	63.7	0.783
Echocardiography	63.0	46.2	<b>0.031*</b>
Ophthalmic exam	28.8	14.3	<b>0.023*</b>
Empiric therapy			
Fluconazole	40.6	30.9	0.236
Voriconazole	1.4	1.5	1.00
Itraconazole	4.3	2.9	1.00
Caspofungin	27.5	29.4	0.808
Anidulafungin	7.2	19.1	<b>0.04*</b>
Micafungin	10.1	8.8	0.792
Lipid formulation of amphotericin B	8.7	7.4	0.772
Central venous catheter removed	91.3	53.6	<b>&lt;0.001*</b>
Tip sent for culture	88.1	78.4	0.245