

**Table S1.** Local treatment and evolution of simple CL cases.

Local Therapy 24 (59%)	N=24	Effectiveness 3 months	Rescue Treatment	Effectiveness 12 months
Cryotherapy	8 (33%)	Cured		Cured
Intralesional Sb <sup>v</sup>	5 (21%)	Cured		Cured
Combination previous	8 (33%)	3 (13%) treatment failure	1 Imiquimod 5% 2 Fluconazol	Cured Cured
Surgery	1 (4%)	Cured		Cured
Observation	2 (8%)	Cured		Cured
		1 (4%) Lost follow-up		1 (4%) Lost follow-up

**Table S2.** Age, comorbidities, concomitant medication, type and grade of adverse events presented in the cohort of complex CL treated with L-AmB.

Patient	Age	Comorbidity	Concomitant Medication	Adverse event	Grade	Finish Treatment	Hospitalization
1	49	Arterial hypertension Psoriatic arthritis	Antihypertensive Infliximab	Fever Nausea Diarrhea	III	Yes	No
2	66	Inflammatory bowel disease	Infliximab	Infusional reaction Acute kidney failure	IV	Yes	Yes
3	84	Arterial hypertension	Antihypertensive	Acute kidney failure	III	Yes	No
4	39	Inflammatory bowel disease	Infliximab Azathioprine	Nausea/Vomiting Acute kidney failure	IV	No	Yes
5	55	-	-	Infusional reaction	III	Yes	No
6	66	Inflammatory bowel disease	Infliximab	Infusional reaction Acute kidney failure	III	Yes	No
7	8	-	Primary CD4 deficiency	Nausea/Vomiting	III	No	No
8	11	-	-	Nausea/Vomiting	III	No	No
9	43	-	-	-	-	Yes	No
10	69	Psoriatic arthritis	Golimumab	-	-	Yes	No
11	45	-	-	-	-	Yes	No
12	73	Arterial hypertension Dyslipidemia Psoriasis	Antihypertensive Hypolipemiant Adalimumab	-	-	Yes	No
13	36	Primary CD4 deficiency	-	-	-	Yes	No
14	42	-	-	-	-	Yes	No
15	6	-	-	-	-	Yes	No
16	31	-	-	-	-	Yes	No