

Table S4. Summary of cross agglutinin absorption tests (CAAT) performed on strain 782.

Antiserum	Antigen	Cross- agglutination (%)	CAAT, live antigen ( %)	CAAT, dead antigen ( %)
<b>Bafani</b>	782	6.25	Nil	Nil
<b>Benjamini</b>	782	3.13	Nil	Nil
<b>Malaya H6</b>	782	< 3.13	Nil	Nil
<b>Galtoni</b>	782	6.25	Nil	Nil
<b>Kuwait</b>	782	3.13	Nil	Nil
<b>Vleermus qoc</b>	782	6.25	Nil	Nil
<b>Qunjian</b>	782	3.13	Nil	Nil
<b>Hond Utrecht</b>	782	50	25	25
<b>Yonsis</b>	782	50	25	25
<b>Portlandvere</b>	782	50	25	50
<b>Sumneri</b>	782	50	25	25
<b>Kamituga</b>	782	12/62*	Nil	Nil
<b>782</b>	Kamituga	12/62*	100	50
<b>Broomi patane</b>	782	50	0.2	0.2
<b>782</b>	Broomi patane	50	6.3	6.3
<b>782</b>	Broomi patane	50	12.5	12.5
<b>Broomi patane</b>	782	50	3.1	3.1
<b>782</b>	Broomi patane	50	12.5	12.5
<b>782</b>	Broomi patane	50	12.5	12.5
<b>Bindjei</b>	782	50	0.8	< 0.4
<b>782</b>	Bindjei	50	6.3	0.4
<b>782</b>	Bindjei	50	3.1	6.3
<b>Bindjei</b>	782	50	3.1	1.6
<b>782</b>	Bindjei	50	1.6	1.6

\*Results were inconclusive, as repeat measures yielded both values.