

VP1 78% Nucleotide Cutoff Frequency of Identity 400-300-300-200-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-100-1 Nucleotide Percent Identity Inter-genotype Intra-genotype






Figure S1. Nucleotide pairwise identity frequency graphs of rotavirus B genes. The cutoff values are represented by

