

Table S2 Resistance to 0.3% bile salts at 15°C and 30°C of the examined *Lp. plantarum* strains (23V, 33V, 36V, 37V, 64V, 65V, 66V, 67V, 68V, 73V). The data (mean ± SD; n=3) were expressed as log CFU/mL. Different lowercase letters (a–e) in each row indicate significant differences ($p < 0.05$).

Temperature (°C)	Time (h)	<i>Lp. plantarum</i> strains									
		23V	33V	36V	37V	64V	65V	66V	67V	68V	73V
15	3	8.83±0.04 ^a	8.85±0.05 ^a	8.72±0.05 ^b	8.75±0.04 ^b	8.71±0.01 ^b	8.27±0.02 ^c	8.62±0.01 ^c	8.80±0.03 ^a	8.33±0.02 ^d	8.31±0.03 ^c
	24	8.71±0.05 ^a	8.62±0.03 ^a	8.55±0.04 ^b	8.11±0.05 ^d	8.20±0.01 ^d	8.25±0.04 ^c	8.33±0.02 ^c	8.58±0.03 ^b	8.12±0.04 ^d	8.08±0.07 ^e
30	3	8.84±0.06 ^a	8.11±0.03 ^c	8.79±0.01 ^a	8.79±0.03 ^a	8.17±0.02 ^c	8.15±0.01 ^c	8.14±0.03 ^c	8.68±0.01 ^b	8.14±0.04 ^c	8.73±0.02 ^b
	24	8.34±0.06 ^b	8.36±0.02 ^b	8.78±0.01 ^a	8.32±0.02 ^b	8.35±0.04 ^b	8.34±0.04 ^b	8.10±0.04 ^c	8.06±0.03 ^c	8.07±0.03 ^c	8.36±0.01 ^b