



Supplementary Figure S1. Kinetics of infections in childhood allogeneic and autologous hematopoietic stem cell transplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is represented with a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.




(a). Kinetics of infections in patient 1 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is represented with a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.




(b). Kinetics of infections in patient 2 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 3 allo-HSCT




(c). Kinetics of infections in patient 3 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 4 allo-HSCT




(d). Kinetics of infections in patient 4 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 5 allo-HSCT




(e) Kinetics of infections in patient 5 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 6 allo-HSCT




(f) Kinetics of infections in patient 6 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 7 allo-HSCT




(g) Kinetics of infections in patient 7 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 8 allo-HSCT




(h) Kinetics of infections in patient 8 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 9 allo-HSCT




(i) Kinetics of infections in patient 9 who received allogeneic hematopoietic stem cell transplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is represented with a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 10 allo-HSCT




(j) Kinetics of infections in patient 10 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is represented with a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 11 allo-HSCT




(k) Kinetics of infections in patient 11 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 12 allo-HSCT




(l) Kinetics of infections in patient 12 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 13 allo-HSCT




(m) Kinetics of infections in patient 13 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 14 allo-HSCT




(n) Kinetics of infections in patient 14 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 15 allo-HSCT




(o) Kinetics of infections in patient 15 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 16 allo-HSCT




(p) Kinetics of infections in patient 16 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 17 allo-HSCT




(q) Kinetics of infections in patient 17 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 18 allo-HSCT




(r) Kinetics of infections in patient 18 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 19 allo-HSCT




(s) Kinetics of infections in patient 19 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 20 allo-HSCT




(t) Kinetics of infections in patient 20 who received allogeneic hematopoietic stem celtransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 21 allo-HSCT




(u) Kinetics of infections in patient 21 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 22 allo-HSCT




(v) Kinetics of infections in patient 22 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 23 allo-HSCT




(w) Kinetics of infections in patient 23 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 24 allo-HSCT




(x) Kinetics of infections in patient 24 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 25 allo-HSCT




(y) Kinetics of infections in patient 25 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 26 allo-HSCT




(z) Kinetics of infections in patient 26 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 27 allo-HSCT




(aa) Kinetics of infections in patient 27 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 28 allo-HSCT




(ab) Kinetics of infections in patient 28 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 29 allo-HSCT




(ab) Kinetics of infections in patient 29 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 30 allo-HSCT




(ad) Kinetics of infections in patient 30 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 31 allo-HSCT




(ae) Kinetics of infections in patient 31 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 32 allo-HSCT




(af) Kinetics of infections in patient 32 who received allogeneic hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 1 auto-HSCT




(A) Kinetics of infections in patient 1 who received autologous hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 2 auto-HSCT




(B) Kinetics of infections in patient 2 who received autologous hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 3 auto-HSCT




(C) Kinetics of infections in patient 3 who received autologous hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 4 auto-HSCT




(D) Kinetics of infections in patient 4 who received autologous hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma

Patient 5 auto-HSCT




(E) Kinetics of infections in patient 5 who received autologous hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 6 auto-HSCT




(F) Kinetics of infections in patient 6 who received autologous hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 7 auto-HSCT




(G) Kinetics of infections in patient 7 who received autologous hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Patient 8 auto-HSCT



(H) Kinetics of infections in patient 8 who received autologous hematopoietic stem celltransplant (HSCT). The follow up is shown in the X axis and the viral load in the Y axis. Each virus is representedwith a specific color. Viral loads are expressed as DNA copies/ ug DNA in leukocytes or ml plasma.

Supplementary Figure S2. Viral loads for HCMV in patients with allogeneic and autologous hematopoietic stem cell transplantation (HSCT). Viral loads are expressed as DNA copies/ ug of DNA in leukocytes or ml of plasma. The red line indicates the clinical threshold used at the Children's Hospital of Mexico.



Supplementary Table S1. Pre-transplant serology for donors and recipients. Patients (n=40).

Type of transplant	Pre-transplant EBV serology		Pre-transplant HCMV serology		
	Donor	Recipient	Donor	Recipient	
Allogeneic (N=32)	Patient 1 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: + IgM: -
	Patient 2 Allo-HSCT	Without data	Without data	Without data	IgG: + IgM: -
	Patient 3 Allo-HSCT	IgG: + IgM: -	IgG: + IgM: -	IgG: + IgM: -	IgG: + IgM: -
	Patient 4 Allo-HSCT	IgG: + IgM: -	IgG: + IgM: -	IgG: + IgM: -	IgG: + IgM: -
	Patient 5 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 6 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 7 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: + IgM: -
	Patient 8 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: + IgM: -
	Patient 9 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 10 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: + IgM: -
	Patient 11 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 12 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 13 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: + IgM: -
	Patient 14 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 15 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: + IgM: -
	Patient 16 Allo-HSCT	IgG: + IgM: -	IgG: + IgM: -	IgG: + IgM: -	IgG: + IgM: -
	Patient 17 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 18 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: + IgM: -
	Patient 19 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 20 Allo-HSCT	Without data	IgG: - IgM: -	Without data	IgG: + IgM: -
	Patient 21 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 22 Allo-HSCT	IgG: + IgM: -	IgG: - IgM: -	IgG: + IgM: -	IgG: - IgM: -
	Patient 23 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 24 Allo-HSCT	Without data	Without data	IgG: - IgM: -	Without data
	Patient 25 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: + IgM: -
	Patient 26 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 27 Allo-HSCT	Without data	IgG: + IgM: -	Without data	IgG: Ind IgM: -
	Patient 28 Allo-HSCT	IgG: + IgM: -	Without data	IgG: - IgM: -	Without data
	Patient 29 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 30 Allo-HSCT	Without data	IgG: + IgM: -	IgG: + IgM: -	IgG: + IgM: -
	Patient 31 Allo-HSCT	Without data	Without data	Without data	Without data
	Patient 32 Allo-HSCT	Without data	Without data	Without data	Without data
Autologous (N=8)	Patient 1 Auto-HSCT	IgG: - IgM: -		IgG: - IgM: -	
	Patient 2 Auto-HSCT	IgG: + IgM: -		IgG: - IgM: -	
	Patient 3 Auto-HSCT	IgG: + IgM: -		IgG: + IgM: -	
	Patient 4 Auto-HSCT	IgG: + IgM: -		IgG: + IgM: -	
	Patient 5 Auto-HSCT	IgG: + IgM: -		IgG: + IgM: -	

Patient 6 Auto-HSCT	IgG: + IgM: -	IgG: + IgM: -
Patient 7 Auto-HSCT	IgG: + IgM: -	IgG: + IgM: -
Patient 8 Auto-HSCT	Without data	Without data

Ind means an indeterminate result. Patients who received autologous transplantation receive their own stem cells.

Supplementary Table S2. Leukocyte counts in blood samples.

Patient	Sample ID	Pre/Post transplant	Follow up days	Total of leukocytes (millions)
1 allo-HSCT	1.00	Pre-transplant	-5	0.8
	1.03	Post-transplant	21	16.2
	1.04		35	19.4
	1.05		49	12.6
	1.06		68	16.5
	1.07		78	3.6
	1.09		110	2.1
	1.10		138	4.0
	1.12		160	13.8
	1.13		182	5.4
2 allo-HSCT	2.00	Post-transplant	24	1.2
	2.01		37	9.7
	2.02		52	6.5
	2.03		68	13.5
	2.04		85	8.2
	2.05		92	3.3
	2.06		106	3.2
	2.09		155	11.1
	2.10		162	8.6
	2.11		187	9.7
	2.13		207	12.7
	2.14		241	2.1
3 allo-HSCT	3.00	Pre-transplant	-7	8.5
	3.01	Post-transplant	7	11.3
	3.02		21	8.9
	3.03		30	0.7
	3.04		62	0.7
	3.07		86	1.2
4 allo-HSCT	4.00	Pre-transplant	-10	8.0
	4.02	Post-transplant	20	10.3
	4.05		74	7.7
	4.06		81	8.2
	4.07		113	3.6

	4.09		151	0.7
	4.10		165	2.5
	4.11		172	0.0
	4.12		193	2.9
	4.13		207	2.7
	4.15		280	2.3
	4.16		292	6.1
	4.17		330	4.9
	4.18		340	1.4
	4.19		351	2.8
5 allo-HSCT	6.00	Pre-transplant	-7	4.8
	6.02	Post-transplant	22	2.4
	6.04		48	2.9
	6.06		80	9.0
	6.07		94	1.4
	6.08		104	0.0
	6.09		132	4.6
	6.10		143	2.0
	6.11		157	2.0
	6.13		212	3.1
	6.14		233	1.1
	6.15		259	5.1
	6.16		294	1.0
	6.19		350	0.9
	6.20		364	2.8
6 allo-HSCT	9.00	Pre-transplant	-7	3.1
	9.02	Post-transplant	13	4.3
	9.03		21	5.2
	9.04		25	7.4
	9.05		75	7.4
	9.06		95	1.4
	9.07		106	2.5
	9.08		120	7.0
7 allo-HSCT	10.00	Pre-transplant	-8	0.2
	10.02	Post-transplant	21	7.8
	10.03		36	7.3
	10.04		49	7.3
	10.05		60	3.2
	10.06		88	6.9

	10.07		117	2.2
	10.09		152	3.4
	10.10		189	4.2
	10.11		202	0.7
	10.12		249	3.2
	10.13		277	2.2
	10.14		291	2.6
	10.15		305	2.1
	10.16		320	2.0
	10.17		344	3.2
8 allo-HSCT	12.00	Pre-transplant	-12	0.6
	12.02	Post-transplant	21	11.6
	12.03		33	16.1
	12.04		51	7.6
	12.05		65	7.9
	12.06		79	5.2
	12.07		89	4.5
	12.08		104	11.2
	12.09		110	7.5
	12.10		126	7.5
	12.11		138	3.4
	12.12		159	17.7
	12.13		173	10.4
	12.15		222	3.9
	12.16		245	11.9
	12.17		264	7.8
	12.18		279	19.9
	12.19		293	16.6
	12.20		314	1.8
	12.21		328	2.1
	12.22		342	2.5
9 allo-HSCT	13.00	Pre-transplant	-9	1.6
	13.02	Post-transplant	19	7.6
	13.03		32	2.7
	13.04		53	1.5
	13.05		68	1.8
	13.06		74	1.8
	13.07		95	2.0
	13.08		111	71.6

	13.09		131	13.3
	13.10		152	2.8
	13.11		168	9.6
	13.12		186	2.5
	13.13		202	5.5
	13.14		221	3.5
	13.15		236	2.5
	13.16		249	3.9
	13.17		271	2.6
	13.18		299	2.8
	13.19		319	5.4
10 allo-HSCT	14.00	Pre-transplant	-8	0.1
	14.02	Post-transplant	21	0.1
	14.03		31	11.4
	14.09		115	6.3
	14.10		136	0.4
	14.11		150	1.0
	14.12		168	2.2
	14.13		192	0.1
	14.14		206	2.8
	14.15		220	7.5
	14.16		235	0.1
	14.17		248	3.0
	14.18		270	0.1
	14.19		291	2.1
	14.20		327	3.9
11 allo-HSCT	15.01		7	0.1
	15.02		54	1.5
	15.03		65	3.1
	15.05		85	2.0
12 allo-HSCT	19.00	Pre-transplant	-7	5.3
	19.03	Post-transplant	35	3.5
	19.04		48	12.3
	19.05		61	2.3
	19.07		75	7.3
	19.09		121	1.4
	19.10		138	1.1
	19.11		152	4.8
	19.12		173	2.3

	19.13		187	2.4
	19.14		201	2.5
	19.15		236	3.6
13 allo-HSCT	20.00	Pre-transplant	-8	0.8
	20.03	Post-transplant	21	3.3
	20.05		55	1.2
	20.06		66	6.9
	20.07		76	2.3
14 allo-HSCT	22.00	Pre-transplant	-8	0.4
	22.01	Post-transplant	7	4.8
	22.02		26	9.1
	22.03		40	36.0
	22.04		46	1.7
	22.06		78	0.8
	22.08		102	4.1
	22.09		116	8.4
	22.10		130	5.9
	22.11		158	2.6
	22.12		172	44.8
	22.13		200	6.0
	22.14		215	0.9
	22.15		228	4.2
	22.16		252	21.8
15 allo-HSCT	23.00	Pre-transplant	-7	1.2
	23.01	Post-transplant	0	19.1
	23.03		21	17.2
	23.04		38	5.0
	23.05		56	1.8
16 allo-HSCT	24.00	Pre-transplant	-6	1.8
	24.01	Post-transplant	7	31.7
	24.02		21	3.1
	24.03		36	0.5
	24.04		50	20.1
	24.05		61	1.5
	24.06		75	1.7
	24.07		88	1.9
	24.08		102	5.9
17 allo-HSCT	25.00	Pre-transplant	-8	0.8
	25.02	Post-transplant	21	6.6

	25.03		33	0.8
	25.04		46	2.1
	25.05		61	2.6
	25.06		75	4.0
	25.07		95	12.6
	25.08		116	5.8
	25.09		131	4.2
	25.10		151	13.1
	25.11		183	5.1
	25.12		201	6.1
	25.13		214	3.8
	25.14		228	10.8
	25.15		242	5.1
	25.16		257	4.4
	25.17		271	7.0
	25.18		284	2.4
	25.19		299	3.1
	25.20		319	4.5
	25.21		334	4.4
18 allo-HSCT	26.00	Post-transplant	2	0.8
	26.02		30	4.0
	26.03		44	6.8
	26.04		62	1.9
	26.05		83	0.5
	26.06		96	2.9
	26.07		112	2.2
	26.08		132	5.4
	26.09		153	2.1
	26.10		170	3.5
	26.11		188	3.0
	26.12		202	0.8
	26.13		215	0.1
19 allo-HSCT	27.00	Pre-transplant	-7	5.8
	27.01	Post-transplant	11	0.4
	27.02		29	12.3
	27.03		41	6.4
	27.04		53	5.6
	27.05		67	3.7
	27.06		88	8.4

20 allo-HSCT	28.00	Pre-transplant	-7	6.4
	28.03	Post-transplant	48	40.0
	28.04		63	4.3
21 allo-HSCT	29.00		17	2.4
	29.02		21	5.6
	29.03		34	5.0
	29.04		48	0.6
	29.05		62	4.2
	29.06		76	13.4
	29.07		89	2.7
	29.08		104	1.3
	29.09		119	4.6
	29.10		132	2.7
	29.11		160	1.4
	29.12		174	3.7
	29.13		188	2.5
	29.14		202	2.5
	29.15		216	1.6
	29.16		230	3.4
	29.17		251	6.4
	29.18		272	3.6
	29.19		309	5.5
	29.20		351	3.5
22 allo-HSCT	30.00	Pre-transplant	-7	0.5
	30.01	Post-transplant	7	8.7
	30.02		21	11.8
	30.03		47	4.1
	30.04		61	3.2
	30.05		75	1.3
	30.06		90	6.8
	30.07		124	3.4
	30.08		133	3.4
	30.09		166	1.6
	30.10		180	1.7
	30.11		201	0.9
	30.12		236	3.2
	30.13		250	5.1
	30.14		272	5.5
	30.15		293	2.0

23 allo-HSCT	31.00	Pre-transplant	-2	0.2
	31.01	Post-transplant	14	2.2
	31.02		21	11.5
	31.03		35	4.7
	31.04		54	8.1
24 allo-HSCT	33.00	Pre-transplant	-9	3.0
	33.02	Post-transplant	21	9.7
	33.03		33	0.2
	33.04		47	0.5
	33.05		68	2.9
	33.06		89	2.9
25 allo-HSCT	34.00	Pre-transplant	-8	3.1
	34.02	Post-transplant	20	4.3
	34.03		36	3.5
	34.04		55	2.6
	34.05		69	2.0
	34.06		97	0.4
	34.07		112	1.2
	34.08		126	0.6
	34.09		146	0.5
	34.10		160	0.2
	34.11		191	2.7
	34.12		209	1.9
	34.13		223	1.2
	34.14		251	1.8
	34.15		322	1.5
	34.16		339	2.3
26 allo-HSCT	37.00	Post-transplant	6	1.4
	37.02		27	0.8
	37.03		46	1.1
	37.04		60	0.5
	37.05		82	0.3
	37.07		117	1.1
	37.08		131	2.0
	37.09		145	0.5
	37.10		159	3.3
	37.11		174	1.6
	37.12		187	5.3
	37.13		202	3.7

	37.14		215	0.7
	37.15		231	3.1
	37.16		287	0.6
27 allo-HSCT	40.00	Pre-transplant	-7	5.5
	40.03	Post-transplant	35	4.6
	40.04		49	1.1
	40.05		63	2.6
	40.06		77	0.4
	40.07		91	3.4
	40.08		104	2.0
	40.09		125	2.3
	40.10		181	1.3
	40.11		259	3.3
	40.12		293	6.0
	40.13		307	5.4
	40.14		323	1.9
	40.15		359	3.3
28 allo-HSCT	41.00	Pre-transplant	-7	3.5
	41.01	Post-transplant	4	0.3
	41.02		20	2.6
	41.03		40	5.6
	41.04		52	1.3
	41.05		68	3.3
	41.06		81	1.8
	41.07		95	1.4
	41.08		109	1.1
	41.09		171	3.6
	41.10		199	4.7
	41.11		227	3.8
	41.12		245	4.4
	41.13		292	7.0
	41.14		319	6.5
29 allo-HSCT	42.00	Pre-transplant	-9	0.9
	42.02	Post-transplant	21	0.8
	42.03		34	2.9
	42.04		55	2.8

	42.05		69	3.7
30 allo-HSCT	44.00	Pre-transplant	-7	5.5
	44.02	Post-transplant	25	1.7
	44.03		35	1.8
	44.04		80	2.1
	44.05		101	3.8
	44.06		136	2.6
	44.07		171	7.5
	44.08		185	2.4
	44.09		199	4.2
	44.10		227	9.4
	44.11		238	5.0
31 allo-HSCT	45.00		0	3.3
	45.02		21	1.5
	45.03		35	0.2
	45.04		50	15.6
	45.05		61	15.0
	45.06		75	6.8
	45.07		85	1.5
	45.08		111	1.0
	45.09		132	2.3
	45.10		146	7.4
	45.11		169	0.8
	45.12		180	0.5
	45.13		202	1.2
	45.14		211	2.3
32 allo-HSCT	46.00	Pre-transplant	-4	0.5
	46.01	Post-transplant	7	0.5
	46.02		22	2.4
	46.03		34	1.8
	46.04		48	0.8
	46.05		63	4.0
	46.06		97	8.7
	46.07		111	4.2
	46.08		125	1.6
	46.09		134	0.1
	46.10		152	6.7
1 auto-HSCT	5.00	Pre-transplant	-8	5.4
	5.01	Post-transplant	1	1.3

	5.03		21	7.5
	5.04		56	5.7
2 auto-HSCT	7.00	Pre-transplant	-7	6.7
	7.02	Post-transplant	19	6.7
	7.03		24	1.2
3 auto-HSCT	8.00	Pre-transplant	-7	4.3
	8.03	Post-transplant	11	0.8
	8.04		31	16.3
	8.05		87	4.9
4 auto-HSCT	17.00	Pre-transplant	-7	1.6
	17.02	Post-transplant	22	9.5
5 auto-HSCT	32.00	Pre-transplant	-7	1.0
	32.01	Post-transplant	15	0.8
	32.02		21	1.8
	32.03		41	0.6
	32.04		57	8.7
	32.05		170	12.9
6 auto-HSCT	35.00	Pre-transplant	-7	6.0
	35.01	Post-transplant	4	1.1
	35.02		21	2.4
	35.03		32	2.4
	35.04		60	7.6
7 auto-HSCT	38.00	Pre-transplant	-7	0.6
	38.01	Post-transplant	7	5.5
	38.02		41	5.5
8 auto-HSCT	43.00	Pre-transplant	-7	1.7
	43.01	Post-transplant	7	1.8
	43.02		21	2.5