

Supplementary Table S5. Differences in resistance of ExPEC (n=14) versus non-ExPEC (n=126) isolates.

Antimicrobial agent	No. (%) of isolates resistant in		P-value by Chi-square test
	ExPEC	Non-ExPEC	
Cefepime	7 (50.0)	63 (50.0)	1.00
Cephalothin	14 (100.0)	117 (92.6)	0.600
Ceftazidime	8 (57.1)	59 (46.8)	0.58
Cefoxitin	5 (35.7)	25 (19.8)	0.18
Cefotaxime	14 (100.0)	105 (83.3)	0.13
Ceftriaxone	8 (57.1)	34 (27.0)	0.03
Aztreonam	7 (50.0)	34 (27.0)	0.12
Ertapenem	1 (7.1)	3 (2.4)	0.31
Meropenem	0 (0.0)	2 (1.6)	1.00
Imipenem	1 (7.1)	10 (7.9)	1.00
Gentamicin	1 (7.1)	6 (4.8)	0.53
Streptomycin	8 (57.1)	70 (55.6)	1.00
Amikacin	3 (21.4)	17 (13.5)	0.42
Amoxyclav	9 (64.3)	72 (57.1)	1.00
Ampicillin/Sulbactam	4 (28.6)	46 (36.5)	0.57
Piperacillin/Tazobactam	5 (35.7)	70 (55.6)	0.17
Ampicillin	10 (71.4)	97 (77.0)	0.74
Piperacillin	13 (92.9)	105 (83.3)	0.70
Trimethoprim	5 (35.7)	55 (43.7)	0.78
Co-Trimoxazole	3 (21.4)	55 (43.7)	0.15
Tetracycline	7 (50)	74 (58.7)	0.58
Ciprofloxacin	8 (57.1)	38 (30.2)	0.07
Fosfomycin	1 (7.1)	3 (2.4)	0.35
Azithromycin	3 (21.4)	10 (7.9)	0.13
Chloramphenicol	4 (28.6)	23 (18.3)	0.47
Sulphafurazole	6 (42.9)	60 (47.6)	0.79
Polymyxin B	4 (28.6)	21 (16.7)	0.28
Colistin	2 (14.3)	2 (1.6)	0.05