

Table S1. Susceptibility of *E. coli* isolated in cloacal and skin sample.

		Sample			
		Skin		Cloacal swabs	
		Count	%	Count	%
AMP	Resistant	158a	78.2%	153a	75.0%
	Intermediate	15a	7.4%	13a	6.4%
	Susceptible	29a	14.4%	38a	18.6%
CAZ	Resistant	22a	10.9%	22a	10.8%
	Intermediate	10a	5.0%	15a	7.4%
	Susceptible	170a	84.2%	167a	81.9%
CTX	Resistant	62a	30.7%	62a	30.4%
	Intermediate	4a	2.0%	13b	6.4%
	Susceptible	136a	67.3%	129a	63.2%
AMC	Resistant	74a	36.6%	71a	34.8%
	Intermediate	48a	23.8%	18b	8.8%
	Susceptible	80a	39.6%	115b	56.4%
NA	Resistant	83a	41.1%	94a	46.1%
	Intermediate	15a	7.4%	3b	1.5%
	Susceptible	104a	51.5%	107a	52.5%
CIP	Resistant	52a	25.7%	68a	33.3%
	Intermediate	26a	12.9%	41b	20.1%
	Susceptible	124a	61.4%	95a	46.6%
SXT	Resistant	83a	41.1%	80a	39.2%
	Intermediate	5a	2.5%	8a	3.9%
	Susceptible	114a	56.4%	116a	56.9%
TET	Resistant	136a	67.3%	142a	69.6%
	Intermediate	6a	3.0%	4a	2.0%
	Susceptible	60a	29.7%	58a	28.4%
CN	Resistant	22a	10.9%	26a	12.7%
	Intermediate	7a	3.5%	8a	3.9%
	Susceptible	173a	85.6%	170a	83.3%

Ampicillin (AMP), cefotaxime (CTX), ceftazidime (CAZ), amoxicillin/acid clavulanic (AMC), nalidixic acid (NA), ciprofloxacin (CIP), trimethoprim/sulfamethoxazole (SXT), tetracycline (TET) and gentamicin (CN)