

[illegible]

6	1	1	2	0	0	1	5
	1	1	2	0	0	1	5
	LIFUP_1						
	1	2	2	0	0	1	6
	4	5	0	0	0	1	10
	1	1	2	0	0	1	5
7							
	0	1	2	0	0	1	4
	0	1	2	1	0	1	5
	LIFUP_1						
	1	1	2	1	0	1	6
	0	1	2	1	0	1	5
	0	1	2	1	0	1	5
	0	1	2	1	0	1	5
	LIFUP_2						
	1	1	2	1	0	1	6
	1	1	2	1	0	1	6
	1	1	2	1	0	1	6
8	1	0	2	1	0	0	4
	1	0	2	1	0	0	4
	LIFUP_1						
	0	1	0	1	0	0	2
	0	1	2	1	0	0	4
9	1	0	1	1	0	2	5
	3	1	2	0	0	0	6
	LIFUP_1						
	1	2	1	1	0	1	6
	3	3	2	1	0	1	10
	3	3	2	1	0	1	10
10	1	1	2	0	0	1	5
	3	1	2	1	0	1	8
	LIFUP_1						
	0	1	2	1	0	0	4
	3	2	2	1	0	1	9
11	1	0	0	0	0	0	1
	0	0	1	0	0	0	1
	0	0	1	0	0	0	1
	LIFUP_1						
	0	0	1	1	0	0	2
	0	0	1	1	0	0	2

Supplementary Table S1. Here, all behavioral data is supplied for each subject. Though the number of assessments varies some between subjects, each subject received at least two CRS-R assessments prior to and following LIFU. These include, necessarily, 1 hour and 1 day prior to and following LIFU. Assessments are supplied chronologically. If additional assessments are present, they represent an assessment taken 1 week prior to and/or following LIFU.