



Supplementary Table 1. Correlations among markers of oxidative stress in Crohn's patients.

	FRAP	CO	TBARS	AGEs	AOPP
FRAP		0.2477 (39)	0.4830 (39)	0.2553 (39)	0.2863 (39)
		0.1284	0.0018	0.1167	0.0773
CO	0.2477 (39)		0.4100 (39)	0.2849 (39)	0.4789 (39)
		0.1284	0.0095	0.0788	0.0020
TBARS	0.4830 (39)	0.4100 (39)		0.3352 (39)	0.5298 (39)
		0.0018	0.0095	0.0370	0.0005
AGEs	0.2553 (39)	0.2849 (39)	0.3352 (39)		0.1256 (39)
		0.1167	0.0788	0.0370	0.4462
AOPP	0.2863 (39)	0.4789 (39)	0.5298 (39)	0.1256 (39)	
		0.0773	0.0020	0.0005	0.4462

Data reported in table are correlation coefficients, (sample size: number of complete data) and p-values.

Supplementary Table 2. Correlations among markers of oxidative stress in healthy volunteers.

	FRAP	CO	TBARS	AOPP	AGEs
FRAP		0.5149 (16)	0.1760 (16)	0.3106 (16)	0.1350 (16)
		0.0413	0.5144	0.2417	0.6180
CO	0.5149 (16)		0.3851 (16)	0.5335 (16)	0.1092 (16)
		0.0413	0.1407	0.0333	0.6873
TBARS	0.1760 (16)	0.3851 (16)		0.6053 (16)	0.2402 (16)
		0.5144	0.1407	0.0130	0.3703
AOPP	0.3106 (16)	0.5335 (16)	0.6053 (16)		-0.2030 (16)
		0.2417	0.0333	0.0130	0.4508
AGEs	0.1350 (16)	0.1092 (16)	0.2402 (16)	-0.2030 (16)	
		0.6180	0.6873	0.3703	0.4508

Data reported in table are correlation coefficients, (sample size: number of complete data) and p-values.