

Table S1. The condition of patients with UC before and after FMT according to the survey.

Patient	Sex	Age	Clinical remission or re response at week 8	Improvements after FMT (interview)	Total duration of improvements, months	Current number of defecations per day*		Presence of blood in the stool at the time of survey*		Suggested cause of recurrence	Supporting medications used
						Before FMT	After FMT	Before FMT	After FMT		
1	F	29	rem	+	18**	2-3	1-2	+	-		Mesalazine
2	F	42	rem	+	6	1-2	1-2	+	-	Severe stress	Azathioprine, Mesalazine
3	M	57	rem	+	24	5	2-3	+	-	Surgery (cholecystostomy) or treatment of pneumonia with a triple course of antibiotics	
4	M	34	res	+	48**	7-8	Up to 3	-	-		Sulfasalazine, Mesalazine
6	F	34	res	-	2	4	2	+	+		Mesalazine
8	M	42	rem	+	6**	2-3	1	+	-		Mesalazine
9	F	25	rem	+	6**	1-2	1-2	-	-		Mesalazine, Kolofort, Prednisolone, 6-Mercaptopurine, Zakofalk
12	M	36	rem	+	24**	1	1-2	-	-		Mesalazine
14	F	34	res	-	2	2-8	Up to 16	+	+		Mesalazine, Zakofalk
15	M	46	res	+	4	1	1	+	-		
17	M	36	rem	+	4	1	1	-	+	Diet violation	Mesalazine
19	F	26	rem	+	24**	Up to 4	1	+	-		Mesalazine, Trimedat
20	M	34	rem	+	6	Up to 5	2-12	+	+	Preparation for sigmoidoscopy	GCS, Mesalazine

* At the time of the survey. ** No recurrences of the disease at the time of the survey.

Table S2. Shannon indices for patients before and after FMT

Patient	Shannon index before FMT	Shannon index after FMT
1	5.03	3.92
2	6.16	3.40
3	5.90	5.76
4	2.77	5.94
5	3.27	3.37

6	3.28	5.10
7	5.03	3.33
8	5.82	6.14
9	3.84	6.49
10	3.08	5.62
11	4.86	5.57
12	4.44	5.63
13	4.72	5.77
14	5.76	5.55
15	4.83	6.16
16	5.23	5.78
17	4.63	5.94
18	4.88	4.92
19	4.87	5.59
20	5.77	6.49