



**Figure S1.** The change in N<sub>2</sub>O concentration after all bottles were opened and placed in a fuming cupboard for 5 minutes. The N<sub>2</sub>O concentration at 1 min indicated the initial concentration.

**Table S1.** Primers and the thermal cycling conditions of functional genes related to N<sub>2</sub>O.

Gene	Name	Sequence	Thermal profile	Reference
AOA <i>amoA</i>	CrenamoA23f	ATGGTCTGGCTWAGACG	15 s-95°C, 53°C-45 s, 72°C-45 s, 83°C-15 s, 45 cycles	Nicol and Schleper, 2006
	CrenamoA616r	GCCATCCATCTGTATGTCCA		
AOB <i>amoA</i>	<i>amoA</i> -1F	GGGGTTTCTACTGGTGGT	15 s-95°C, 54°C-40 s, 72°C-45 s,	Rotthauwe et al., 1997
	<i>amoA</i> -2R	CCCCTCKGSAAAGCCTCTTC	84°C-15 s, 45 cycles	
<i>nirK</i>	<i>nirK</i> -F1aCu	ATCATGGTSCTGCCGCG	10 s-95°C, 53°C-45 s, 72°C-45 s,	Henry et al., 2004
	<i>nirK</i> -R3Cu	GCCTCGATCAGRTTGTGGTT	86°C-15 s, 45 cycles	
<i>nirS</i>	<i>nirS</i> -Cd3aF	GTSAACGTSAAAGGARACSGG	15 s-95°C, 50°C-45 s, 72°C-45 s,	Throbäck et al., 2004
	<i>nirS</i> -R3cd	GASTTCGGRTGSGTCTTGA	88°C-15 s, 50 cycles	
<i>nosZ</i>	<i>nosZ</i> -F	CGYTGTTCMTCGACAGCCAG	15 s-95°C, 50°C-30 s, 72°C-30 s, 83°C-15 s, 55 cycles	Scala and Kerkhof, 1998
	<i>nosZ</i> -1662R	CGSACCTTSTTGCCSTYGCG		

## References

- Henry, S., Baudoin, E., López-Gutiérrez, J.C., Martin-Laurent, F., Brauman, A., Philippot, L., 2004. Quantification of denitrifying bacteria in soils by nirK gene targeted real-time PCR. *Journal of Microbiological Methods* 59, 327–335. <https://doi.org/10.1016/j.mimet.2004.07.002>
- Nicol, G.W., Schleper, C., 2006. Ammonia-oxidising Crenarchaeota: important players in the nitrogen cycle? *Trends in Microbiology* 14, 207–212. <https://doi.org/10.1016/j.tim.2006.03.004>
- Rotthauwe, J.H., Witzel, K.P., Liesack, W., 1997. The ammonia monooxygenase structural gene amoA as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Applied and environmental microbiology* 63, 4704–4712. <https://doi.org/10.1128/AEM.63.12.4704-4712.1997>
- Scala, D.J., Kerkhof, L.J., 1998. Nitrous oxide reductase (*nosZ*) gene-specific PCR primers for detection of denitrifiers and three *nosZ* genes from marine sediments. *FEMS Microbiology Letters* 162, 61–68. <https://doi.org/10.1111/j.1574-6968.1998.tb12979.x>
- Throbäck, I.N., Enwall, K., Jarvis, Å., Hallin, S., 2004. Reassessing PCR primers targeting nirS, nirK and nosZ genes for community surveys of denitrifying bacteria with DGGE. *FEMS Microbiology Ecology* 49, 401–417. <https://doi.org/10.1016/j.femsec.2004.04.011>