

Light Conversion upon Photoexcitation of NaBiF₄:Yb³⁺/Ho³⁺/Ce³⁺ Nanocrystalline Particles

Enrico Trave ^{1,*}, Michele Back ^{1,*}, Davide Pollon ¹, Emmanuele Ambrosi ¹ and Leonardo Puppulin ^{1,2}

¹ Department of Molecular Sciences and Nanosystems, Università Ca' Foscari Venezia, Via Torino 155, 30172 Venice, Italy; pollon.davide@gmail.com (D.P.); emmanuele.kizito@gmail.com (E.A.); leonardo.puppulin@unive.it (L.P.)

² WPI Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kanazawa, Ishikawa 920-1192, Japan

* Correspondence: enrico.trave@unive.it (E.T.); michele.back@unive.it (M.B.)




Figure S1. Trend of cell volume vs. Ce³⁺ content; cell volume estimation was obtained from fitting of XRPD patterns reported in Figure 1(a).

Citation: Trave, E.; Back, M.; Pollon, D.; Ambrosi, E.; Puppulin, L. Light Conversion upon Photoexcitation of NaBiF₄:Yb³⁺/Ho³⁺/Ce³⁺ Nanocrystalline Particles. *Nanomaterials* **2023**, *13*, 672. <https://doi.org/10.3390/nano13040672>

Academic Editors: Vincenzo Vaiano

Received: 29 December 2022

Revised: 1 February 2023

Accepted: 5 February 2023

Published: 9 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure S2. FE-SEM images of Ce₀, Ce₂, Ce₅ and Ce₂₀ samples.




Figure S3. EDS spectrum of Ce_10 sample.




Figure S4. FE-SEM image and elemental maps of Ce_10 sample.