



Article

Comparison of Different Approaches to Surface Functionalization of Biodegradable Polycaprolactone Scaffolds

Elizaveta S. Permyakova ¹, Philipp V. Kiryukhantsev-Korneev ¹, Kristina Yu. Gudz ¹, Anton S. Konopatsky ¹, Josef Polčák, ^{2,3} Irina Y. Zhitnyak ⁴, Natalia A. Gloushankova ⁴, D. V. Shtansky ¹ and Anton M. Manakhov ^{5,*}

¹ National University of Science and Technology "MISIS", Leninsky prospect 4, Moscow, 119049, Russia
permyakova.elizaveta@gmail.com (E.S.P.); kiruhancev-korneev@yandex.ru (P.V.K-K.);
kristinagudz@mail.ru (K.Y.G.); konopatskiy@misis.ru (A.S.K.); shtansky@shs.misis.ru (D.V.S.)

² CEITEC—Central European Institute of Technology, Brno University of Technology, Purkyňova 123, Brno, Czech Republic; polcak@fme.vutbr.cz

³ Institute of Physical Engineering, Brno University of Technology, Technická 2896/2, 616 69 Brno, Czech Republic

⁴ N.N. Blokhin Russian Cancer Research Center, Kashirskoe shosse 24, Moscow 115478, Russia;
irishaz@mail.ru (I.Y.Z); natglu@hotmail.com (N.A.G)

⁵ Scientific Institute of Clinical and Experimental Lymphology—Branch of the ICG SB RAS, 2 Timakova str., 630060 Novosibirsk, Russian Federation

* Correspondence: ant-manahov@ya.ru; Tel.: +7-915-8494059

Content:

- 1) Additional FTIR spectra
- 2) Microscope image for cell quantification analysis

- 1) Additional FTIR spectra




Figure S1. The comparison of PCL-ref-SBF, PCL-COOH-SBF and PCL-TiCaPCON-SBF after immersion into SBF for 7 days for identification of phosphate ions peaks.




Figure S2. The example of the images for demonstration of the cell quantification.