

Table S1. Minimal inhibitory concentrations (MIC) of antibiotics and non-antibiotic agents against bacteria used in the study, µg /ml.

		<i>E.coli</i>	<i>P. aeru- ginosa</i>	<i>K. pneumonia</i>	<i>E. faecium</i>	<i>S. aureus</i>	<i>C. acnes</i>
Antibiotics ^a	Azythromycin	-	-	8.0	-	0.5	-
	Claritromycin	-	-	64.0	-	0.25	-
	Amoxiclav	-	-	8.0	-	0.13	-
	Gentamicin	1.0	0.25	0.25	1,0	0.25	-
Non-antibiotic agents ^b	Benzoyl peroxide	31.0	-	-	-	1250.0	80.0
	Chlorhexidine	16.0	1.0	-	-	1.0	1.0

^a Azithromycin dihydrate (Sigma-Aldrich, St. Louis, MO, USA), clarithromycin (Sigma-Aldrich, St. Louis, MO, USA), amoxiclav (Sandoz, Kundl, Austria; amoxicillin 875 mg, acid clavulanic 125 mg), gentamicin (Sigma-Aldrich, St.Louis, MO, USA). ^b Benzoyl peroxide (Saules Aptieka, Riga, Latvia), chlorhexidine (Saules aptieka, Riga, Latvia).