



Article Understanding the Effect of Conserved Charges on the Coalescence Sum Rule of Directed Flow

Kishora Nayak ^{1,*}, Shu-Su Shi ², and Zi-Wei Lin ³

- ¹ Department of Physics, Panchayat College, Sambalpur University, Bargarh 768028, India
- ² Key Laboratory of Quark & Lepton Physics (MOE) and Institute of Particle Physics, Central China Normal University, Wuhan 430079, China
- ³ Department of Physics, East Carolina University, Greenville, NC 27858, USA; linz@ecu.edu

* Correspondence: k.nayak1234@gmail.com

Abstract: Recently, the rapidity-odd directed flow (v_1) of produced hadrons $(K^-, \phi, \overline{p}, \overline{\Lambda}, \overline{\Xi}^+, \Omega^-,$ and $\overline{\Omega}^+$) has been studied. Several combinations of these produced hadrons, with very small mass differences but differences in the net electric charge (Δq) and net strangeness (ΔS) on the two sides, have been considered. A difference in v_1 between the two sides of these combinations (Δv_1) has been proposed as a consequence of the electromagnetic field produced in relativistic heavy-ion collisions, especially if Δv_1 increases with Δq . Our study is performed to understand the effect of the coalescence sum rule (CSR) on Δv_1 . We point out that the CSR leads to $\Delta v_1 = c_q \Delta q + c_S \Delta S$, where the coefficients c_q and c_S reflect the Δv_1 of produced quarks. Equivalently, one can write $\Delta v_1 = c_q \Delta q + c_B \Delta B$, involving the difference in the net baryon number ΔB , where the CSR gives $c_B = -3c_S$. We then propose two methods to extract the coefficients for the Δq and ΔS dependences of Δv_1 .

Keywords: directed flow; coalescence sum rule

1. Introduction

The properties of strongly interacting quark-gluon plasma, produced by relativistic heavy-ion collisions, can be studied using anisotropic flows, including the directed flow (v_1) [1–3]. v_1 has been found to be a sensitive probe of the equation of state of the produced medium [4,5]. The anisotropic flows of identified hadrons are expected to follow the coalescence sum rule (CSR) when the produced matter is initially in parton degrees of freedom and hadronizes via quark coalescence. The CSR states that the sum of constituent quarks' flow equals the corresponding flow of the hadron; it leads to the number-of-constituent-quark scaling and provides information about the particle production mechanism and partonic collectivity [6,7]. This paper is based on our recent study [8], which was motivated by Ref. [9].

A new method of testing the coalescence sum rule using the difference in v_1 in different combinations of produced hadrons $[K^-(\bar{u}s), \phi(s\bar{s}), \bar{p}(\bar{u}\bar{u}\bar{d}), \bar{\Lambda}(\bar{u}\bar{d}\bar{s}), \Xi^+(\bar{d}\bar{s}\bar{s}), \Omega^-(sss)$ and $\bar{\Omega}^+(\bar{s}\bar{s}\bar{s})]$ has been previously proposed [9]. These selected hadrons are all produced, consisting of \bar{u}, \bar{d}, s , and \bar{s} quarks. In contrast to the produced hadrons, transported hadrons such as π^{\pm} , p, and Λ receive contributions from initial-state (incoming) u and d quarks, along with the produced quarks, which complicates the interpretations of the CSR. Table 1 shows a selection of five independent hadron sets [8]. Sets 1–3 are identical, with the same quark contents on both sides, whereas sets 4–5 are non-identical. Indeed, different choices of five independent hadron sets can be made [8]. One can also obtain all ten sets from Table 1 in Ref. [9]; however, they are not all independent. In a given set, the difference in v_1 between the left and right sides (after including the weighting factors of each hadron as listed in Table 1) is termed Δv_1 . Similarly, the differences in the net electric charge (Δq or Δq_{ud}) and net strangeness (ΔS) are obtained.



Citation: Nayak, K.; Shi, S.-S.; Lin, Z.-W. Understanding the Effect of Conserved Charges on the Coalescence Sum Rule of Directed Flow. *Universe* 2024, *10*, 112. https:// doi.org/10.3390/universe10030112

Academic Editor: Carlos Pajares

Received: 29 November 2023 Revised: 6 February 2024 Accepted: 20 February 2024 Published: 1 March 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). As v_1 develops in the early stage of collisions, it is sensitive to the strong electromagnetic field produced by incoming protons in the two colliding nuclei. The correlation between the difference in v_1 on Δq has been considered as breaking the coalescence sum rule and being a consequence of the electromagnetic fields [9,10], especially when the difference in v_1 increases with Δq . This motivated us to critically examine the consequence of the coalescence sum rule on the difference in v_1 [8]. This paper is organized as follows. The derivation of the CSR relation is given in Section 2. Two methods for extracting the dependence of Δv_1 on Δq and ΔS are presented in Section 3. A summary is finally given in Section 4.

2. The CSR Relation for the Difference in v_1 of a Hadron Set

The relation between the v_1 of a hadron (H) and those of its constituent quarks from the CSR can be written as

$$v_{1,\mathrm{H}}(p_{\mathrm{T,H}}) = \sum_{j} v_{1,j}(p_{T,j}),\tag{1}$$

which is a sum over each constituent quark *j* of the hadron. The simplest case for the coalescence of comoving equal-mass quarks gives the usual relation [7] $v_{1,H}(N_{cq} p_{T,q}) = N_{cq} v_{1,q}(p_{T,q})$, where N_{cq} is the number of constituent quarks of the hadron.

For each of the five hadron sets in Table 1, the number of quarks of flavor $i(N_i)$ for each side (L: left; R: right) is calculated as the sum of the number of quarks of flavor i in each hadron, multiplied by the weighting factor. With $\Delta N_i = N_i^L - N_i^R$, each set satisfies the following relations:

$$\Delta N_{\bar{u}} + \Delta N_{\bar{d}} = 0, \ \Delta N_s + \Delta N_{\bar{s}} = 0.$$
⁽²⁾

In other words, the two sides have the same number of $\bar{u} + \bar{d}$ quarks and the same number of $s + \bar{s}$ quarks. However, they may have different $N_{\bar{u}}$ and/or N_s . Therefore, they can have a different total net electric charge (*q*), total net strangeness (*S*), or total net electric charge in light quarks (q_{ud}), defined as $\Delta q = q^L - q^R$, $\Delta S = S^L - S^R$, and $\Delta q_{ud} = q^L_{ud} - q^R_{ud}$, respectively. For the hadron sets in Table 1, Equation (2) leads to

$$\Delta q_{ud} = \Delta N_{\bar{d}}, \ \Delta S = 2\Delta N_{\bar{s}}, \ \Delta q = \Delta q_{ud} + \Delta S/3.$$
(3)

Table 1. List of five independent hadron sets, including identical ($\Delta q = 0$ and $\Delta S = 0$) and non-identical ($\Delta q \neq 0$ and/or $\Delta S \neq 0$) constituent quark combinations with similar mass.

Set	Δq	Δq_{ud}	ΔS	ΔB	Left Side (L)	Right Side (R)
1	0	0	0	0	$v_1[K^-(\bar{u}s)] + v_1[\overline{\Lambda}(\bar{u}d\bar{s})]$	$v_1[\bar{p}(\bar{u}\bar{u}\bar{d})] + v_1[\phi(s\bar{s})]$
2	0	0	0	0	$v_1[\overline{\Lambda}(ar{u}ar{d}ar{s})]$	$\frac{v_1}{2}[\overline{\Xi}^+(\overline{d}\overline{s}\overline{s})] + \frac{v_1}{2}[\overline{p}(\overline{u}\overline{u}\overline{d})]$
3	0	0	0	0	$\frac{v_1}{3}[\Omega^-(sss)] + \frac{v_1}{3}[\overline{\Omega}^+(\bar{s}\bar{s}\bar{s}\bar{s})]$	$v_1[\phi(sar s)]$
4	1/3	0	1	-1/3	$\frac{v_1}{2}[\phi(s\bar{s})]$	$\frac{v_1}{3}[\Omega^-(sss)]$
5	2/3	1/3	1	-1/3	$\frac{v_1}{2}[\phi(s\bar{s})] + \frac{v_1}{3}[\bar{p}(\bar{u}\bar{u}\bar{d})]$	$v_1[K^-(\bar{u}s)]$

The total v_1 from each side of a hadron set can be written as $v_1^{L,R} = \sum_i N_i^{L,R} v_{1,i}$. It is important to note that we neglect the effect of different constituent quark masses in this study. Therefore, the above relation applies to light and strange (anti)quarks at the same p_T value or p_T range, and consequently, it also applies to baryons at a p_T value or range that is 3/2 times that of the meson's. The difference in v_1 between the two sides is then given by

$$\Delta v_{1} = v_{1}^{L} - v_{1}^{R} = \sum_{i} \Delta N_{i} v_{1,i}$$

= $(v_{1,\bar{d}} - v_{1,\bar{u}}) \Delta q_{ud} + \left(\frac{v_{1,\bar{s}} - v_{1,s}}{2}\right) \Delta S$ (4)

$$= \left(v_{1,\bar{d}} - v_{1,\bar{u}}\right)\Delta q + \left(\frac{v_{1,\bar{s}} - v_{1,s}}{2} - \frac{v_{1,\bar{d}} - v_{1,\bar{u}}}{3}\right)\Delta S.$$
 (5)

Equations (4) and (5) show the linear dependence of Δv_1 of a hadron set on both Δq and ΔS , where the coefficients are simply determined by the quark-level differences in v_1 . We can also see that the coefficient for the ΔS dependence in Equation (5) is not as clean as the corresponding coefficient in Equation (4).

The difference in v_1 in the hadron sets also depends on ΔB , which represents the difference in the net baryon number between the two sides of a hadron set. Under the condition from Equation (2), we have

$$\Delta B = -2\Delta N_{\bar{s}}/3 = -\Delta S/3. \tag{6}$$

Therefore, Δv_1 depends linearly on both Δq and ΔB as follows:

$$\Delta v_1 = (v_{1,\bar{d}} - v_{1,\bar{u}})\Delta q_{ud} - 3\left(\frac{v_{1,\bar{s}} - v_{1,s}}{2}\right)\Delta B \tag{7}$$

$$= \left(v_{1,\bar{d}} - v_{1,\bar{u}}\right) \Delta q - 3 \left(\frac{v_{1,\bar{s}} - v_{1,s}}{2} - \frac{v_{1,\bar{d}} - v_{1,\bar{u}}}{3}\right) \Delta B.$$
(8)

Assuming that the rapidity of a hadron formed by quark coalescence is the same as that of the coalescing quarks, the v_1 slope at y = 0, denoted as $v'_1 = dv_1/dy(y = 0)$, satisfies exactly the same relations as Equations (4), (5), (7) and (8), where one just needs to replace v_1 with v'_1 [8].

3. Extracting the Δq and ΔS Dependences of the Coefficients

The coalescence sum rule may not be satisfied for certain collision energies and/or systems, e.g., in cases where v_1 is not dominated by parton dynamics. Since Equations (4) and (5) from the coalescence sum rule predict $\Delta v_1 = 0$ for hadron sets with $\Delta S = \Delta q_{ud} = 0$ or $\Delta S = \Delta q = 0$, we propose that the following modified equations are used to fit the Δv_1 data from the five independent sets:

$$\Delta v_1 = c_0 + c_q \,\Delta q_{ud} + c_S \,\Delta S, \tag{9}$$

$$\Delta v_1 = c_0^* + c_q^* \,\Delta q + c_S^* \,\Delta S. \tag{10}$$

This way, a non-zero value of the intercept parameter, denoted as c_0 or c_0^* , would mean breaking the coalescence sum rule. Note that even for a given collision system, the coalescence sum rule may be satisfied around midrapidity but not satisfied at large rapidities; therefore, these coefficients are rapidity-dependent.

To extract the coefficients of the Δq and ΔS dependences of Δv_1 for a given collision system, one can utilize the five-set method by simply fitting the 5 data points using Equation (9) or Equation (10). It should be noted that the fit function represents a two-dimensional plane over the $\Delta q - \Delta S$ space. Therefore, one should not simply fit the Δv_1 data with a one-dimensional function of Δq without taking into account the different ΔS values of different hadron sets.

Alternatively, we can use the three-set method [8], where we take the average of sets 1–3, because they have the same Δq and ΔS values, and denote the average as set A. Let us also denote set 4 and set 5 as set B and set C, respectively. Equation (9) then leads to $(\Delta v_1)_A = c_0$, $(\Delta v_1)_B = c_0 + c_s$, and $(\Delta v_1)_C = c_0 + c_g/3 + c_s$. Thus, we obtain

$$c_0 = (\Delta v_1)_A; \ c_q = 3[(\Delta v_1)_C - (\Delta v_1)_B]; \ c_S = (\Delta v_1)_B - (\Delta v_1)_A.$$
(11)

Similarly, the three-set method yields the coefficients in Equation (10) as

$$c_{0}^{\star} = (\Delta v_{1})_{A} = c_{0}; \quad c_{q}^{\star} = 3[(\Delta v_{1})_{C} - (\Delta v_{1})_{B}] = c_{q}; c_{S}^{\star} = 2(\Delta v_{1})_{B} - (\Delta v_{1})_{A} - (\Delta v_{1})_{C} = c_{S} - c_{q}/3.$$
(12)

We see that the three-set method has an advantage over the five-set method in that we can extract the Δq and ΔS coefficients analytically instead of performing a two-dimensional fit. It should be noted that the CSR predicts the following coefficients [8]:

$$c_0 = c_0^* = 0; \ c_q = c_q^* = v_{1,\bar{d}} - v_{1,\bar{u}};$$
 (13)

$$c_{S} = (v_{1,\bar{s}} - v_{1,s})/2, \ c_{S}^{\star} = (v_{1,\bar{s}} - v_{1,s})/2 - (v_{1,\bar{d}} - v_{1,\bar{u}})/3.$$
(14)

When the difference in v_1 of the hadron sets is expressed in terms of ΔB , we can use the following modified equations, which are similar to Equations (9) and (10):

$$\Delta v_1 = c_0 + c_q \,\Delta q_{ud} + c_B \,\Delta B, \tag{15}$$

$$\Delta v_1 = c_0^* + c_q^* \,\Delta q + c_B^* \,\Delta B. \tag{16}$$

Here, the CSR predicts the following ΔB coefficients:

$$c_B = -3(v_{1,\bar{s}} - v_{1,s})/2 = -3 c_S, \ c_B^{\star} = -3(v_{1,\bar{s}} - v_{1,s})/2 + (v_{1,\bar{d}} - v_{1,\bar{u}}) = -3 c_S^{\star}.$$
 (17)

Obviously, the CSR relates the Δq and ΔS (or ΔB) coefficients to the quark-level differences in v_1 . It should be noted that the quark v_1 in Equations (13), (14) and (17) refers to its value after the partonic evolution just before the quark coalescence. Therefore, these coefficients could be non-zero due to the flavor dependence of the strong interaction or the effect of the electromagnetic field on the produced quarks. Additionally, the extraction methods outlined in this section apply to the v'_1 data in exactly the same way. Furthermore, in the full study [8], we demonstrated these extraction methods for the Δq and ΔS coefficients using results from the AMPT model for Au+Au collisions at different energies.

4. Summary

In this study, we derived the relations for Δv_1 , representing the difference in v_1 between the two sides of a hadron set, using the coalescence sum rule. In earlier studies, seven produced hadron species $(K^-, \phi, \overline{p}, \overline{\Lambda}, \overline{\Xi}^+, \Omega^-, \text{ and } \overline{\Omega}^+)$ were considered, and a non-zero Δv_1 dependence on the difference in the net electric charge (Δq) was considered as breaking the coalescence sum rule and being a consequence of the electromagnetic fields. Our study showed that the coalescence sum rule only leads to a zero Δv_1 for a hadron set if its two sides have identical constituent quark contents (i.e., $\Delta q = \Delta S = 0$). In general, Δv_1 depends linearly on both Δq and ΔS , or on both Δq_{ud} and ΔS , and the same applies to the difference in the v_1 slopes at midrapidity ($\Delta v'_1$). Since there are only five such independent hadron sets, there will be five independent data points from the measurement of a given collision system. We propose using a two-dimensional function such as $c_0 + c_a \Delta q_{ud} + c_s \Delta S$ to extract the coefficients of the Δq_{ud} and ΔS dependences, where a non-zero value of the coefficient c_0 indicates the breaking of the coalescence sum rule. In the five-set method, one fits the five data points using this function. On the other hand, in the three-set method, the three data points with the same Δq and ΔS values are combined into one set, allowing us to obtain the coefficients analytically. The coalescence sum rule relates these coefficients to the quark-level differences in v_1 just before quark coalescence. Therefore, the coefficients can be affected by both the strong interaction and the electromagnetic fields. Equivalently, one can also express Δv_1 or $\Delta v'_1$ in terms of the net baryon difference ΔB , where the function $c_0 + c_q \Delta q_{ud} + c_B \Delta B$ can be used to extract the coefficients, and the coalescence sum rule yields $c_B = -3c_S$.

Author Contributions: Conceptualization, Z.-W.L.; Methodology, Z.-W.L.; Software, K.N.; Formal analysis, K.N.; Resources, S.-S.S.; Writing—original draft, Z.-W.L.; Writing—review and editing, K.N.; Visualization, K.N.; Supervision, Z.-W.L.; Project administration, S.-S.S.; Funding acquisition, S.-S.S. All authors have read and agreed to the published version of the manuscript.

Funding: S.S. was supported in part by the National Key Research and Development Program of China under Grant Nos. 2022YFA1604900 and 2020YFE0202002, and the National Natural Science

Foundation of China under Grant Nos. 12175084 and 11890710 (11890711). K.N. was supported by OSHEC, Department of Higher Education, Government of Odisha, Index No. 23EM/PH/124, under MRIP 2023. Z.-W.L. was supported by the National Science Foundation under Grant Nos. 2012947 and 2310021. The APC was funded by the National Science Foundation under Grant No. 2012947.

Data Availability Statement: Data are contained within the article.

Acknowledgments: We thank Sandeep Chatterjee for the discussions.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Li, B.-A.; Ko, C.M. Probing the softest region of the nuclear equation of state. *Phys. Rev. C* 1998, *58*, R1382. [CrossRef]
- 2. Stöcker, H. Collective Flow signals the Quark Gluon Plasma. Nucl. Phys. 2005, 750, 121–147. [CrossRef]
- 3. Nara, Y.; Niemi, H.; Ohnishi, A.; Stöcker, H. Examination of directed flow as a signature of the softest point of the equation of state in QCD matter. *Phys. Rev. C* 2016, *94*, 034906. [CrossRef]
- 4. Luo, X.; Shi, S.; Xu, N.; Zhang, Y. A Study of the Properties of the QCD Phase Diagram in High-Energy Nuclear Collisions. *Particle* **2020**, *3*, 278. [CrossRef]
- 5. Sorge, H. Elliptical Flow: A Signature for Early Pressure in Ultrarelativistic Nucleus-Nucleus Collisions. *Phys. Rev. Lett.* **1997**, 78, 2309. [CrossRef]
- 6. Das, S.K.; Plumari, S.; Chatterjee, S.; Alam, J.; Scardina, F.; Greco, V. Directed flow of charm quarks as a witness of the initial strong magnetic field in ultra-relativistic heavy ion collisions. *Phys. Lett. B* 2017, *768*, 260–264. [CrossRef]
- Molnar, D.; Voloshin, S.A. Elliptic Flow at Large Transverse Momenta from Quark Coalescence. *Phys. Rev. Lett.* 2003, 91, 092301. [CrossRef]
- 8. Nayak, K.; Shi, S.; Lin, Z.-W. Coalescence sum rule and the electric charge- and strangeness-dependences of directed flow in heavy ion collisions. *Phys. Lett. B* 2024, *849*, 138479. [CrossRef]
- 9. Sheikh, A.I.; Keane, D.; Tribedy, P. Testing the impact of electromagnetic fields on the directed flow of constituent quarks in heavy-ion collisions. *Phys. Rev. C* 2022, 105, 014912. [CrossRef]
- 10. Adamczyk, L. et al. [STAR Collaboration] Beam-Energy Dependence of Directed Flow of Λ , $\bar{\Lambda}$, K^{\pm} , K_s^0 and ϕ in Au+Au Collisions. *Phys. Rev. Lett.* **2018**, 120, 062301. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.