



Conventional and Emerging Extraction Processes of Flavonoids

Mónica L. Chávez-González ^{1,*}^(D), Leonardo Sepúlveda ¹^(D), Deepak Kumar Verma ²^(D), Hugo A. Luna-García ¹, Luis V. Rodríguez-Durán ³^(D), Anna Ilina ¹ and Cristobal N. Aguilar ^{1,*}^(D)

- ¹ Bioprocesses and Bioproducts Research Group, Food Research Department, School of Chemistry, Universidad Autónoma de Coahuila, 25280 Saltillo, Mexico; leonardo_sepulveda@uadec.edu.mx (L.S.); hugoluna@uadec.edu.mx (H.A.L.-G.); annailina@uadec.edu.mx (A.I.)
- ² Agricultural and Food Engineering Department, Indian Institute of Technology Kharagpur, Kharagpur 721302, India; deepak.verma@agfe.iitkgp.ernet.in
- ³ Department of Biochemical Engineering, Unidad Académica Multidisciplinaria Mante, Universidad Autónoma de Tamaulipas, 89840 Ciudad Mante, Mexico; luis.duran@docentes.uat.edu.mx
- * Correspondence: monicachavez@uadec.edu.mx (M.L.C.-G.); cristobal.aguilar@uadec.edu.mx (C.N.A.); Tel.: +52-844-416-1238 (M.L.C.-G.)

Received: 14 January 2020; Accepted: 24 February 2020; Published: 7 April 2020



Abstract: Flavonoids are a group of plant constituents called phenolic compounds and correspond to the nonenergy part of the human diet. Flavonoids are found in vegetables, seeds, fruits, and beverages such as wine and beer. Over 7000 flavonoids have been identified and they have been considered substances with a beneficial action on human health, particularly of multiple positive effects because of their antioxidant and free radical scavenging action. Although several studies indicate that some flavonoids have provident actions, they occur only at high doses, confirming in most investigations the existence of anti-inflammatory effects, antiviral or anti-allergic, and their protective role against cardiovascular disease, cancer, and various pathologies. Flavonoids are generally removed by chemical methods using solvents and traditional processes, which besides being expensive, involve long periods of time and affect the bioactivity of such compounds. Recently, efforts to develop biotechnological strategies to reduce or eliminate the use of toxic solvents have been reported, reducing processing time and maintaining the bioactivity of the compounds. In this paper, we review, analyze, and discuss methodologies for biotechnological recovery/extraction of flavonoids from agro-industrial residues, describing the advances and challenges in the topic.

Keywords: flavonoids; extraction methods; biotransformation; human health

1. Introduction

Flavonoids are natural pigments present in the plant or microbial sources and correspond to a specific group of chemical constituents called phenolic compounds [1,2]. They are found in vegetables, seeds, fruit, and various fruits and alcoholic beverages [3]. Flavonoids have important positive effects on human health especially due to their antioxidant and free radical scavenging. Although several studies have shown that some flavonoids have a pro-oxidant effect, they only occur at high doses, most of which confirm the existence of anti-inflammatory effects, antiviral or anti-allergic, and their protective role against cardiovascular disease, cancer, and various pathologies [2,4].

Flavonoids protect the human body from damage caused by oxidizing agents such as ultraviolet rays, environmental pollution, food chemicals, etc. The human organism cannot produce these chemicals in a protective manner, so they must be obtained by means of feed or as supplements. These compounds have been discovered by Nobel Prize winner Szent-György, who in 1930 isolated a

substance, citrine, which regulates the permeability of the capillaries from the lemon peels. Flavonoids were first identified as vitamin P (because of the ability to increase capillary permeability) and vitamin C2 (because some flavonoids had similar properties to vitamin C) [1]. However, the fact that flavonoids were vitamins could not be confirmed, and both names remain around 1950. Flavonoids contain in their chemical structure a variable number of phenolic hydroxyl groups and excellent properties of iron chelation and other transition metals, which give them a high antioxidant capacity; therefore, they play an essential role in the protection against oxidative damage and have therapeutic effects in a wide range of conditions, including heart disease ischemic, or atherosclerosis cancer [5–7]. Antifree radical properties of flavonoids are primarily aimed at hydroxyl and superoxide radicals, highly reactive species involved in the onset of lipid chain peroxidation and described their ability to modify eicosanoid synthesis (with antiprostanoid and anti-inflammatory reactions) to prevent platelet aggregation (antithrombotic effects) and to protect low-density lipoproteins from oxidation [8,9].

In addition to its known antioxidant effects, flavonoids have other properties, including stimulation of communication through gap junctions, effects on the regulation of cell growth and induction of enzymes, detoxification such as dependent monooxygenase Cytochrome P-450, among others. [10]. However, most of the biological properties of flavonoids are strongly determined by the mode of extraction for their recovery. Efforts have recently been reported to develop biotechnological strategies to reduce or eliminate the use of toxic solvents, reduce processing time, and maintain the bioactivity of the compounds. This paper examines, analyzes, and discusses the biotechnological methodologies and the recovery/extraction of flavonoids from agro-industrial residues, describing the advances and challenges in the field.

2. Flavonoids

Flavonoids are a type of polyphenolic compound, its chemical structure is varied but the general skeleton structure is composed of 15 carbones (C_6 - C_3 - C_6), which are grouped in two aromatic rings (A and B) connected by a 3-carbon bridge that gives rise to an oxygenated heterocycle (C) [11–15] (Figure 1). Flavonoids are derivatives of 1, 3-diphenylpropan-1-one and their biosynthetic pathway is the condensation of three malonyl-CoA molecules with one *p*-coumaroyl-CoA molecule to the intermediate chalcone [16,17]. Flavonoids are water-soluble pigments present in the plant kingdom as secondary plant metabolites [2,18,19], which can be found specifically in the cytosol and stored in the plant cell vacuole [12,17].



Figure 1. Basic chemical structures of flavonoids and their different class.

Flavonoids are classified according to differences in the structure of the heterocyclic C ring; these differences may be caused by the oxidation state and the degree of unsaturation of the heterocyclic ring (or the lack thereof in the case of chalcones). It has been estimated that the number of identified flavonoids exceeds 7000 and that the number of flavonoids continues to increase due to their important biological activities [17]. Variations in the basic structure of flavonoids give rise to six different classes of this group of compounds: Isoflavones, flavanone, flavanone, flavan-3-ol, flavonol, and anthocyanidin, each of which has particular characteristics [18,20].

2.1. Flavanones

Flavanones (dihydroflavones) have a structure that differs in the lack of a double bond (C_2 - C_3) in the C-ring of the flavonoid structure. This type of flavonoid can be found in aromatic plants (such as mint), tomatoes, citrus (especially grapefruit) [12]. Flavanones can be found in nature as forms of aglycones and glycosides, some examples of such compounds are naringenin, hesperetin, and eriodictyol [21].

2.2. Flavonols

Flavonols are called 3-hydroxyflavones and are the most commonly found flavonoids in the plant kingdom [22]. A double bond between C_2 and C_3 and a hydroxyl group is included in their structures [13]. Some of the most important phytochemical compounds that represent this group are as follows: Myricetin, quercetin, isorhamnetin, and kaempferol [12,22,23]. They can be found in a variety of colors (from white to yellow). In nature, flavonols can be found in two forms: Glycosides and aglycone (quercetin and kaempferol) [12].

2.3. Flavones

Flavones can be found in all parts of the plants, above-and belowground, in vegetative and generative organs; stem, leaves, buds, bark, heartwood, thorns, roots, rhizomes, flowers, farina, fruit, seeds, and also in root and leaf exudates or resin. They result from the introduction of a double bond between C_2 and C_3 by the abstraction of two hydrogen atoms [16,18]. Flavones are present in all major land-plant lineages. The plant species that contain flavones belong to over 70 different families in the plant kingdom [16].

2.4. Anthocyanins

Anthocyanins are primarily found in nature in glycosidic form. This type of flavonoid is responsible for plant pigment (such as blue, red, pink, and purple) by the formation of weak covalent bonding complexes with other organic compounds [23,24]. More than 500 anthocyanins have been reported and are the product of methoxylation, hydroxylation, and glycosylation patterns in the B ring. The most representative compounds of this subclass of flavonoids are pelargonidin, cyanidin, and delphinidin [12].

2.5. Flavanols

Flavanols (flavan-3-ols) are also called catechins, which have a typical flavonoid structure but have different hydroxylation patterns of rings A and B and asymmetrical carbon stereochemistry of ring C (C_2 and C_3) [25,26]. The catechins are classified into two groups; free catechins and esterified catechins [27], and constitute the most complex class of flavonoids due to their size, monomers (catechin), or polymeric forms (condensed tannins) [21]. They can be found as the main ingredient in green tea [12].

2.6. Isoflavones

Another type of flavonoids, isoflavones are commonly referred to as phytoestrogens due to their considerable estrogen activity. They are characterized by the fusion of their ring B with the C_3 position of the ring C [21,28–30]. They are an important group in a variety of fields, such as medicine, cosmetics, and nutrition. These flavonoids can be found in plants of the Leguminosae family (soybeans, alfalfa sprouts, and red clover leaves) [30,31].

3. Isolation and Extraction Methods

There is a general methodology consisting of three stages for the isolation, extraction, and identification of phytochemicals from natural sources. Pretreatment or preparation of a sample is the first step in which the centrifuge, filtration, or drying process and others can be used. In the second stage, the extraction, isolation, and purification of flavonoid compounds from different plant samples are most notably. In this step, phytochemicals are extracted using processes such as soxhlet, maceration, water infusion, supercritical fluid extraction, solid microphase extraction, microwave extraction, ultrasound, autohydrolysis, etc. In the last step, the purified and extracted extracts are normally used for further study by chromatography techniques, usually involving the identification, quantification, and recovery of flavonoid compounds.

Details of each method, such as conventional and emerging methods used by a number of researchers for flavonoid extraction, are given in the following sections:

3.1. Conventional Methods

Flavonoid extraction and recovery have been booming over recent years because of population trends in healthier lifestyles and the integration of antioxidants into the diet. Therefore, several methods for extracting flavonoids to increase the extraction yields of these major bioactive compounds have been implemented.

Various extractive methods have been proposed, including maceration, percolation, hydro-distillation, boiling, reflux, soaking, and soxhlet [32]. Soxhlet was the most commonly used method for the extraction of flavonoids due to its simplicity and ease of maintenance, low cost, and lower solvent content compared to other methods such as soaking, boiling, or maceration [14,33,34]. Various solvents such as ethanol, methanol, benzene, chloroform, ethyl acetate, etc. have been tested in this extraction method to compare the effect on extraction yields [15,32].

In general, liquid–liquid or solid–liquid extraction is the most widely used process for the extraction of flavonoids. Although maceration and water infusion are conventional extraction processes, they are still used today [35,36]. These methodologies have adopted the use of solvents such as ethanol, methanol, acetone and not just water for the extraction of bioactive compounds [14,37,38]. These conventional extraction methods are characterized by the use of large amounts of solvent, lower extraction yields, and long extraction times compared to other methods. It has been reported that when extraction methodologies involve heat treatments, degradation in the chemical structures of the extracted flavonoids can result in a reduction in bioactivity [39].

Parameters such as time, particle size, type of solvents, mass to volume ratio, temperature, etc. have been evaluated in conventional extraction methods of flavonoid (Table 1) [40–42]. The nature of the extracting agent (solvent) will affect the type of flavonoid extracted and will directly influence the biological activity of the recovered compounds. Of the solvents tested, ethanol and methanol are the most widely used for the extraction of flavonoids due to higher yields achieved in the recovery of flavonoids [43,44].

Table 1. Summary of studies of isolation and extraction of flavonoids from different plant source	es.
---	-----

Туре	Substrate	Solvent	Temperature (°C)	Time	Analysis	Yields	Bioactivities/ Bioactives	References
ASE	Impatiens glandulifera (roots)	Methanol (80%)	80	30 min	LC-MS	257.34 μg PAC/g of dw	Phenolic acid content	[45]
ASE	Impatiens glandulifera (flowers)	Methanol (80%)	80	30 min	LC-MS	188.86 µg PAC/g of dw	Phenolic acid content	[45]
ASE	Broccoli	Acetone/water/ acetic acid (70:29,5:0.5 v/v/v)	70–80	5 min	-	3377 ± 62 mg GAE/100 g edp	AoA	[46]
ASE	Cabbage common	Acetone/water/ acetic acid (70:29,5:0.5 v/v/v)	70–80	5 min	-	2037 ± 31 mg GAE/100 g edp	AoA	[46]
ASE	Cabbage red	Acetone/water/ acetic acid (70:29,5:0.5 v/v/v)	70–80	5 min	-	2547 ± 18 mg GAE/100 g edp	AoA	[46]
ASE	Cauliflower	Acetone/water/ acetic acid (70:29,5:0.5 v/v/v)	70–80	5 min	-	274 mg GAE/100 g edp	AoA	[46]
ASE	Lepidium sativum	Ethanol (96%)	50	5 min	GC-MS	58 mg RuE/g dm of flavonoid content	AmA and CtA	[47]
ASE	Impatiens glandulifera (leaves)	Methanol (80%)	80	30 min	LC-MS	244.73 μg PAC/g of dw	Phenolic acid content	[45]
HWE	Pine (Pinus rigida × taedaand Pinus koraiensis) bark	Boiling water	100	1 h	-	111–862 mg CAE/g dw	AoA	[48]
Maceration	Broccoli	Acetone/water (70:30 v/v)	4	24 h	LC-MS	82.2 ± 8.9 mg GAE/100 g edp	AoA	[49]
Maceration	Cauliflower	Acetone/water (70:30 v/v)	4	24 h	LC-MS	27.8 ± 71.5 mg GAE/100 g edp	AoA	[49]
Maceration	Chinese cabbage	Acetone/water (70:30 v/v)	4	24 h	LC-MS	118.9 ± 712.5 mg GAE/100 g edp	AoA	[49]
Maceration	White cabbage	Acetone/water (70:30 v/v)	4	24 h	LC-MS	15.37 ± 2.1 mg GAE/100 g edp	AoA	[49]
Maceration	Broccoli	Acetone/water (80:20 v/v)	Room temperature	8 min	-	80.87 ± 1.2 mg GAE/100 g edp	AoA and ApA	[50]
Maceration	Cabbage	Acetone/water (80:20 v/v)	Room temperature	8 min	-	36.77 ± 6.9 mg GAE/100 g edp	AoA and ApA	[50]
Maceration	Broccoli	Methanol/water (80/20 v/v)	Room temperature	-	-	34.571.0 mg GAE/100 g edp	AoA	[51]
Maceration	Solanum scabrum leaves	Acetone	-	72 h	-	34.2 g GAE/100g	AoA	[52]
Maceration	Lepidium sativum	Ethanol (96%)	50	24 h	GC-MS	25 mg RuE/g dw	AmA and CtA	[47]
Maceration	Banana	Water	-	-	-	8.51 μg QuE/g dw	AoA	[42]
Maceration	Pitanga	Ethanol (75%) and Hexane	25	4 h	-	232.2 mg GAE/g and 12.4 mg GAE/g dw, respectively	AoA and AbA	[53]
Maceration	Artocarpus heterophyllus wastes	Ethanol (70%) and ethanol pure	25	72 h	LC-MS	871.4 mg QuE/g dw	AoA	[44]
Maceration	Kinnow mandarin	Methanol (80%)	-	-	LC-MS	28.40 mg GAE/g dw	AoA and AmA	[41]
Maceration	Apple tree wood residues	50% Ethanol	55	2 h	LC-PdAD	43.2 mg GAE/g dw	Food, pharmaceutical and cosmeticapplications	[54]
Maceration	Pinus radiata bark	Acetone:water 70:30 v/v	40	180 min	-	412 ± 0 mg CAE/g	ArA	[55]
Maceration	<i>Quercus (Q. robur</i> <i>L.)</i> bark	Water	25	120 min	-	3.7 ± 0.6 mg GAE/g	AoA and AbA	[56]
Maceration	Chokeberry (Aronia melanocarpa)	50% Ethanol. Ratio 1:20	Ambient temperature	60 min	LC-MS	27.7 mg GAE/g dw	Extraction of bioactive compounds	[35]

Туре	Substrate	Solvent	Temperature (°C)	Time	Analysis	Yields	Bioactivities/ Bioactives	References
Percolation	Artocarpus heterophyllus wastes	Ethanol (70%)	25	1 h	LC-MS	511.6 mg QuE/g dw	AoA	[44]
Reflux	Portulaca oleracea L.	Ethanol–water (70:30, v/v)	-	150 min	-	6.8 mg RuE/g	Flavonoid content	[57]
RSISE	Tomatoes	Ethanol (60%)	Room temperature	15 h	HPLC	602.91 mg GAE/100 g dw (TPC)	AoA	[58]
SDE	Dried leaves of basil (Ocimum basilicum L.), and epazote (Chenopodium ambrosioides L.).	Water	-	30 min	GC-MS	0.47 y 0.39% yield of EO from basil and epazote	Essential oil extraction	[59]
SIE	Vitis vinifera waste	Methanol or with ethanol	25	19 h	HPLC	67.88 mg GAE/g dw	ArA	[60]
Soxhlet	Portulaca oleracea L.	Ethanol–water (70:30, v/v)	-	300 min	-	7.0 mg RuE/g	Flavonoid content	[57]
Soxhlet	Buddleia officinalis Maxim	Ethanol (95%)	-	2 h	-	62.56 mg CAE/g dw	AoA	[61]
Soxhlet	Morus nigra (dried)	Petroleum ether	50	3 h	-	58.94% of flavonoid yield	AoA	[62]
Soxhlet	Fresh leaves of Vernoniaamygdalina	Water	100	8 h	GC-MS	-	AoA	[63]
Soxhlet	Vernonia cinerea leaves	Ethanol (60%)	-	2 h	LC-Q-TOF	-MS dw	AoA	[32]
Soxhlet	Artocarpus heterophyllus wastes	Ethanol (70% and pure)	Boiling point	5 h	LC-MS	381.4 mg QuE/g dw	AoA	[44]
Soxhlet	Impatiens glandulifera (leaves)	Chloroform, 80% Methanol and pure Methanol	-	72 h	LC-ESI-MS	286.39 mg PAC/g dw	AoA	[45]
Soxhlet	Impatiens glandulifera (roots)	Chloroform, 80% Methanol and pure Methanol	-	72 h	LC-MS	281.82 mg PAC/g dw	AoA	[45]
Soxhlet	Impatiens glandulifera (flowers)	Chloroform, 80% Methanol and pure Methanol	-	72 h	LC-MS	188.07 mg PAC/g dw	AoA	[45]
Soxhlet	Pinus radiata bark	Acetone:water 70:30 v/v	82	60, 120, 180, and 360 min	-	622 ± 40 mg CAE/g	ArA	[55]
Soxhlet	Spearmint (<i>Mentha spicata</i> L.)	Methanol	40	6 h	HPLC	0.144 mg CAE/g dw	Flavonoid extraction	[64]
Soxhlet	Knotwood (Populus tremula)	Methanol	-	48 h	LC-MS	11.5 mg/g	Flavonoid extraction	[65]

Table 1. Cont.

Abbreviations: AbA: Antibacterial activity; AmA: Antimicrobial activity; AoA: Antioxidant activity; ApA: Antiproliferative activities; ArA: Antiradical activity; ASE: Accelerated solvent extraction; CAE: Catechin equivalents; CtA: Cytotoxicity activity; dm: Dry matter; dw: Dry weight; edp: Edible portion; ESI: Electrospray ion source; GAE: Gallic acid equivalents; GC-MS: Gas chromatography mass spectrometer; HPLC: High performance liquid chromatography; HWE: Hot water extraction; LC: Liquid chromatography; LC-MS: Liquid chromatography mass spectrometer; MS: Mass spectrometer; PAC: Phenolic acid content; PdAD: Photodiode array detector; Q-TOF: Quadrupole-time of flight; QuE: Quercetin equivalents; RSISE: Rotary solid–liquid solvent extraction as Traditional Method; RuE: Rutin equivalents; SDE: Steam distillation extraction; SIE: Solid–liquid extraction; TPC: Total phenolic content.

The high demand for antioxidants gave way to the search for methodological alternatives that would increase the yield of flavonoid extraction and reduce process costs. In addition, it has been found that the methodologies implemented are cleaner and environmentally friendly.

3.2. Emerging and Advanced Methods

Two of the most widely used techniques for the extraction of flavonoids are emerging microwave (MAE) and ultrasound-assisted extraction (UAE) technologies. Table 2 shows some of the published works on the extraction of flavonoids using these emerging methods.

 Table 2. Studies on emerging methods of isolation and extraction of flavonoids from different plant sources.

Substrate	Extra	ction Conditio	ons	Analysis	Bioactive	Bioactivity	References
Jubbune	Solvent	Temperature (°C)	Time	, , , , , , , , , , , , , , , , , , ,	Dioneti i e	,	
			(A) M	icrowave-Assisted Ext	raction		
Fresh leaves of Vernonia amygdalina	Water	100	7 min	GC-MS	87.05 mg QuE/g total flavonoid content	AoA	[63]
Black rice (<i>Oryza</i> sativa cv. <i>Poireton</i>) husk	Ethanol 40%–70% (relation m/v (1:20, 1:35, 1:50)	40-60	20–60 s	HPLC	Gallic acid, <i>p</i> -coumaric acid, ferulic acid, quercetin, salicylic acid, quimic acid, apigenin, syringic acid, chlorogenic acid, catechin	AoA	[66]
Apple tree wood residues	Ethanol (60%)	100	20 min	HPLC-PdAD	47.7 mg GAE/g dw	Pharmaceutical and cosmetic applications	[54]
Moringa oleifera leaves	Water, ethanol:water, and etanol	50–180	3–20 min	HPLC-ESI-Q- TOF-MS	Quercetin sambubioside/ Quercetin-3-vicianoside, kaempferol diglycoside, multiflorin B, kaempferol-3-O-glucosidde, quercetin-3-O-glucosidde, quercetin hydroxyl methylglucoside, quercetin hydroxyl methylglutaroyl, glycoside, quercetin triacetylgalactoside, quercetin acetyl glycoside, isorhamnetin-3-O-glucosid quercetin, kaempferol	e, AoA le,	[67]
Coriander (Coriandrum sativum L.) seeds	Ethanol (52%)		35 min	-	382.32 mg GAE/100 g dw	AoA	[68]
<i>Quercus</i> (Q. robur L) bark	Water	100	120 min	HPLC-PdAD-ESI- MS/MS	16.50 ± 0.07 mg GAE/g dw	AoA	[69]
Canola seed cake	Ethanol (10%)	70	20 min	-	-	Polyphenols extraction	[70]
Tomatoes	Ethanol	Room temperature	10 min	HPLC	646.40 mg GAE/100 g dw (TPC)	AoA	[58]
Pinus radiata bark	acetone:water 70:30 v/v	-	3 min	-	479 ± 49 mg CAE/g dw	ArA	[55]
Grape skins	40% methanol	100	5 min	HPLC	1.858 mL AntE/g of MEC/MoS	Anthocyanins extraction	[71]
Uncaria sinensis	Ultra pure water	100	20 min	LC-MS	44 mg EpiCAE/100 g	Quality of medicinal herbs	[72]
Buddleia officinalis Maxim	Ethanol (65%–100%)	40-78	10–30 min	-	75.33 mg CAE/g dw	AoA	[61]

Substrate	Extrac	ction Conditio	ons	Analysis	Bioactive	Bioactivity	References
Jubbunc	Solvent	Temperature (°C)	Time	, in the second s	21040110	,	
			(A) <i>Mi</i>	icrowave-Assisted Ex	traction		
Citrus unshiu	Ethanol (70%)	140	8 min	HPLC-PdAD	47.7 mg HspE/g of MEC/MoS	ТрР	[73]
Defatted residue of yellow horn	Etanol (40%)	50	7 min, 3 extraction cycles	-	11.62% (triterpene saponins) MEAC × 100/MoS	Food and pharmaceutical industries	[74]
Peanut skins	Ethanol (30%)	-	30 s	HPLC and LC-MS-MS	144 mg PAC/g of MEC/MoS	AoA	[75]
Pigeonpea (Cajanus cajan) leaves	Ethanol (80%)	65	1 min (2 min total), 2 extraction cycles	RP-HPLC-PdAD	18.8 mg AAE/g and 3.5 mg PinE/g of MEC/MoS	TpP	[76]
Portulaca oleracea L.	Ethanol–water (70:30, v/v)	50	9 min	_	7.1 mg RuE/g	Flavonoid content	[57]
Purple corn	15 M HCl: 95% ethanol in 15:85 ratio	55	19 min	LC-MS	1.851 mg AntE/g of MEC/MoS	Anthocyanin extraction	[77]
Radix puerariae	Ethanol (70%)	-	6 min	-	8.37 mg RuE/g	Flavonoid content	[78]
Sea buckthorn (Hippophae rhamnoides) food by-products	Water	20–100	15 min	HPLC	Flavonol isorhamnetin 30-rutinoside	AoA	[79]
Tea residues (oolong)	Water	230	2 min	GC-MS	144.0 mg GAE/g dw	AoA	[80]
Tea residues (green)	Water	230	2 min	GC-MS	87.2 mg GAE/g dw	AoA	[80]
Vitis vinifera seed	Methanol	110	60 min	HPLC	86.2 mg GAE/g and 46.8 mg CAE/g dw	ArA	[60]
Alpinia zerumbet (Pers.) Burtt et Smith leaves	Ethanol (70%)	60–70	3s	HPLC	11% w/w	Flavonoid extraction	[81]
Bark of Phyllanthus emblica L.	Aqueous ethanol (75%)	45	25 min	-	19.78 %	AoA	[82]
Citrus mandarin peels	Methanol (66%)	1–120	49 s	HPLC	3779.37 μg PAC/g of MEC/MoS	Phenolic acids extraction	[83]
Milk thistle seed	Ethanol (82%)	112	60 min	-	56.67 mg SilE/g of MEC/MoS	Silymarin extraction	[84]
<i>Morus alba</i> L. leaves	Ethanol (60%)	100	5 min	-	2.4% flavonoid	AfA	[85]
Onion (<i>Allium cepa</i> L.)	Methanol (80%)	2–100	Up to 60 min	HPLC	330.46 mg of flavonol content	Food application	[86]
Herba epimedii	Ethanol	-	-	HPLC	921 peak Area of total flavonoids	Flavonoid extraction	[87]
Myrica rubra leaves	-	60	20 min	HPLC	-	Polyphenol extraction	[88]
Radix astragali	Ethanol (90%)	110	50 min	HPLC	1.190 mg flavonoids/g	Flavonoid extraction	[89]
Radix astragali roots	Ethanol (90%)	110	25 min (50 min total), 2 extraction cycles	-	1.19 mg/g (flavonoids) of MEC/MoS	Flavonoid extraction	[89]

Table 2. Cont.

Substrate	Extrac	tion Condition	ons	Analysis	Bioactive	Bioactivity	References
Substrate	Solvent	Temperature (°C)	Time	j	Dieucuire	,	
			(A) Mid	crowave-Assisted Ex	xtraction		
Platycladus orientalis leaves	Methanol (80%)	-	5 min	-	1.72% (flavonoids) MEAC × 100/MoS	Flavonoid extraction	[90]
Epimedium koreamum Nakai	Ethanol (40%)	-	15 min	LC-ESI-MS	280 m AU/min	Flavonoid extraction	[91]
Olive leaves	Ethanol (80%)	-	8 min	HPLC-PdAD	95% (oleuropein) MEAC × 100/MTACcS	Biophenols extraction	[92]
Acanthopanax senticosusleave	Ethanol s (50%)		10 min	ESI-MS	-	Flavonoid extraction	[93]
Eucommia ulmodies oliv.	Methanol:water acid (20:80:1.0, v/v)	r:acetic -	30–40 s	HPLC	75.6%–83.2% (geniposidic acid) and 77.4%–86.3% (chlorogenic acid) of MEAC × 100/MTACcS	TpP	[94]
Capsicum fruit	Acetone	-	15 min	GC-MS	0.48 mg CpE/g fw	Food aditives	[95]
			(B) Ultr	rasound-Assisted Ex	xtraction		
Dysphania ambrosioides (L)	Ethanol (57% w/w)	57	60 min	-	1.09% of flavonoids equivalents of rutin	Flavonoid extraction	[96]
Lepidium sativum seeds	Ethanol	50	24 h	GC-MS	97 mg GAE/g dm	AmA and CtA	[47]
Fruit of rugose rose (<i>Rosa rugose</i> Thumb)	Ethanol (50%)	50	40 min	-	31.88 mg /g dw	AoA	[97]
Impatiens glandulifera (flowers)	Methanol (80 %)	30	60 min	LC-MS	216.03 μg/g dw	AoA	[45]
Impatiens glandulifera (leaves)	Methanol (80 %)	30	60 min	LC-MS	291.55 μg/g dw	AoA	[45]
Impatiens glandulifera (roots)	Methanol (80 %)	30	60 min	LC-MS	286.04 μg/g dw	AoA	[45]
Kinnow mandarin	Ethanol (80%)	35, 45, 55	40–70 min	HPLC	28.40 mg GAE/g extract	AoA and AmA	[41]
Nephelium lappaceum <i>L. fruit peel</i>	Solid–liquid ratio 1:18.6 g/mL	50	20 min	-	10.26 ± 0.69 mg AntE/100 g; 552.64 ± 1.57 mg GAE/100 g; 104 ± 1.13 mg RuE/100 g	Flavonoid extraction	[98]
<i>Curry leaf</i> (Murraya koenigii L.)	Methanol 80% 55.9% 145.49 W	55.9	-	UHP-LC	0.482 mg CAE/g dw; 0.517 mg NrgE/g dw; 0.394 mg QuE/g dw	Pharmaceutical application	[99]
Portulaca oleracea L	Ethanol 39.01%	55.25	15 min	-	16.25 mg RuE/g dw	-	[100]
<i>Pinus radiata</i> bark	acetone:water 70:30 v/v	-	3-12 min	-	388 ± 7 mg CAE/g bark	ArA	[55]
Morus laevigata W. M. alba L. and M. nigra L	Methanol (80%)	-	-	HPLC	3.89 to 11.79 μmoL GAE/100 g	AoA	[101]
Portulaca oleracea L.	Ethanol–water (70:30, <i>v</i> / <i>v</i>)	25	60 min	-	6.7 mg RuE/g	Flavonoid content	[57]
<i>Vitis vinifera</i> seed	Methanol	25	60 min	HPLC	55.9 mg GAE/g and 39.5 mg CAE/g dw	ArA	[60]

Table 2. Cont.

Substrate	Extraction Conditions			Analysis	Bioactive	Bioactivity	References
	Solvent	Temperature (°C)	Time	ý		2	
			(C) 5	Supercritical Fluid Ext	raction		
Dried bilberry fruits (V. <i>myrtillus</i> L.)	CO ₂ + Ethanol (10%)	-	30 min	HPLC-PdAD-ESI-	MS/ M52 mg AntE/g dw	AoA	[102]
Spearmint (Mentha spicata L.)	Absolute ethanol (EtOH) Flow rate: 3 g/min Pressure: 200 bar	60	60 min	HPLC	0.140 mg CAE/g dw	Bioactive flavonoid extraction	[64]
Pueraria lobata	Ethanol Flow rate: 3 g/min Pressure: 20.04 MPa	50.24	90 min	-	16.95 ± 0.43 mg flavonoid/g dw	Flavonoid extraction	[103]
Ganoderma atrum	CO ₂ + ethanolFlow rate: 30 L/h (80 g sample) Pressure: 25 MPa	55	3 h	-	1.52% (triterpenoid saponins) MEAC × 100/MoS	Triterpenoid saponins extraction	[104]
Lepidium sativum	CO ₂ + ethanol	50	70 min	GC-MS MALDI-TOF-MS	58 mg RuE/g dm (Total flavonoid)	AmA and CtA	[47]

Table 2. Cont.

Abbreviations: AAE: Ajaninstilbene acid equivalents; AfA: Antifatigue activity; AmA: Antimicrobial activity; AntE: Anthocyanins equivalents; AoA: Antioxidant activity; ArA: Antifatigue activity; CAE: Catechin equivalents; CpE: Capsaicin equivalents; CtA: Cytotoxicity activity; dm: Dry matter; dw: Dry weight; EpiCAE: Epicatechin equivalents; ESI: Electrospray ion source; GAE: Gallic acid equivalents; GC-MS: Gas chromatography mass spectrometer; HPLC: High performance liquid chromatography; HspE: Hesperidin equivalents; LC: Liquid chromatography provide the spectrometer; MAE: Microwave assisted extraction; MALDI: Matrix-assisted laser desorption ionization; MEAC: Mass of extracted active compound; MEC: Mass of extracted compound; MoS: Mass of sample; MS: Mass spectrometer; MTACCS: Mass of total active compound content in the sample; NrgE: Narengine equivalents; PAC: Phenolic acid content; PdAD: Photodiode array detector; PinE: Pinostrobin equivalents; SilE: Silymarin equivalents; TOF: Time of flight; TPC: Total phenolic content; TpP: Therapeutic potential; UHP: Ultra high performance.

3.2.1. Microwave-Assisted Extraction

(96 %)

Microwave is an electromagnetic spectrum of radiation ranging from 300 MHz (radio radiation) to 300 GHz (infrared radiation). This heating technique uses microwave energy and is based on the direct effect of microwaves on dipole polarization and ion conduction molecules [105–107] (Figure 2). The extraction of flavonoids may be affected by a large number of parameters, among the most important of which are: Time, temperature, plant material-solvent ratio, solvent concentration, solvent polarity, irradiation, frequency of intensity, and microwave power [63,106,108–110].

It has been reported that MAE allows for a significant reduction in the extraction times of a wide variety of compounds, also reduces the volumes of solvents used, and it has been shown that the extraction yields of bioactive compounds are superior to conventional methods such as maceration, Soxhlet, or heat reflux [66,106,112,113]. Reduction of extraction times and the use of solvents are employed to improve the cost of extraction [39].



Figure 2. Schematic setup of microwave-assisted extraction (MAE) [111].

The choice of solvent is not only important in this methodology; the dielectric properties of the solvent must be taken into account [106]. The most commonly used solvents for MAE are ethanol and methanol, both of which have shown the best extraction yields, although water has also shown positive effects [66,68,107]. The efficacy of MAE will also depend on the type of flavonoid to be recovered. For example, the polarity of the flavonoid will be a very important parameter to consider; the type of solvent used will be the polarity of the desired recovery. Moreover, the solvent used in the extraction process may have an effect on the bioactivity of the recovered flavonoids [66]. Apolar solvents such as dichloromethane, ethyl acetate, diethyl ether, chloroform are commonly used for the extraction of isoflavones, flavones, and methylated flavones due to their apolar nature. In contrast, solvents such as ethanol or methanol are used to extract polar flavonoids such as flavonoid glycosides and aglycones (Table 2A).

3.2.2. Ultrasound-Assisted Extraction

Ultrasound-assisted extraction is a technique that is used to rupture the plant material and extract the bioactive compounds with applications in industries such as food and pharmaceuticals [108]. This technique is based on the phenomenon of acoustic cavitation, which consists of the formation of bubbles and the subsequent rupture, which causes the release of bioactive compounds, and this rupture depends on the extraction conditions [97,114] (Figure 3).

The cavitation effect produced by this methodology not only enables the destruction of the cell walls of the plant material but also promotes the reduction of the particle size that benefits the solvent–substrate interaction [97]. There are many variables that can have an effect on flavonoid extractive processes and therefore on the number of experiments; in order to optimize a particular process, experimental matrices are usually used to perform the optimization process in order to determine the conditions that favor the recovery of the maximum flavonoid content. Most of the published works have opted to use the surface response methodology to achieve this objective [63].



Figure 3. Schematic setup of ultrasound-assisted extraction (UAE) [111].

The UAE increased the yield of bioactive compounds and the yield of flavonoids were shown to be variable depending on the method of extraction, as well as on the type of plant material (Table 2B). Mainly phenolic compounds and flavonoids, as well as the reduction of extraction times, and another of the advantages of this methodology is that it improves the biological properties of the extracts [115]. The extraction of bioactive compounds depends on a variety of factors, such as the frequency used and usually between 20 kHz and 100 MHz [116], solvent selection, solvent concentration, solid–solvent ratio, temperature, and time extraction (Table 2B).

Among the flavonoid compounds that have been extracted and recovered using this methodology are the following: Rutin, narcissin, nicotiflorin, epicatechin, epicatechin gallate, catechin, procyanidin B₂, apiofuranosyl(1^{'''} \rightarrow 2^{''})- β -D-glucopyranosyl] rhamnocitrin, quercetin-3 -*O*-rhamnoglucoside, quercetin-3-*O*- β -D-glucopyranoside, myricetin-5[']-*O*- β -D-glucopyranoside, 4[']-*O*-(3^{'''}-*O*-dihydrophaseoyl- β -D-glucopyranosyl) rhamnocitrin, formononetin-7-*O*-glucoside, myricomplanoside, kaempferol-3-*O*-glucosylrutinoside, complantoside A, quercetin-3-*O*-acylglycoside, etc. [117–119].

3.2.3. Supercritical Fluid Extraction

Any substance at a temperature and pressure above its thermodynamic critical point is a supercritical fluid. Under these conditions, the properties of the fluids generate a high diffusivity and low viscosity of the solvents used to improve the process of transfer of the matter [120]. Due to this, the SFE methodology (Figure 4) has reported flavonoid extraction yields much higher than those used in other techniques [121].

The most commonly used solvent in this extraction method is carbon dioxide (CO_2) due to its numerous advantages, such as that it is flammable, nontoxic, cheap, and very easy to remove due to its volatility [123,124]. It is a strong solvent for supercritical extraction with all these features.

Certain advantages of this extraction methodology are low temperatures that maintain the integrity of the products, high volatility of the solvents which keep the waste low, the extraction is carried out without phase changes, easy separation of volatile and nonvolatile compounds. However, they present some limitations such as the difficult equilibrium between solute and solvent, may require other separation processes, high pressures hinder the continuous addition of solids to the extract, operating costs are high, equipment is low, maintenance cost is high, etc. [125,126].





Figure 4. Schematic setup of supercritical fluid extraction (SFE) [122].

Temperature is one of the most important factors in the process. In this methodology, the use of reduced temperatures is intended to keep the product as stable as possible. In one study, for example, the valorization of agro-industrial residues (expellers) from the extraction of soybean oil by pressing was analyzed. Extractions were carried out at 40 MPa and at 35 or 40 °C using CO₂ as a solvent. Moreover, the expellers were impregnated with ethanol. The best results showed the flavonoids content of 65.0 and 31.3 QE/100 gdm [127]. On the other hand, flavonoids were extracted from Odontonema strictum leaves with supercritical carbon dioxide and ethanol. The effect of temperature (55–65 °C) on the total flavonoid content was optimized. The total flavonoid content and flavonoid recovery vary respectively from 99.33 to 247.78 mg/g of dried extract and 10.68–18.92 mg/g of dried leaves powder [128]. In another study, supercritical CO₂ extraction with/without ethanol from *Citrus unshiu* peels was examined at a temperature of 59.85 °C and a pressure of 30 MPa. The best results obtained were that the extracts, including nobiletin, increased the concentration of ethanol in supercritical CO₂ and increased the time of extraction. On the other hand, the role of pressure in these methods is very important and depends on the solvent used. For example, an effective method of extracting Medicago sativa using enzyme-assisted supercritical fluid was developed in another study. The design of the Box-Behnken was chosen to optimize the extraction process parameters, including pressure (100–300 bar). Optimal extraction parameters for total polyphenol content were: 68 °C, 205 bar, and 15.5% for temperature, pressure and co-solvent content, respectively. This methodology provides effective enzyme-assisted supercritical fluid extraction for the enhanced release of polyphenol compounds [129]. Supercritical fluids are a more efficient, safer, and environmentally friendly method for extracting and recovering flavonoids for the purpose of bioactive compounds study (Table 2C).

3.2.4. Enzyme and Microorganism-Assisted Extraction

There are several different techniques for the extraction of flavonoids, including conventional solid–liquid extraction, pressurized fluid extraction, pressurized hot water extraction. Supercritical fluid extraction, MAE, UAE, and pulsed electrical field extraction are among the most sustainable alternatives to these traditional methods [130]. The process of extraction of flavonoids is usually performed using organic solvents mixed with or without water. The extraction can be controlled by different parameters, such as the selection of extraction procedures, as well as the flow rate used for

extraction, solvent and temperature, pressure and time. In addition, other techniques have yielded similar results; these techniques involve the use of microorganism-and enzyme-assisted extractions.

Enzyme-assisted extraction is a promising alternative to conventional extraction methods where a high amount of solvent is used. The advantages of this method are regioselectivity and specificity of enzyme action, their ability to catalyze reactions in aqueous solutions under mild conditions [131]. Furthermore, the application of enzyme is an environmentally friendly method for the extraction of flavonoids due to a decrease in solvent quantity in order to reduce extraction times and increase extraction yield and quality of flavonoids. Enzymes and microorganisms may also be used for pretreatment in different agro-industrial and food industries.

Enzymes obtained from fungi, bacteria, vegetable extracts and animal organs, such as pectinases, different glucanases, hemicellulases, cellulases, etc., in mixtures or alone, break down the cell wall by hydrolysis of biopolymeric components to increase the permeability of the cell wall and also increase the yield of flavonoids and other physiologically active extractants [132,133].

Hydrolytic enzymes can break down the cell wall of polysaccharides to improve intracellular release. Cellulose is hydrolyzed by four classes of enzymes: Endo-and exoglucanases, cellobiohydrolases, and β -glucosidases. The main chain of hemicellulose can be hydrolyzed by endooxylanases, endomannanases, β -xylosidases, and β -mannanases. Degradation of the backbone of pectin requires a number of enzymes such as pectin lyases, pectate lyases, endoand exo-polygalacturonases, endo- and exo-rhamnogalacturonases and rhamnogalacturan lyases. In addition, to hydrolyze the side chains of polysaccharides, by-product enzymes (such as arabinases, galactosidases and feruloyl, esterases) are needed [134].

Several authors have utilized enzymatic treatments with commercial preparations for the extraction of flavonoids from plant material (Table 3). These commercial enzymes usually have one or more main hydrolytic activities and a number of side activities. Commercial pectinases have been used for enzyme-assisted extraction of black currant phenols (*Ribes nigrum*) juice press residues [135]. Mixtures of pectinases and cellulases have been used for the extraction of polyphenols from grape pomace [136,137]. A combination of enzymatic hydrolysis and ultrasonic-assisted extraction was used for the extraction of flavonoids from pigeon pea (*Cajanus cajan*) leaves [138], celery (*Apium graveolens*) leaves [139], shepherd's purse (*Capsella bursa-pastoris*) pulp [140], and mulberry (*Morus nigra*) must [141]. A set of combinations of different extraction methods (*viz.* enzymatic hydrolysis, ultrasonic, and microwave-assisted extraction) that were used for the recovery of flavonoids from *Nitraria tangutorun* juice by-product [142].

Special attention must be paid to the presence of undesirable enzyme activities during the enzyme-assisted extraction of flavonoids. These activities may be caused by endogenous enzymes of plant material or by side activities of commercial preparations. Polyphenol oxidases may induce coupled oxidative browning reactions; β -glucosidases, β -galactosidases, and α -L-arabinosidases may hydrolyze native glycosylated anthocyanins and cause unstable aglycons [135]. For example, Kammerer et al. (2005) [136] observed a low yield of anthocyanin recovery (2.9%) during enzyme-assisted extraction of polyphenols from grape pomace (Vitis vinifera) at the pilot plant level. This low yield was associated with the action of endogenous enzymes. In subsequent research, thermal inactivation of endogenous enzymes to be recovered from grape pomace [137].

Landbo and Meyer (2001) **[135]** found that two commercial pectinases had a negative effect on the recovery of anthocyanins during enzyme-assisted extraction of phenolic compounds from black currant (*Ribes nigrum*) pomace. This negative effect was associated with the presence of glycosidase activities in enzyme preparations. On the other hand, Xu et al. (2013b) **[145]** used the activity of glycosylase present in commercial enzyme preparations to improve the extraction of bioactive compounds from two medicinal plants (*Glycyrrhizae radix* and *Scutellariae radix*). Bifunctional enzymes were used for the simultaneous degradation of the cell wall and deglycosylation of native flavonoid glycosides.

Source	Enzyme (s)	Compound (s)	Reference
Ginkgo biloba leaves	Cellulase	Flavonols	[143]
Grape (<i>Vitis vinifera</i>) skins	Oenological preparation (pectinase + cellulase + hemicellulase)	Anthocyanins, flavonol glycosides, and flavan-3-ols	[144]
Mulberry (<i>Morus nigra</i>) must	Pectinase	Anthocyanins and nonanthocyanin flavonoids	[141]
<i>Nitraria tangutorum</i> juice by-products	Cellulase	Anthocyanins and nonanthocyanin flavonoids	[142]
Shepherd's purse (Capsella bursa-pastoris) pulp	Pectinase and cellulase	Flavonoids	[140]
Glycyrrhizae radix	Cellulase	Liquiritigenin and isoliquiritigenin	[145]
Celery (Apium graveolens) leaves	Pectinase	Luteolin and apigenin	[139]
Scutellariae radix	Naringinase	Bacalein and wogonin	[146]
Grape (Vitis vinifera) pomace	Pectinase and cellulase	Anthocyanins and nonanthocyanin flavonoids	[137]
Pigeonpea (<i>Cajanus cajan</i>) leaves	Pectinase	Luteolin and apigenin	[138]
Grape (Vitis vinifera) pomace	Pectinase and cellulase	Anthocyanins and nonanthocyanin flavonoids	[136]
Black currant (<i>Ribes nigrum</i>) pomace	Pectinase	Anthocyanins	[135]

Table 3. Studies on enzyme assisted isolation and extraction of flavonoids from different plant sources.

Chen et al. (2011) [146] developed a method for enzyme-assisted extraction of flavonoids from Ginkgo biloba leaves in which the enzyme was used not only for cell wall degradation but also to increase the solubility of target compounds. They used commercial *Penicillium decumbens* cellulase with high transglycosylation activity. The presence of maltose *P. decumbens* cellulase transglycosylated flavonol aglycones in more polar glucosides with higher solubility in polar solvents improves the extraction yield.

The parameters to be taken into account for increasing the extraction yield of flavonoids are treatment time, pH, and temperature, as well as the enzyme quantity in relation to the concentration of the substrate. Various studies reported the optimization of these parameters [143,144].

There are also disadvantages of enzymatic methods, which have been reported in many reports: To date, the enzyme tested has not been able to achieve complete plant cell wall hydrolysis due to the major difficulties of scale-up enzyme-assisted extraction at industrial level and the relatively high cost of biocatalyst for large volumes of raw materials [132]. The use of microorganisms as enzyme producers may replace the use of food-grade enzymes in the extraction of flavonoids. However, their growth and activity are sensitive to changes in environmental conditions (temperature, percentage of dissolved oxygen, agitation rate, design of reactors, and availability of nutrients) [147].

4. Transformation of Flavonoids

4.1. Microbial Biotransformation

Biotransformation of flavonoids to increase the biological activity of the recovered compounds is a trend. Biotransformation is a process in which the chemical structure of the compounds is modified by the use of microorganisms. The objective of biotransformation is to produce fine chemical compounds (high added value) that are difficult to produce by chemical synthesis under low-severity reaction conditions. These processes have good production yields and the recovery of flavonoids. The main biotransformations reported in flavonoids are as follows: Dehydroxylation, dehydrogenation, hydrogenation, glycosylation, O-methylation, O-demethylation, deglycosylation, cyclization, sulfation, and carbonyl reduction [148]. This section discusses some biotransformation studies on flavonoids using filamentous fungi and bacteria.

Filamentous fungi and 20 strains of *Streptomyces* for the production of flavonoids were evaluated and two derivatives of quercetin were obtained from *Beauveria bassiana* ATCC7159 in the bioconversion of quercetin. Furthermore, the bioconversion of rutin was obtained by *Cunninghamella echinulata* ATCC 9244 rutin sulfate, rutin glucuronide, and rutin methylation. This biotechnology method was appropriate to produce biologically-active flavonoids [149]. In another study, the biodegradation of isoflavones into 4'-fluoroisoflavone were evaluated by *Aspergillus niger* and *Cunninghamella elegans* strains and obtained more than 20 metabolites. *A. niger* was the microbial strain that has the most ability to degrade isoflavones which could be used as bioactive compounds [150]. Furthermore, the biotransformation of icariin, epimedin C, epimedoside A, epimedin A, and epimedin B were evaluated from the *Epimedium koreanum* plant using *Cunninghamella blakesleana*. This process generated flavonoids that posed potential applications in the pharmaceutical and food industries [151].

A new method was developed to produce genistein from roots of pigeon pea (*Cajanus cajan*) using immobilized strains of *Aspergillus oryzae* and *Monacus anka*. This biotransformation method was a good alternative to the production of genistein from plants with food industry potential [152]. Different filamentous fungi used in solid-state fermentation were evaluated for the biotransformation of phenolic compounds in cauliflower leaves. *A. sojae* strain was best suited to high yields of flavonoids, including kaempferol-derived metabolites. This bioprocess was proposed as an alternative for the development of the concept of bio-refinery and the use of agricultural by-products [153]. In another study, a process of biotransformation of phenolic compounds from citrus waste using solid-state fermentation by *Peacilomyces variotii* was developed, where remarkable production of naringenin and hesperetin was achieved and the antioxidant capacity increased to 73%. These compounds are of high added value that can be used in the food sector [154].

Moreover, a strategy for the biodegradation pathway from tyrosine to the production of fisetin using *Escherichia coli* has been developed. The production of this flavonoid is of great interest because it has different biological properties for human health, such as antiviral and anticancer [155]. In another study, the biotransformation of soy isoflavones in *ortho*-hydroxyisoflavones was evaluated using CYP105D7 from *Streptomyces avermitilis* MA4680 and expressed in *Pichia pastoris*. This study provides evidence of the great potential of the use of genetic microorganisms for the production of isoflavonoids for food industry applications [156]. Puerarin catalyzed by Bacillus cereus NT02 was evaluated for its biotransformation. The results showed that puerarin phosphorylation innovations in medicinal chemistry have particular importance [157].

4.2. Enzyme-Catalyzed Transformation

Flavonoids are characterized by their low solubility and stability in aqueous and lipid phases. For instance, aglycones are less soluble than their derivatives. Aglycones are readily absorbed by passive diffusion through biological membranes, while flavonoid glucosides can be introduced into cells by means of a sodium-dependent glucose transporter 1. Therefore, deglycosylation is important for the assimilation of dietary flavonoids [158]. They may be modified chemically, enzymatically, or chemo-enzymatically to enhance these properties.

Glycosylation and acylation are the most important transformations of flavonoids catalyzed by enzymes. Glycosylation allowed flavonoids to enhance their hydrophilic character by adding sugars, while acylation makes them more hydrophobic due to the combination of fatty acids. Chemical acylation is not regioselective [159] and results in the modification of phenolic groups responsible for the antioxidant activity of flavonoids [158]. The enzyme groups used for flavonoid transformation are presented in Table 4.

Enzymes Type		Transformation	Reference	
Esterases	Esterases (carboxyl esterases)	Reaction similar to lipases, but with short-chain fatty acids and difference in the interfacial activation. Low practical applications in enzymatic transformation of flavonoids with a short aliphatic chain length such as acetate, propionate, and butyrate.	[160]	
Isomerase	Chalcone isomerase (CHI)	Cyclization of chalcone to form flavanone, transformation of chalcone and 6'-deoxychalcone into (2 <i>S</i>)-naringerin and (2 <i>S</i>)-5-deoxyflavanone. Soybean CHIs do not require the 4'-hydroxy moiety on the substrate for high enzyme activity.	[161]	
Laccase	Laccase from Myceliophthora	Synthesis of a flavonoid polymer and high molecular fraction of extracted flavonoids from rutin as substrate in the mixture of methanol and buffer. Oxidation of catechin in the presence of gelatin and synthesize the gelatin-catechin conjugate.	[162,163]	
Lipase	<i>Candida antarctica</i> lipase B	Acetylation only on the primary 6'-OH of the isoquercitrin glucose and the secondary 4'-OH of the rutin rhamnose were expected to be acetylated.	[164]	
	Pseudomonas cepacea lipase	Pseudomonas cepacea lipaseAcetylation occurred only on 3'-OH, 5'-OH, and 7-OH hydroxyls.		
Pectinase and Cellulase	Commercial Cellulases from Trichoderma viride	Transglucosylation activity toward (+)-catechin and (–)-epigallocatechin gallate (EGCG) using dextrin as a glucosyl donor. EGCG glucosides were functionally superior to EGCG as food additives.	[146,165,166]	
	Pectinolytic and Cellulolytic Enzymes	Hydrolysis of main- and side-chain of polysaccharides, and glycosidase activities.		
Perovidase	Chloroperoxidase	Halogenation of naringenin and hersperetin, at C-6 and C-8 with chloride and bromide ions.	[167 168]	
1 croxicuse	Horseradish peroxidase	Conjugation of green-tea catechin with amine substituted octahedral silsesquioxane.	[107,100]	
	Alkaline protease from Bacillus subtilis	Synthesis of 3"-O-substituted vinyl rutin esters in pyridine.		
	Novozym 435	Synthesis of 4"- O-substituted vinyl rutin esters in tert-butanol.		
Protease	Proteases	Hydrolytic and synthetic functions. Enzymatic transformations of flavonoids were affected by the type, origin and concentration of enzymes, nature of flavonoids, donor and optimal conditions (temperature, substrates, and solvent).	[169–171] 	
	Subtilisin (serine protease)	Flavonoid ester synthesis, the selective rutin acylation in organic solvents with excellent selectivity. The structure of the sugar moiety affected the regioselectivity.		

 Table 4. Flavonoids transformation catalyzed by different enzymes.

Table 4. Cont.

Enzymes Type		Transformation		
	Glycosyltransferases (GTs)	Glycosylation of on one or more of five hydroxyl groups of flavonol quercetin, as well as formation of hesperetin-7-glucoside. Some natural GTs is characterized by low specificity and other by the stringent specificity for glycosylation patterns. Positions, number, and length of the sugar moieties are significant factors for yield reaction.		
Transferase	Prenyltransferase (from Morus nigra)	Exclusive prenylation of chalcones (1, 2, 3, bearing two hydroxyl groups (C-2', C-4') on ring A) with a 2', 4' dihydroxy substitution and the isoflavone genistein. The position of substituents in ring B appeared to be critical for the prenylation.	[95,172–175]	
	Prenyltransferase (NovQ):	Transferring of a dimethylallyl group to the B-ring of flavonoids. Genistein and naringenin and yielded two products with a dimethylallyl group at C-3' or O-4'.		

Lipase (Table 4) catalyzed flavonoids acylation with phenolic acids leads to increased solubility, stability, and antioxidant activity of flavonoids in different media [176]. In addition, the presence of electron-donating or withdrawing substituents in the aromatic ring of flavonoids appears to be essential for their activity in the central nervous system as anxiolytic, anticonvulsant, and sedative and skeletal muscle relaxant drugs.

Chemical modifications are complex and laborious work that requires specific conditions. Therefore, enzyme modification can be a very promising alternative technique in this way. It is a well-known enzymatic halogenation of organic compounds by means of chloroperoxidase (EC 1.11.10). Chloroperoxidase from *Caldariomyces fumago* (Table 4) and whole microbial cells were applied to halogenate the flavones, naringenin, and hesperetin, at C-6 and C-8 in the presence of chloride and bromide ions [168]. Biomodified compounds have shown similar properties compared to derivatives obtained by chemical modification using highly aggressive agents such as molecular halogens and hypohalous acid.

The use of different enzymes (Table 4) has been studied in order to find the most potent biocatalyst for the selective transformation of flavonoids (acylation, deacylation, etc.). Applications of thermostable enzymes have been reported [166]. Enzyme immobilization has been performed to increase enzyme stability, facilitate enzyme reuse, and product insulation [177].

The regioselective synthesis of phloridzin-6'-O-cinnamate has been carried out using *Candida antarctic* lipase B immobilized with a macroporous acrylic resin [178,179]. Different enzymes have been used in immobilized forms to reduce the cost of enzymatic modifications of structural flavonoids due to their advantages such as easy isolation and re-use, increased enzyme stability, and regioselectivity in nonaqueous media [164].

Oxidative and conjugative biocatalyzed transformations of flavonoids may be performed in the presence of different microorganisms (*Bacillus* sp., *Aspergillus* sp., *Saccharomyces cerevisiae*, *E. coli*) [132]. Biotransformations performed by microorganisms consist of very complex mechanisms, including cyclization reactions, condensation, dehydroxylation, hydroxylation, O-dealkylation, alkylation, dehydrogenation, halogenation, double-bond reduction, carbonyl reduction, glycosylation, sulfation, dimerization, or various types of ring degradation. They were described earlier [132,180,181].

5. Human Health and Biological Properties

There is a strong link between a high intake of flavonoids and low cardiovascular disease, neurodegeneration, and cancer in the population in which a food that includes high levels of these phytochemicals is included in a daily diet [18,22,182]. These properties are based on protein interaction (enzyme, factor transcription, and receptor transcription) [22]. The use of herbs with a high content of flavonoids has been used as part of traditional remedies to treat various diseases that improve the immune system, as shown by antioxidants, anti-inflammatory, anti-allergenic, and antithrombotic drugs (Figure 5) [19,183,184]. There are some reports indicating the pro-oxidant activity of flavonoids [185]. Flavonoids generally act to protect plants from UV-B radiation and have the potential to reduce oxidative damage, act as photoreceptors, and have the power to attract flower pigment pollinators and protect against pathogen attack [19,23,186].



Figure 5. Different biological properties of plant flavonoids for human health and disease.

In particular, isoflavones have shown a variety of effects on human health, especially in women's health, such as the prevention and reduction of climate symptoms (such as hot flushes). The control of diseases such as diabetes, osteoporosis, and breast cancer has also been reported to be associated with the consumption of food rich in isoflavones [28,30,31]. In addition, even for prostate cancer isoflavones have shown their chemical-prevention effect [29].

6. Concluding Remarks

Flavonoids are a category of compounds with diverse biological activities that are of interest to various industrial areas, such as food, health pharmaceuticals, and cosmetics. Due to consumers demands today for products that benefit their quality of life substantially. Several extraction methods for the recovery of these flavonoids have been proposed for its wide range of applications. Today, microwave, ultrasound, and supercritical fluid technologies are among the most used methods of flavonoid extraction by most researchers worldwide. In addition, the use of biotechnological alternatives has attracted the attention of research to finding cleaner and more efficient ways to recover flavonoids. These methodologies have been distinguished from conventional methods by a marked increase in extraction yields and the displacement of organic solvents by "green" solvents, as well as a significant decrease in extraction times. Most studies have reported the extraction of flavonoids from vegetable sources such as fruits, seeds, roots or by-products of food, and/or beverage processing such as peels. Although progress has been made in this area, it is necessary to redefine the extractive techniques, since the majority is focused on increasing the yield of total flavonoids, leaving aside the purity of the extracts. The acquisition of extracts and/or purer flavonoid compounds would allow us to

know more precisely the biological activities and thus to understand the mechanisms of action and the effects on their consumption, allowing for greater diversification of their applications and generating more knowledge in this area.

Author Contributions: Conceptualization, M.L.C.-G. and C.N.A.; literature survey, organization, and critical analysis of data, M.L.C.-G., L.S., H.A.L.-G., L.V.R.-D., A.I., and C.N.A.; initial writing, M.L.C.-G.; revisions and proofreading, M.L.C.-G., D.K.V., L.V.R.-D., and C.N.A.; funding acquisition, M.L.C.-G. and C.N.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Guo, L.; Gao, L.; Ma, X.; Guo, F.; Ruan, H.; Bao, Y.; Wang, Y. Functional analysis of flavonoid 3'-hydroxylase and flavonoid 3',5'-hydroxylases from tea plant (*Camellia sinensis*), involved in the B-ring hydroxylation of flavonoids. *Gene* 2019, 717, 144046. [CrossRef]
- 2. Raffa, D.; Maggio, B.; Raimondi, M.V.; Plescia, F.; Daidone, G. Recent discoveries of cancer flavonoids. *EJMC* **2017**, 142, 213–228.
- 3. Aherne, S.A.; O'Brien, N.M. Dietary flavonols: Chemistry, food content, and metabolism. *Nutrition* **2002**, *18*, 75–81. [CrossRef]
- 4. Amri, F.S.; Hossain, M.A. Comparison of total phenols, flavonoids and antioxidant potential of local and imported ripe bananas. *EJBAS* **2018**, *5*, 245–251. [CrossRef]
- 5. Miao, M.; Cao, L.; Xu, K.; Xin, W.; Zheng, Y. Intervention action of total flavonoids from root of Ilex pubescens in cerebral ischemic tolerance with blood stasis. *Saudi J. Biol. Sci.* **2017**, *24*, 729–736. [CrossRef]
- 6. Moon, K.M.; Lee, B.; Cho, W.K.; Lee, B.S.; Kim, C.K.; Ma, J.Y. Swertiajaponin as an anti-browning and antioxidant flavonoid. *Food Chem.* **2018**, 252, 207–214. [CrossRef]
- 7. Diwan, V.; Brown, L.; Gobe, G.C. The flavonoid rutin improves kidney and heart structure and function in an adenine-induced rat model of chronic kidney disease. *J. Funct. Foods* **2017**, *33*, 85–93. [CrossRef]
- Geleijnse, J.M.; Launer, L.J.; Van der Kuip, D.A.; Hofman, A.; Witteman, J.C. Inverse association of tea and flavonoid intakes with incident myocardial infarction: The Rotterdam study. *AJCN* 2002, 75, 880–886. [CrossRef]
- 9. Spagnuolo, C.; Moccia, S.; Ruso, G.L. Anti-inflammatory effects of flavonoids in neurodegenerative disorders. *EJMC* **2018**, 153, 105–115. [CrossRef]
- 10. Stahl, W.; Ale-Agha, N.; Polidori, M.C. Non-antioxidant properties of carotenoids. *JBC* **2002**, *383*, 553–558. [CrossRef]
- 11. Xu, M.S.; Chen, S.; Wang, W.Q.; Liu, S.Q. Employing bifunctional enzymes for enhanced extraction of bioactives from plants: Flavonoids as an example. *J. Agric. Food Chem.* **2013**, *61*, 7941–7948. [CrossRef]
- Zuiter, A.S.; Zarqa, J. Proanthocyanidin: Chemistry and biology: From phenolic compounds to proanthocyanidins. In *Reference Module in Chemistry, Molecular Sciences and Chemical Engineering*; Reedijk, J., Ed.; Elsevier: Amsterdam, The Netherlands, 2014; pp. 1–29.
- 13. Karabin, M.; Hudcova, T.; Jelinek, L.; Dostalek, P. Biotransformation and biological activities of hop flavonoids. *Biotech. Adv.* **2015**, *33*, 1063–1090. [CrossRef]
- 14. Sharma, V.; Janmeda, P. Extraction, isolation and identification of flavonoid from *Euphorbia neriifolia* leaves. *AJC* **2017**, *10*, 509–514. [CrossRef]
- 15. Arora, S.; Itankar, P. Extraction, isolation and identification of flavonoid from *Chenopodium album* aerial parts. *JTCM* **2018**, *8*, 476–482. [CrossRef]
- 16. Martens, S.; Mithöfer, A. Flavones and flavone synthases. Phytochemistry 2005, 66, 2399–2407. [CrossRef]
- 17. De Villiers, A.; Venter, P.; Pasch, H. Recent advances and trends in the liquid-chromatography-mass spectrometry analysis of flavonoids. *J. Chromatogr. A* **2016**, *1430*, 16–78. [CrossRef]
- 18. Olagaray, K.E.; Bradford, B.J. Plant flavonoids to improve productivity of ruminants—A review. *Anim. Feed Sci. Tech.* **2019**, *251*, 21–36. [CrossRef]

- Ververidis, F.; Trantas, E.; Douglas, C.; Vollmer, G.; Kretzschmar, G.; Panopoulos, N. Biotechnology of flavonoids and other phenylpropanoid-derived natural products. Part I: Chemical diversity, impacts on plant biology and human health. *Biotechnol. J.* 2007, *2*, 1214–1234. [CrossRef]
- 20. Kale, A.; Gawande, S.; Kotwal, S. Cancer phytotherapeutics: Role for flavonoids at the cellular level. *Phytother. Res.* **2008**, *22*, 567–577. [CrossRef]
- 21. Cassidy, A.; Kay, C. Phytochemicals: Classification and occurrence. In *Encyclopedia of Human Nutrition*; Caballero, B., Ed.; Elsevier: Amsterdam, The Netherlands, 2013; pp. 39–46.
- 22. Rybarczyk-Plonska, A.; Wold, A.B.; Bengtsson, G.B.; Borge, G.I.A.; Hansen, M.K.; Hagen, S.F. Flavonols in broccoli (*Brassica oleracea* L. var. *italica*) flower buds as affected by postharvest temperature and radiation treatments. *Postharvest Biol. Technol.* **2016**, *116*, 105–114. [CrossRef]
- Cui, B.; Hu, Z.; Zhang, Y.; Hu, J.; Yin, W.; Feng, Y.; Xie, Q.; Chen, G. Anthocyanins and flavonols are responsible for purple color of *Lablab purpureus* (L.) sweet pods. *Plant Physiol. Biochem.* 2016, 103, 183–190. [CrossRef] [PubMed]
- 24. Xu, Y.; Simon, J.E.; Ferruzzi, M.G.; Ho, L.; Pasinetti, G.M.; Wu, Q. Quantification of anthocyanidins in the grapes and grape juice products with acid assisted hydrolysis using LC/MS. *JFF* **2012**, *4*, 710–717. [CrossRef]
- 25. Oliveira, J.; Mateus, N.; de Freitas, V. Flavanols: Catechins and proanthocyanidins. In *Natural Products*; Ramawat, K.G., Mérillon, J.M., Eds.; Springer: Berlin/Heidelberg, Germany, 2013; pp. 1753–1801.
- 26. Pons, Z.; Margalef, M.; Bravo, F.I.; Arola-Arnal, A.; Muguerza, B. Grape seed flavanols decrease blood pressure *via* Sirt-1 and confer a vasoprotective pattern in rats. *JFF* **2016**, *24*, 164–172. [CrossRef]
- 27. Vuong, Q.V.; Golding, J.B.; Nguyen, M.; Roach, P.D. Extraction and isolation of catechins from tea. *JSS* **2010**, 33, 3415–3428. [CrossRef] [PubMed]
- Zhao, C.C.; Kim, P.H.; Eun, J.B. Influence of high-intensity ultrasound application on the physicochemical properties, isoflavones composition and antioxidant activity of tofu whey. *LWT-Food Sci. Technol.* 2020, 117, 108618. [CrossRef]
- 29. Duru, K.C.; Kovaleva, E.; Danilova, I.; Belousova, A. Production and assessment of novel probiotic fermented oat flour enriched with isoflavones. *LWT-Food Sci. Technol.* **2019**, *117*, 108618. [CrossRef]
- 30. Nemitz, M.C.; Argenta, D.F.; Koester, L.S.; Bassani, V.L.; von Poser, G.L.; Teixeira, H.F. The international scenario of patents concerning isoflavones. *Trends Food Sci. Technol.* **2016**, *49*, 85–95. [CrossRef]
- Ganzera, M. Supercritical fluid chromatography for the separation of isoflavones. *J. Pharm. Biomed. Anal.* 2015, 107, 364–369. [CrossRef]
- 32. Alara, O.R.; Abdurahman, N.H.; Ukaegbu, C.I. Soxhlet extraction pf phenolic compounds from *Vernonia cinereal* leaves and its antioxidant activity. *J. Appl. Res. Med. Aromat. Plants* **2018**, *11*, 12–17.
- 33. Sati, P.; Dhyani, P.; Bhatt, I.D.; Pandey, A. Ginkgo biloba flavonoid glycosides in antimicrobial perspective with reference to extraction method. *JTCM* **2019**, *9*, 15–23. [CrossRef]
- Mackela, I.; Andriekus, T.; Venskutonis, P.R. Biorefinning of buckwheat (Fagopyru, esculentum) hulls by using supercritical fluid, Soxhlet, pressurized liquid and enzyme-assisted extraction methods. *JFE* 2017, 213, 38–46. [CrossRef]
- 35. Cujic, N.; Svikin, K.; Jankovic, T.; Pljevljakusic, D.; Zdunic, G.; Ibric, S. Optimization of polyphenols extraction from dried chokeberry using maceration as traditional technique. *Food Chem.* **2016**, *194*, 135–142. [CrossRef]
- Pereira, S.V.; Reis, R.A.S.P.; Garbuio, D.C.; de Freitas, L.A.P. Dynamic maceration of *Matricaria chamomilla* inflorescences: Optimal conditions for flavonoids and antioxidant activity. *Rev. Bras. Farmacogn.* 2018, 28, 111–117. [CrossRef]
- 37. Agustin-Salazar, S.; Medina-Juárez, L.A.; Soto-Valdez, H.; Manzanares-López, F.; Gámez-Meza, N. Influence of the solvent system on the composition of phenolic substances and antioxidant capacity of extracts of grape (*Vitis vinifera* L.) marc. *AJGWR* **2014**, *R20*, 208–213. [CrossRef]
- Albuquerque, B.R.; Prieto, M.A.; Vazquez, J.A.; Barreirio, M.F.; Barros, L.; Ferreira, I.C.F.R. Recovery of bioactive compounds form Arbutus unedo L. fruits: Comparative optimization study of maceration/ microwave/ultrasound extraction techniques. *Food Res. Int.* 2018, 109, 455–471. [CrossRef]
- 39. Farzaneth, V.; Carvalho, I.S. Modelling of Microwave Assisted Extraction (MAE) of Anthocyanins (TMA). *JARMAP* 2017, *6*, 92–100.
- 40. Ye, C.L.; Liu, X.G. Extraction of flavonoids from *Tetrastigma hemsleyanum* diels et gilg and their antioxidant activity. *JFPP* **2015**, *39*, 2197–2205.

- Safdar, M.N.; Kausar, T.; Jabbar, S.; Mumtaz, A.; Ahad, K.; Saddozai, A.A. Extraction and quantification of polyphenols from kinnow (*Citrus reticulate* L.) peel using ultrasound and maceration techniques. *JFDA* 2017, 25, 488–500. [CrossRef]
- 42. Bergeron, C.; Gafner, S.; Clausen, E.; Carrier, D.J. Comparison of the Chemical Composition of Extracts from Scutellaria lateriflora Using Accelerated Solvent Extraction and Supercritical Fluid Extraction versus Standard Hot Water or 70% Ethanol Extraction. *J. Agric. Food Chem.* **2005**, *53*, 3076–3080. [CrossRef]
- 43. Yu, M.; Wang, B.; Qi, Z.; Xin, G.; Li, Q. Response Surface method was used to optimize the ultrasonic assisted extraction of flavonoids from *Crinum asiaticum*. *Saudi J. Biol. Sci.* **2019**, *26*, 2079–2084. [CrossRef]
- 44. Daud, M.N.H.; Fatanah, D.N.; Abdullah, N.; Ahmad, R. Evaluation of antioxidant potential of *Artocarpus heterophyllus* L. J33 variety fruit waste from different extraction methods and identification of phenolic constituents by LCMS. *Food Chem.* **2017**, 232, 621–632. [CrossRef] [PubMed]
- 45. Szewczyk, K.; Olech, M. Optimization of extraction method for LC-MS based determination of phenolic acid profiles in different Impatiens species. *Phytochem. Lett.* **2017**, *20*, 322–330. [CrossRef]
- Wu, X.; Beecher, G.R.; Holden, J.M.; Haytowitz, D.B.; Gebhardt, S.E.; Prior, R.L. Lipophilic and Hydrophilic Antioxidant Capacities of Common Foods in The United States. J. Agric. Food Chem. 2004, 52, 4026–4037. [CrossRef]
- 47. Rafińska, K.; Pomastowski, P.; Rudnicka, J.; Krakowska, A.; Maruska, A.; Narkute, M.; Buszewski, B. Effect of solvent and extraction technique on composition and biological activity of Lepidium sativum extracts. *Food Chem.* **2019**, *289*, 16–25. [CrossRef] [PubMed]
- 48. Ku, C.S.; Jang, J.P.; Mun, S.P. Exploitation of polyphenol-rich pine barks for potent antioxidant activity. *J. Wood Sci.* **2007**, *53*, 524–528. [CrossRef]
- Bahorun, T.; Luximon-Ramma, A.; Crozier, A.; Aruoma, O.I. Total phenol, flavonoid, proanthocyanidin and vitamin C levels and antioxidant activities of Mauritian vegetables. *J. Sci. Food Agric.* 2004, *84*, 1553–1561. [CrossRef]
- 50. Chu, Y.-F.; Sun, J.; Wu, X.; Liu, R.H. Antioxidant and Antiproliferative Activities of Common Vegetables. *J. Agric. Food Chem.* **2002**, *50*, 6910–6916. [CrossRef]
- 51. Zhang, D.; Hamauzu, Y. Phenolics, ascorbic acid, carotenoids and antioxidant activity of broccoli and their changes during conventional and microwave cooking. *Food Chem.* **2004**, *88*, 503–509. [CrossRef]
- 52. Anokwuru, C.P.; Anyasor, G.N.; Ajibaye, O.; Fakoya, O.; Okebugwu, P. Effect of extraction solvents on phenolic, flavonoid and antioxidant activities of three nigerian medicinal plants. *Nat. Sci.* **2011**, *9*, 53–61.
- 53. Vieitez, I.; Maceiras, L.; Jachmanián, I.; Alborés, S. Antioxidant and antibacterial activity of different extracts from herbs obtained by maceration or supercritical technology. *JSF* **2018**, *133*, 58–64. [CrossRef]
- 54. Moreira, M.M.; Barroso, M.F.; Boeykens, A.; Withouck, H.; Morais, S.; Delerue-Matos, C. Valorization of apple tree wood residues by polyphenols extraction: Comparison between conventional and microwave-assisted extraction. *Ind. Crop. Prod.* **2017**, *104*, 210–220. [CrossRef]
- 55. Aspé, E.; Fernández, K. The effect of different extraction techniques on extraction yield, total phenolic, and anti-radical capacity of extracts from *Pinus radiata* Bark. *Ind. Crop. Prod.* **2011**, *34*, 838–844. [CrossRef]
- 56. Lamounier, K.C.; Cunha, L.C.S.; de Morais, S.A.L.; de Aquino, F.J.T.; Chang, R.; do Nascimento, E.A.; de Souza, M.G.M.; Martins, C.H.G.; Cunha, W.R. Chemical analysis and study of phenolics, antioxidant activity, and antibacterial effect of the wood and bark of *Maclura tinctoria* (L.) D. Don ex Steud. *Altern. Med.* **2012**, 2012, 1–7.
- 57. Zhu, H.B.; Wang, Y.Z.; Liu, Y.X.; Xia, Y.L.; Tang, T. Analysis of flavonoids in *Portulaca oleracea* L. by uv-vis spectrophotometry with comparative study on different extraction technologies. *Food Anal. Methods* **2010**, *3*, 90–97. [CrossRef]
- 58. Li, H.; Deng, Z.; Wu, T.; Liu, R.; Loewen, S.; Tsao, R. Microwave-assisted extraction of phenolics with maximal antioxidant activities in tomatoes. *Food Chem.* **2012**, *130*, 928–936. [CrossRef]
- 59. Cardoso-Ugarte, G.A.; Juárez-Becerra, G.P.; Sosa-Morales, M.E.; López-Malo, A. Microwave-assisted Extraction of Essential Oils from Herbs. *JMPEE* **2013**, *47*, 63–72. [CrossRef]
- 60. Casazza, A.A.; Aliakbarian, B.; Mantegna, S.; Cravotto, G.; Perego, P. Extraction of phenolics from *Vitis vinifera* wastes using non-conventional techniques. *J. Food Eng.* **2010**, *100*, 50–55. [CrossRef]
- 61. Pan, Y.M.; He, C.H.; Wang, H.S.; Ji, X.W.; Wang, K.; Liu, P.Z. Antioxidant activity of microwave-assisted extract of *Buddleia officinalis* and its major active component. *Food Chem.* **2010**, *121*, 497–502. [CrossRef]

- 62. Feng, R.Z.; Wang, Q.; Tong, W.Z.; Xiong, J.; Wei, Q.; Zhou, W.H.; Yin, Z.Q.; Yin, X.Y.; Wang, L.Y.; Chen, Y.Q.; et al. Extraction and antioxidant activity of flavonoids of *Morus nigra*. *Int. J. Clin. Exp. Med.* **2015**, *8*, 22328–22336.
- Alara, O.R.; Abdurahman, N.H.; Olalere, O.A. Optimization of microwave-assisted extraction of flavonoids antioxidants from *Vernonia amygdalina* leaf using response surface methodology. *Food Bioprod. Process.* 2018, 107, 36–48. [CrossRef]
- 64. Bimakr, M.; Rahman, R.A.; Taip, F.S.; Ganjloo, A.; Salleh, L.M.; Selamat, J.; Hamid, A.; Zaidul, I.S.M. Comparison of different extraction methods for the extraction of major bioactive flavonoid compounds from spearmint (*Mentha spicata* L.) leaves. *Food Bioprod. Process.* **2011**, *89*, 67–72. [CrossRef]
- Hartonen, K.; Parshintsev, J.; Sandberg, K.; Bergelin, E.; Nisula, L.; Riekkola, M.L. Isolation of flavonoids from aspen knotwood by pressurized hot water extraction and comparison with other extraction techniques. *Talanta* 2007, 74, 32–38. [CrossRef] [PubMed]
- 66. Jha, P.; Das, A.J.; Deka, S.C. Optimization of ultrasound and microwave assisted extractions of polyphenols from black rice (*Oryza sativa* cv. Poireton) husk. *J. Food Sci. Technol.* **2017**, *54*, 3847–3858. [CrossRef] [PubMed]
- Rodríguez-Pérez, C.; Gilbert-López, B.; Mendiola, J.A.; Quirantes-Piné, R.; Segura-Carretero, A.; Ibáñez, E. Optimization of microwave-assisted extraction and pressurized liquid extraction of phenolic compounds from *Moringa oleifera* leaves by multi-response surface methodology. *Electrophoresis* 2016, *37*, 1938–1946. [CrossRef]
- Zekovic, Z.; Vladic, J.; Vidovic, S.; Adamovic, D.; Pavlic, B. Optimization of microwave-assisted extraction (MAE) of coriander phenolic antioxidants—Response surface methodology approach. *J. Sci. Food Agric.* 2016, *96*, 4613–4622. [CrossRef] [PubMed]
- 69. Bouras, M.; Chadni, M.; Barba, F.J.; Grimi, N.; Bals, O.; Vorobiev, E. Optimization of microwave-assisted extraction of polyphenols from Quercus bark. *Ind. Crop. Prod.* **2015**, *77*, 590–601. [CrossRef]
- 70. The, S.S.; Niven, B.E.; Bekit, A.E.A.; Carne, A.; Birch, E.J. Microwave and pulsed electric field extraction of polyphenols from defatted canola seeds cake. *Int. J. Food Sci. Technol.* **2015**, *50*, 1109–1115.
- 71. Liazid, A.; Guerrero, R.F.; Cantos, E.; Palma, M.; Barroso, C.G. Microwave assisted extraction of anthocyanins from grape skins. *Food Chem.* **2011**, *124*, 1238–1243. [CrossRef]
- Tan, S.N.; Yong, J.W.H.; Teo, C.C.; Ge, L.; Chan, Y.W.; Hew, C.S. Determination of metabolites in Uncaria sinensis by HPLC and GC-MS after green solvent microwave-assisted extraction. *Talanta* 2011, *83*, 891–898. [CrossRef]
- 73. Inoue, T.; Tsubaki, S.; Ogawa, K.; Onishi, K.; Azuma, J.I. Isolation of hesperidin from peels of thinned Citrus unshiu fruits by microwave-assisted extraction. *Food Chem.* **2010**, *123*, 542–547. [CrossRef]
- 74. Li, J.; Zu, Y.G.; Fu, Y.J.; Yang, Y.C.; Li, S.M.; Li, Z.N.; Wink, M. Optimization of microwave-assisted extraction of triterpene saponins from defatted residue of yellow horn (*Xanthoceras sorbifolia* Bunge.) kernel and evaluation of its antioxidant activity. *Innov. Food Sci. Emerg. Technol.* **2010**, *11*, 637–643. [CrossRef]
- 75. Ballard, T.S.; Mallikarjunan, P.; Zhou, K.; O'Keefe, S. Microwave-assisted extraction of phenolic antioxidant compounds from peanut skins. *Food Chem.* **2010**, *120*, 1185–1192. [CrossRef]
- Kong, Y.; Zu, Y.-G.; Fu, Y.-J.; Liu, W.; Chang, F.-R.; Li, J.; Chen, Y.-H.; Zhang, S.; Gu, C.-B. Optimization of microwave-assisted extraction of cajaninstilbene acid and pinostrobin from pigeonpea leaves followed by RP-HPLC-DAD determination. *J. Food Compos. Anal.* 2010, 23, 382–388. [CrossRef]
- Yang, Z.; Zhai, W. Optimization of microwave-assisted extraction of anthocyanins from purple corn (*Zea mays* L.) cob and identification with HPLC–MS. *Innov. Food Sci. Emerg. Technol.* 2010, 11, 470–476. [CrossRef]
- 78. Wang, Y.L.; Xi, G.S.; Zheng, Y.C.; Miao, F.S. Microwave-assisted extraction of flavonoids from Chinese herb *Radix puerariae* (Ge Gen). *J. Med. Plants Res.* **2010**, *4*, 304–308.
- Périno-Issartier, S.; Zille, H.; Abert-Vian, M.; Chemat, F. Solvent free microwave-assisted extraction of antioxidants from sea buckthorn (*Hippophae rhamnoides*) food by-products. *Food Bioprocess. Technol.* 2010, 4, 1020–1028. [CrossRef]
- 80. Tsubaki, S.; Sakamoto, M.; Azuma, J. Microwave-assisted extraction of phenolic compounds from tea residues under autohydrolytic conditions. *Food Chem.* **2010**, *123*, 1255–1258. [CrossRef]
- 81. Victório, C.P.; Lage, C.L.S.; Kuster, R.M. Flavonoid extraction from *Alpinia zerumbet* (Pers.) Burtt et Smith leaves using different techniques and solvents. *Eclet. Quim.* **2009**, *34*, 19–24.

- Yang, L.; Jiang, J.-G.; Li, W.-F.; Chen, J.; Wang, D.-Y.; Zhu, L. Optimum extraction Process of polyphenols from the bark of Phyllanthus emblica L. based on the response surface methodology. *J. Sep. Sci.* 2009, 32, 1437–1444. [CrossRef]
- 83. Hayat, K.; Hussain, S.; Abbas, S.; Farooq, U.; Ding, B.; Xia, S.; Jia, C.; Zhang, X.; Xia, W. Optimized microwave-assisted extraction of phenolic acids from citrus mandarin peels and evaluation of antioxidant activity *in vitro*. *Sep. Purif. Technol.* **2009**, *70*, 63–70. [CrossRef]
- Zheng, X.; Wang, X.; Lan, Y.; Shi, J.; Xue, S.J.; Liua, C. Application of response surface methodology to optimize microwave-assisted extraction of silymarin from milk thistle seeds. *Sep. Purif. Technol.* 2009, 70, 34–40. [CrossRef]
- 85. Li, W.; Li, T.; Tang, K. Flavonoids from mulberry leaves by microwave-assisted extract and anti-fatigue activity. *Afr. J. Agric. Res.* **2009**, *4*, 898–902.
- 86. Vian, M.A.; Maingonnat, J.F.; Chemat, F. Clean recovery of antioxidant flavonoids from onions: Optimising solvent free microwave extraction method. *J. Chroma A* **2009**, *1216*, 7700–7707.
- 87. Chen, L.; Jin, H.; Ding, L.; Zhang, H.; Li, J.; Qu, C. Dynamic microwave-assisted extraction of flavonoids from *Herba epimedii*. *Sep. Purif. Technol.* **2008**, *59*, 50–57. [CrossRef]
- 88. Wang, J.X.; Xiao, X.H.; Li, G.K. Study of vacuum microwave-assisted extraction of polyphenolic compounds and pigment from Chinese herbs. *J. Chroma A* **2008**, *1198–1199*, 45–53. [CrossRef] [PubMed]
- Xiao, W.; Han, L.; Shi, B. Microwave-assisted extraction of flavonoids from Radix astragali. *Sep. Purif. Technol.* 2008, *62*, 614–618. [CrossRef]
- Chen, L.G.; Ding, L.; Yu, A.M.; Yang, R.L.; Wang, X.P.; Li, J.T.; Jin, H.Y.; Zhang, H.Q. Continuous determination of total flavonoids in *Platycladus orientalis* (L.) Franco by dynamic microwave-assisted extraction coupled with on-line derivatization and ultraviolet-visible detection. *Anal. Chim. Acta* 2007, 596, 164–170. [CrossRef]
- Liu, Z.; Ding, L.; Zhang, H.; Hu, X.; Bu, F. Comparison of the different extraction methods of flavonoids in Epimedium koreamum Nakai by HPLC-DAD-ESI-MSn. J. Liq. Chromatogr. Relat. Tech. 2006, 29, 719–731. [CrossRef]
- Japón-Luján, R.; Luque-Rodríguez, J.M.; Luque de Castro, M.D. Multivariate optimisation of the microwave-assisted extraction of oleuropein and related biophenols from olive leaves. *Anal. Bioanal. Chem.* 2006, 385, 753–759.
- 93. Liu, Z.; Yan, G.; Bu, F.; Sun, J.; Hu, X.; Zhang, H. Analysis of chemical composition of *Acanthopanax senticosusleaves* applying high-pressure microwave-assisted extraction. *Chem. Anal.* **2005**, *50*, 851–861.
- 94. Li, H.; Chen, B.; Zhang, Z.; Yao, S. Focused microwave-assisted solvent extraction and HPLC determination of effective constituents in Eucommia ulmodies Oliv. (*E. ulmodies*). *Talanta* **2004**, *63*, 659–665. [CrossRef]
- 95. Williams, G.J.; Zhang, C.; Thorson, J.S. Expanding the promiscuity of a natural-product glycosyltransferase by directed evolution. *Nat. Chem. Biol.* **2007**, *3*, 657–662. [CrossRef] [PubMed]
- 96. Ferreira, T.M.S.; dos Santos, J.A.; Modesto, L.A.; Souza, L.S.; dos Santos, M.P.; Bezerra, D.G.; de Paula, J.A.M. An eco-friendly method for extraction and quantification of flavonoids in *Dysphania ambrosioides*. *Braz. J. Pharmacogn.* 2019, 29, 266–270. [CrossRef]
- Um, M.; Han, T.H.; Lee, J.W. Ultrasound-assisted extraction and antioxidant activity of phenolic and flavonoid compounds and ascorbic acid from rugosa rose (*Rosa rugosa* Thunb.) fruit. *Food Sci. Biotech.* 2018, 27, 375–382. [CrossRef]
- Maran, J.P.; Manikandan, S.; Nivetha, C.V.; Dinesh, R. Ultrasound assisted extraction of bioactive compounds from *Nephelium lappaceum* L. fruit peel using central composite face centered response surface design. *Arab. J. Chem.* 2017, 10, S1145–S1157. [CrossRef]
- Ghasemzadeh, A.; Jaafar, H.; Karimi, E.; Rahmat, A. Optimization of ultrasound-assisted extraction of flavonoid compounds and their pharmaceutical activity from curry leaf (*Murraya koenigii* L.) using response surface methodology. *BMC Complement. Altern. Med.* 2014, 14, 318. [CrossRef] [PubMed]
- Wang, C.; Li, Y.; Yao, L.; Wu, G.; Chang, J.; Shu, C.; Chen, M. Optimization of ultrasonic-assisted extraction of flavonoid from *Portulaca oleracea* L. by response surface methodology and chemical composition analysis. *J. Korean Soc. Appl. Biol. Chem.* 2014, 57, 647–653. [CrossRef]
- 101. Memon, A.A.; Memon, N.; Luthria, D.L.; Bhanger, M.I.; Pitafi, A.A. Phenolic Acids Profiling and Antioxidant Potential of Mulberry (*Morus laevigata W., Morus nigra L., Morus alba L.*) Leaves and Fruits Grown in Pakistan. *Pol. J. Food Nutr. Sci.* 2010, 60, 25–32.

- 102. Babova, O.; Occhipintia, A.; Capuzzo, A.; Maffei, M.E. Extraction of bilberry (Vaccinium myrtillus) antioxidants using supercritical/subcritical CO₂ and ethanol as co-solvent. *J. Supercrit. Fluids* **2016**, 107, 358–363. [CrossRef]
- Wang, L.; Yang, B.; Du, X.; Yi, C. Optimisation of supercritical extraction of flavonoids from Pueraria lobata. *Food Chem.* 2008, 108, 737–741. [CrossRef]
- 104. Chen, Y.; Xie, M.-Y.; Gong, X.F. Microwave-assisted extraction used for the isolation of total triterpenoid saponins from *Ganoderma atrum*. J. Food Eng. 2007, 81, 162–170. [CrossRef]
- Flórez, N.; Conde, E.; Domínguez, H. Microwave assisted water extraction of plant compounds. J. Chem. Technol. Biotech. 2015, 90, 590–607. [CrossRef]
- 106. Krishnan, R.Y.; Chandran, M.N.; Vadivel, V.; Rajan, K.S. Insights on the influence of microwave irradiation on the extraction of flavonoids from *Terminalia chebula*. *Sep. Purif. Technol.* **2016**, 170, 224–233. [CrossRef]
- 107. Cassol, L.; Rodrigues, E.; Noreña, C.P.Z. Extracting phenolic compounds from *Hibiscus sabdariffa* L. calyx using microwave assisted extraction. *Ind. Crop. Prod.* **2019**, 133, 168–177. [CrossRef]
- 108. Akbari, S.; Abdurahman, N.H.; Yunus, R.M.; Fayaz, F. Microwave-assisted extraction of saponin, phenolic and flavonoid compounds from *Trigonella foenum-graecum* seed based on two level factorial design. *J. Appl. Res. Med. Aromat. Plants* 2019, 14, 100212. [CrossRef]
- 109. Ling, Y.Y.M.; Fun, P.S.; Yeop, A.; Yusoff, M.M.; Gimbun, J. Assessment of maceration, ultrasonic and microwave assisted extraction for total phenolic content, total flavonoid content and Kaempferol yield form *Cassia alata* via Microstructures Analysis. *Mater. Today Proc.* 2019, *19*, 1273–1279. [CrossRef]
- 110. Tomaz, I.; Maslov, L.; Stupic, D.; Prenier, D.; Asperger, D.; Karoglan Kontić, J. Multi-response optimization of ultrasound-assisted extraction for recovery of flavonoids from red grape skins using response surface methodology. *Phytochem. Anal.* **2015**, *27*, 13–22. [CrossRef]
- 111. Castro-López, C.; Rojas, R.; Sánchez-Alejo, E.J.; Niño-Medina, G.; Martínez-Ávila, G.C.G. Phenolic Compounds Recovery from Grapefruit and By- Products: An Overview of Extraction Methods, Grape and Wine Biotechnology, Antonio Morata and Iris Loira; IntechOpen: London, UK, 2016. [CrossRef]
- 112. Chludil, H.D.; Corbino, G.B.; Leicach, S.R. Soil Quality Effects on Chenopodium album Flavonoid Content and Antioxidant Potential. *J. Agric. Food Chem.* **2008**, *56*, 5050–5056. [CrossRef]
- 113. Wang, X.H.; Wang, J.P. Effective extraction with deep eutectic solvents and enrichment by microporous adsorption resin of flavonoids from *Carthamus tinctorius* L. J. Pharm. Biomed. **2019**, *176*, 112804. [CrossRef]
- 114. Chen, M.C.A.J.; Zhang, H.; Li, Z.; Zhao, L.; Qiu, H. Effective extraction of flavonoids from Lycium barbarum L. fruits by deep eutectic solvents-based ultrasound-assisted extraction. *Talanta* **2019**, *203*, 16–22.
- 115. Singanusong, R.; Nipornram, S.; Tochampa, W.; Rattanatraiwong, P. Low Power Ultrasound-Assisted Extraction of Phenolic Compounds from Mandarin (*Citrus reticulata* Blanco cv. Sainampueng) and Lime (*Citrus aurantifolia*) Peels and the Antioxidant. *Food Anal. Methods* **2015**, *8*, 1112–1123. [CrossRef]
- 116. Piana, F.; Ciulu, M.; Quirantes-Piné, R.; Sanna, G.; Segura-Carretero, A.; Spano, N.; Mariani, A. Simple and rapid procedures for the extraction of bioactive compounds from Guayule leaves). *Ind. Crop. Prod.* 2018, 116, 162–169. [CrossRef]
- 117. Nagendra-Prasad, K.; Yang, B.; Zhao, M.; Ruenroengklin, N.; Jiang, Y. Application of ultrasonication or high-pressure extraction of flavonoids from litchi fruit pericarp. *J. Food Process. Eng.* 2009, 32, 828–843. [CrossRef]
- Zhang, Q.A.; Fan, X.H.; Li, T.; Zhang, Z.Q.; Liu, Y.K.; Li, X.P. Optimization of ultrasound extraction for flavonoids from semen astragali complanati and its identification by HPLC-DAD-MS/MS. *Int. J. Food Sci. Technol.* 2013, 48, 1970–1976. [CrossRef]
- 119. Xie, Z.; Sun, Y.; Lam, S.; Zhao, M.; Liang, Z.; Yu, X.; Yang, D.; Xu, X. Extraction and isolation of flavonoid glycosides from Flos Sophorae Immaturus using ultrasonic-assisted extraction followed by high-speed countercurrent chromatography. *J. Sep. Sci.* **2014**, *37*, 957–965. [CrossRef] [PubMed]
- Dassoff, E.S.; Li, Y.O. Mechanisms and effects of ultrasound-assisted supercritical CO₂ extraction. *Trends Food Sci. Tech.* 2019, *86*, 492–501. [CrossRef]
- 121. Román Páez, M.; Rivera Narváez, C.; Cardona Bermúdez, L.; Muñoz, L.; Gómez, D.; Passaro Carvalho, C.; Quiceno Rico, J. *Guía de Extracción por Fluidos Supercríticos: Fundamentos y Aplicaciones*; Servicio Nacional de Aprendizaje—SENA: Rionegro, Colombia, 2016.

- 122. Verma, D.K.; Dhakane, J.P.; Mahato, D.K.; Billoria, S.; Bhattacharjee, P.; Srivastav, P.P. Supercritical Fluid Extraction (SCFE) for Rice Aroma Chemicals: Recent and Advance Extraction Method. In *Science and Technology of Aroma, Flavour and Fragrance in Rice*; Verma, D.K., Srivastav, P.P., Eds.; Apple Academic Press: Waretown, NJ, USA, 2018; pp. 179–198.
- Belbaki, A.; Louaer, W.; Menial, A.H. Supercritical CO₂ extraction of oil from Crushed Algerian olives. J. Supercrit. Fluids 2017, 130, 165–171. [CrossRef]
- 124. Kavoura, D.; Kyriakopoulou, K.; Papaefstahiou, G.; Spanidi, E.; Gardikis, K.; Louli, V.; Aligiannis, N.; Krokida, M.; Magoulas, K. Supercritical CO₂ extraction of Salvia fruticosa. *J. Supercrit. Fluids* 2019, 146, 159–164. [CrossRef]
- 125. Song, L.; Liu, P.; Yan, Y.; Huang, Y.; Bai, B.; Hou, X.; Zhang, L. Supercritical CO₂ fluid extraction of flavonoid compounds from Xinjiang jujube (Ziziphus jujube Mill.) leaves and associated biological activities and flavonoid compositions. *Ind. Crop. Prod.* 2019, 139, 111508. [CrossRef]
- 126. Panja, P. Green extraction methods of food polyphenols from vegetable materials. *Curr. Opin. Food Sci.* **2018**, 23, 173–182. [CrossRef]
- 127. Alvarez, M.; Cabred, S.; Ramírez, C.; Fanovich, M. Valorization of an agroindustrial soybean residue by supercritical flfluid extraction of phytochemical compounds. J. Supercrit. Fluids 2019, 143, 90–96. [CrossRef]
- 128. Ouédraogo, J.C.; Dicko, C.; Kini, F.; Bonzi-Coulibaly, Y.; Dey, E. Enhanced extraction of flflavonoids from Odontonema strictum leaves with antioxidant activity using supercritical carbon dioxide flfluid combined with ethanol. *J. Supercrit. Fluids* **2018**, *131*, 66–71. [CrossRef]
- Krakowska, A.; Rafinska, K.; Walczak, J.; Buszewski, B. Enzyme-assisted optimized supercritical fifluid extraction to improve Medicago sativa polyphenolics isolation. *Ind. Crop. Prod.* 2018, 124, 931–940. [CrossRef]
- Kala, H.K.; Mehta, R.; Sen, K.K.; Tandey, R.; Mandal, V. Critical analysis of research trends and issues in microwave assisted extraction of phenolics: Have we really done enough. *Trends Anal. Chem.* 2016, 85, 140–152. [CrossRef]
- 131. Ilina, A.D. The use of enzymes and microorganisms for solving fundamental and applied problems. Herald of the Russian Academy of Sciences. *Rep. Bashortostan* **2013**, *18*, 40–43.
- 132. Zha, J.; Wu, X.; Gong, G.; Koffas, M.A.G. Pathway enzyme engineering for flavonoid production in recombinant microbes. *Metab. Eng. Commun.* **2019**, *9*, e00104. [CrossRef]
- 133. Castro-Vazquez, L.; Alañón, M.E.; Rodríguez-Robledo, V.; Pérez-Coello, M.S.; Hermosín-Gutierrez, I.; Díaz-Maroto, M.C.; Jordán, J.; Galindo, M.F.; Arroyo-Jimenez, M.D.M. Bioactive flavonoids, antioxidant behaviour, and cytoprotective effects of dried grapefruit peels (*Citrus paradisi* macf.). Oxid. Med. Cell. Longev. 2016. [CrossRef]
- 134. Lagaert, S.; Belien, T.; Volckaert, G. Plant cell walls: Protecting the barrier form degradation by microbial enzymes. *Semin. Cell Dev. Biol.* **2009**, *20*, 1064–1073. [CrossRef]
- 135. Landbo, A.K.; Meyer, A.S. Enzyme-assisted extraction of antioxidative phenols from black currant juice press residues (*Ribes nigrum*). J. Agric. Food Chem. 2001, 49, 3169–3177. [CrossRef]
- 136. Kammerer, D.; Claus, A.; Schieber, A.; Carle, R. A novel process for the recovery of polyphenols from grape (*Vitis vinifera* L.) pomace. *J. Food Sci.* **2005**, *70*, C157–C163. [CrossRef]
- 137. Maier, T.; Göppert, A.; Kammerer, D.R.; Schieber, A.; Carle, R. Optimization of a process for enzyme-assisted pigment extraction from grape (*Vitis vinifera* L.) pomace. *EFRT* **2008**, 227, 267–275. [CrossRef]
- 138. Fu, Y.J.; Liu, W.; Zu, Y.G.; Tong, M.H.; Li, S.M.; Yan, M.M.; Efferth, T.; Luo, H. Enzyme assisted extraction of luteolin and apigenin from pigeonpea [*Cajanus cajan* (L.) Millsp.] leaves. *Food Chem.* 2008, 111, 508–512. [CrossRef]
- Zhang, Q.; Zhou, M.M.; Chen, P.L.; Cao, Y.Y.; Tan, X.L. Optimization of ultrasonic-assisted enzymatic hydrolysis for the extraction of luteolin and apigenin from celery. *J. Food Sci.* 2011, 76, C680–C685. [CrossRef] [PubMed]
- 140. Pan, M.; Wang, S.K.; Yuan, X.L.; Xie, R.Y.; Hong, Y.C.; Liu, H.J. Process study on the ultrasonic assisted enzymatic extraction of flavonoids from shepherd's purse. *Adv. Mater. Res.* **2013**, *634*, 1281–1286. [CrossRef]
- 141. Tchabo, W.; Ma, Y.; Engmann, F.N.; Zhang, H. Ultrasound-assisted enzymatic extraction (UAEE) of phytochemical compounds from mulberry (*Morus nigra*) must and optimization study using response surface methodology. *Ind. Crop. Prod.* 2015, *63*, 214–225. [CrossRef]

- 142. Wu, D.; Gao, T.; Yang, H.; Du, Y.; Li, C.; Wei, L.; Zhou, T.; Lu, J.; Bi, H. Simultaneous microwave/ ultrasonic-assisted enzymatic extraction of antioxidant ingredients from *Nitraria tangutorun* Bobr. juice by-products. *Ind. Crop. Prod.* **2015**, *66*, 229–238. [CrossRef]
- 143. Tomaz, I.; Maslov, L.; Stupić, D.; Preiner, D.; Ašperger, D.; Kontić, J.K. Recovery of flavonoids from grape skins by enzyme-assisted extraction. *Sep. Sci. Technol.* **2016**, *51*, 255–268. [CrossRef]
- 144. Chávez-Santoscoy, R.A.; Lazo-Vélez, M.A.; Serna-Sáldivar, S.O.; Gutiérrez-Uribe, J.A. Delivery of flavonoids and saponins from black bean (*Phaseolus vulgaris*) seed coats incorporated into whole wheat bread. *Int. J. Mol. Sci.* 2016, 17, 222. [CrossRef] [PubMed]
- 145. Xu, S.L.; Zhu, K.Y.; Bi, C.W.; Yan, L.; Men, S.W.; Dong, T.T.; Tsim, K.W. Flavonoids, derived from traditional chinese medicines, show roles in the differentiation of neurons: Possible targets in developing health food products. *Birth Defects Res. Part C Embryo Today* **2013**, *99*, 292–299. [CrossRef]
- 146. Chen, S.; Xing, X.H.; Huang, J.J.; Xu, M.S. Enzyme-assisted extraction of flavonoids from *Ginkgo biloba* leaves: Improvement effect of flavonol transglycosylation catalyzed by *Penicillium decumbens* cellulase. *Enzym. Microb. Tech.* 2011, 48, 100–105. [CrossRef]
- 147. Huynh, N.T.; Van Camp, J.; Smagghe, G.; Raes, K. Improved release and metabolism of flavonoids by steered fermentation processes: A review. *Int. J. Mol. Sci.* **2014**, *15*, 19369–19388. [CrossRef]
- Cao, H.; Chen, X.; Jassbim, A.R.; Xiao, J. Microbial biotransformation of bioactive flavonoids. *Biotech. Adv.* 2015, 33, 214–223. [CrossRef] [PubMed]
- Frauzino-Araújo, K.C.; de, M.B.; Costa, E.M.; Pazini, F.; Valadares, M.C.; de Oliveira, V. Bioconversion of quercetin and rutin and the cytotoxicity activities of the transformed products. *Food Chem. Toxicol.* 2013, 51, 93–96. [CrossRef] [PubMed]
- 150. Lee, J.H.; Oh, E.T.; Chun, S.C.; Keum, Y.S. Biotransformation of isoflavones by Aspergillus niger and Cuuninghamella elegans. *J. Korean Soc. Appl. Biol. Chem.* **2014**, *57*, 523–527. [CrossRef]
- 151. Xin, X.; Fan, G.J.; Sun, Z.; Li, Y.; Lan, R.; Chen, L.; Dong, P. Biotransformation of major flavonoid glycosides in herb epimedii by fungus *Cunninghamella blakesleana*. J. Mol. Catal. B Enzym. 2015, 122, 141–146. [CrossRef]
- 152. Jin, S.; Wang, W.; Luo, M.; Mu, F.S.; Li, C.Y.; Fu, Y.J.; Zu, Y.G.; Feng, C. Enhanced extraction genistein from pigeon pea [*Cajanus cajan* (L.) Millsp.] roots with the biotransformation of immobilized edible *Aspergillus oryzae* and *Monacus anka* and antioxidant activity evaluation. *Process Biochem.* **2013**, *48*, 1285–1292. [CrossRef]
- Huynh, T.N.; Smagghe, G.; Gonzales, G.B.; Camp, J.V.; Raes, K. Extraction and bioconversion of kaempferol metabolites from cauliflower outer leaves through fungal fermentation. *Biochem. Eng. J.* 2016, 116, 27–33. [CrossRef]
- 154. Madeira, J.V., Jr.; Mayumi-Nakajima, V.; Alves-Macedo, J.; Alves-Macedo, G. Rich bioactive phenolic extract production by microbial transformation of Brazilian Citrus residues. *Chem. Eng. Res. Des.* **2014**, *92*, 1802–1810. [CrossRef]
- 155. Stahlhut, S.G.; Siedler, S.; Malla, S.; Harrison, S.J.; Maury, J.; Neves, A.R.; Forster, J. Assembly of a novel biosynthetic pathway for production of the plant flavonoid fisetin in *Escherichia coli*. *Metab. Eng.* 2015, *31*, 84–93. [CrossRef]
- 156. Chiang, C.M.; Ding, H.Y.; Lu, J.Y.; Chang, T.S. Biotransformation of isoflavones daidzein and genistein by recombinant *Pichia pastoris* expressing membrane-anchoring and reductase fusion chimeric CYP105D7. *J. Taiwan Inst. Chem. Eng.* **2016**, *60*, 26–31. [CrossRef]
- 157. Yu, L.; Gao, F.; Yang, L.; Xu, L.; Wang, Z.; Ye, H. Biotransformation of puerarin into puerarin-6⁻⁻O-phosphate by *Bacillus cereus. J. Ind. Microbiol. Biotech.* **2012**, *39*, 299–305. [CrossRef]
- 158. Sordon, A.; Poplonski, J.; Tronina, T.; Huszcza, E. Regioselective O-glycosylation of flavonoids by fungi Beauveraia bassiana, Absidia coerulea and Absidia glauca. *Bioorg. Chem.* 2019, 93, 102750. [CrossRef] [PubMed]
- 159. Bok, S.H.; Jeong, T.S.; Lee, S.K.; Kim, J.R.; Moon, S.S.; Choi, M.S.; Hyun, B.H.; Lee, C.H.; Choi, Y.K. Flavanone Derivatives and Composition for Preventing or Treating Blood Lipid Level-Related Diseases Comprising Same. U.S. Patent 6455577, 24 September 2002.
- 160. Hidalgo, A.; Bornscheuer, U.T. Directed evolution of lipases and esterases for organic synthesis. In *Biocatalysis in the Pharmaceutical and Biotechnology Industries*; Patel, R.M., Ed.; CRC Press: Boca Raton, FL, USA; Taylor and Francis: New York, NY, USA, 2006; pp. 159–175.
- 161. Tian, L.; Dixon, R.A. Engineering isoflavone metabolism with an artificial bifunctional enzyme. *Planta* **2006**, 224, 496–507. [CrossRef] [PubMed]

- 162. Kurisawa, M.; Chung, J.E.; Uyama, H.; Kobayashi, S. Enzymatic synthesis and antioxidant properties of poly(rutin). *Biomacromolecules* 2003, *4*, 1394–1399. [CrossRef] [PubMed]
- 163. Mandalari, G.; Bennett, R.N.; Kirby, A.R.; Lo Curto, R.B.; Bisignano, G.; Waldron, K.W.; Faulds, C.B. Enzymatic hydrolysis of flavonoids and pectic oligosaccharides from bergamot (*Citrus bergamia* Risso) peel. J. Agric. Food Chem. 2006, 54, 8307–8313. [CrossRef] [PubMed]
- 164. Jiao, J.; Gai, Q.Y.; Wang, W.; Zang, Y.P.; Niu, L.L.; Fu, Y.J.; Wang, X. Remarjable enhancement of flavonoid production in a co-cultivation system of *Isatis tinctorial* L. hairy root cultures and immobilized *Aspergillus niger. Ind. Crop. Prod.* 2018, 112, 252–261. [CrossRef]
- 165. Ralston, L.; Subramanian, S.; Matsuno, M.; Yu, O. Partial reconstruction of flavonoid and isoflavonoid biosynthesis in yeast using soybean type I and type II chalcone isomerases. *Plant Physiol.* 2005, 137, 1375–1388. [CrossRef]
- 166. Lindahl, S.; Ekman, A.; Khan, S.; Wennerberg, C.; Börjesson, P.; Sjöberg, P.J.; Karlsson, E.N.; Turner, C. Exploring the possibility of using a thermostable mutant of b-glucosidase for rapid hydrolysis of quercetin glucosides in hot water. *Green Chem.* 2010, *12*, 159–168. [CrossRef]
- Ihara, N.; Kurisawa, M.; Chung, J.E.; Uyama, H.; Kobayashi, S. Enzymatic synthesis of a catechin conjugate of polyhedral oligomeric silsesquioxane and evaluation of its antioxidant activity. *Appl. Microbiol. Biotech.* 2005, *66*, 430–433. [CrossRef]
- Yaipakdee, P.; Robertson, L.W. Enzymatic halogenation of flavanones and flavones. *Phytochemistry* 2001, 57, 341–347. [CrossRef]
- 169. Danieli, B.; Luisetti, M.; Riva, S.; Bertinotti, A.; Ragg, E.; Scaglioni, L.; Bombardelli, E. Regioselective enzyme-mediated acylation of polyhydroxy natural compounds. A remarkable, highly efficient preparation of 6'-acetyl and 6'-O-carboxyacetyl ginsenoside Rg1. *J. Org. Chem.* **1995**, *60*, 3637–3642. [CrossRef]
- 170. Chung, J.E.; Kurisawa, M.; Uyama, H.; Kobayashi, S. Enzymatic synthesis and antioxidant property of gelatin-catechin conjugates. *Biotechnol. Lett.* **2003**, *25*, 1993–1997. [CrossRef] [PubMed]
- 171. Xiao, Y.M.; Wu, Q.; Wu, W.B.; Zhang, Q.Y.; Lin, X.F. Controllable regioselective acylation of rutin catalyzed by enzymes in non-aqueous solvents. *Biotechnol. Lett.* **2005**, *27*, 1591–1595. [CrossRef] [PubMed]
- Vitali, A.; Giardina, B.; Delle Monache, G.; Rocca, F.; Silvestrini, A.; Tafi, A.; Botta, B. Chalcone dimethylallyltransferase from *Morus nigra* cell cultures. Substrate specificity studies. *FEBS Lett.* 2004, 557, 33–38. [CrossRef]
- 173. Moon, Y.H.; Lee, J.H.; Ahn, J.S.; Nam, S.H.; Oh, D.K.; Park, D.H.; Chung, H.J.; Kang, S.; Day, D.F.; Kim, D. Synthesis, structure analyses, and characterization of novel epigallocatechin gallate (EGCG) glycosides using the glucansucrase from *Leuconostoc mesenteroides* B-1299CB. *J. Agric. Food Chem.* 2006, 54, 1230–1237. [CrossRef] [PubMed]
- 174. Nielsen, I.L.F.; Chee, W.S.S.; Poulsen, L.; Offord-Cavin, E.; Rasmussen, S.E.; Frederiksen, H.; Enslen, M.; Barron, D.; Horcajada, M.N.; Williamson, G. Bioavailability is improved by enzymatic modification of the citrus flavonoid hesperidin in humans: A randomized, double-blind, crossover trial. *J. Nutr.* 2006, 136, 404–408. [CrossRef] [PubMed]
- 175. Ozaki, T.; Mishima, S.; Nishiyama, M.; Kuzuyama, T. NovQ is a prenyltransferase capable of catalyzing the addition of a dimethylallyl group to both phenylpropanoids and flavonoids. *J. Antibiot.* **2009**, *62*, 385–392. [CrossRef]
- Kobayashi, R.; Itou, T.; Hanaya, K.; Shoji, M.; Hada, N.; Sugai, T. Chemo-enzymatic transformation of naturally abundant naringin to luteolin, a flavonoid with various biological effects. *J. Mol. Catal. B Enzym.* 2013, 92, 14–18. [CrossRef]
- 177. Adamczak, M.; Krishna, S.H. Strategies for improving enzymes for efficient biocatalysis. *Food Technol. Biotechnol.* **2004**, *42*, 251–264.
- 178. Enaud, E.; Humeau, C.; Piffaut, B.; Girardin, M. Enzymatic synthesis of new aromatic esters of phloridzin. *J. Mol. Catal. B Enzym.* **2004**, 27, 1–6. [CrossRef]
- 179. Chebil, L.; Humeau, C.; Falcimaigne, A.; Engasser, J.M.; Ghoul, M. Enzymatic acylation of flavonoids. *Process Biochem.* 2006, 41, 2237–2251. [CrossRef]
- Das, S.; Rosazza, J.P.N. Microbial and enzymatic transformations of flavonoids. J. Nat. Prod. 2006, 69, 499–508.
 [CrossRef] [PubMed]
- Viskupicova, J.; Ondrejovic, M.; Maliar, T. Enzyme-mediated preparation of flavonoid esters and their applications. *Biochemistry* 2012, 10, 263–286.

- Peterson, J.J.; Dwyer, J.T.; Jacques, P.F.; McCullough, M.L. Associations between flavonoids and cardiovascular disease incidence or mortality in European and US populations. *Nutr. Rev.* 2012, 70, 491–508. [CrossRef] [PubMed]
- Koirala, N.; Thuan, N.H.; Ghimire, G.P.; Thang, D.V.; Sohng, J.K. Methylation of flavonoids: Chemical structures, bioactivities, progress and perspectives for biotechnological production. *Enzym. Microb. Technol.* 2016, *86*, 103–116. [CrossRef] [PubMed]
- 184. Chen, G.L.; Fan, M.X.; Wu, J.L.; Li, N.; Guo, M.Q. Antioxidant and anti-inflammatory properties of flavonoids from lotus plumule. *Food Chem.* **2019**, 277, 706–712. [CrossRef] [PubMed]
- Treml, J.; Smejkal, K. Flavonoids as potent scavengers of hydroxyl radicals. *Compr. Rev. Food Sci. Food Saf.* 2016, 15, 720–738. [CrossRef]
- 186. Agati, G.; Tattini, M. Multiple functional roles of flavonoids in photoprotection. *New Phytol.* **2010**, *186*, 786–793. [CrossRef]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).