

Article

GA₃ Treatment Delays the Deterioration of ‘Shixia’ Longan during the On-Tree Preservation and Room-Temperature Storage and Up-Regulates Antioxidants

Tao Luo ^{1,2,3,†}, Xiaolan Lin ^{1,2,†}, Tingting Lai ^{1,2}, Libing Long ^{1,2}, Ziyang Lai ^{1,2}, Xinxin Du ^{1,2}, Xiaomeng Guo ^{1,2}, Liang Shuai ⁴, Dongmei Han ⁵ and Zhenxian Wu ^{1,2,3,*}

¹ College of Horticulture, South China Agricultural University, Guangzhou 510642, China; luotao0502@scau.edu.cn (T.L.); linxl4595@stu.scau.edu.cn (X.L.); ltt7059@stu.scau.edu.cn (T.L.); longlibing@stu.scau.edu.cn (L.L.); laiziying@stu.scau.edu.cn (Z.L.); xinxindu@stu.scau.edu.cn (X.D.); 31000462@scau.edu.cn (X.G.)

² Guangdong Provincial Key Laboratory of Postharvest Science of Fruits and Vegetables, Engineering Research Center of Southern Horticultural Products Preservation, Ministry of Education, Guangzhou 510642, China

³ Key Laboratory of Biology and Genetic Improvement of Horticultural Crops (South China), Ministry of Agriculture and Rural Affairs, College of Horticulture, South China Agricultural University, Guangzhou 510642, China

⁴ College of Chemistry and Food Science, Nanchang Normal University, Nanchang 330032, China; shuailiang1212@hzxy.edu.cn

⁵ Institute of Fruit Tree Research, Guangdong Academy of Agricultural Sciences, Key Laboratory of South Subtropical Fruit Biology and Genetic Resource Utilization, Ministry of Agriculture, Guangzhou 510640, China; handongmei@gdaas.cn

* Correspondence: zhenxwu@scau.edu.cn.

† These authors contributed equally to this work.




Figure S1. The TSS content of the GA₃ treated and control longan pulp in the year 2020 and 2021.




Figure S2. Statistics of metabolites in each class according to the fold change by the GA₃ vs. control longan pulp.




Figure S3. The fold changes of the metabolites from the important pathways. CK: Control; GA₃: 50 mg L⁻¹ GA₃.