

Table S3. Regression equations and the corresponding reliability of four flavonoid standards

<i>Compound</i>	<i>Regression equation</i>	<i>R</i> ²	<i>Linear ranges</i> ($\mu\text{g/mL}$)	<i>LOD</i> ($\mu\text{g/mL}$)	<i>LQD</i> ($\mu\text{g/mL}$)	<i>Precisio</i> (<i>n</i> =6)	<i>Repeatability</i> (<i>n</i> =6)	<i>Accuracy</i> (<i>n</i> =6)	
						<i>RSD</i> (%)	<i>RSD</i> (%)	<i>RSD</i> (%)	<i>Recovery</i> (%)
eriodictyol	Y=32264X-8.1381	0.9998	2.5~250	0.3333	1.0000	1.18%	0.69%	3.22%	94.8%
eriodictyol-7-O-glucoside	Y=23680X+29.012	0.9991	4~400	0.0833	0.3333	1.09%	1.24%	2.24%	96.04%
luteolin	Y=16091X-21.205	0.9995	1.96~196	0.2500	0.8000	1.35%	1.24%	2.13%	98.56%
luteolin-7-O-glucoside	Y=20785X+2.6562	0.9994	2.2~220	0.0270	0.0813	1.70%	1.26%	2.72%	100.58%