

***Chaenomeles sinensis* extract ameliorates ovalbumin-induced allergic rhinitis by inhibiting the IL-33/ST2 axis and regulating epithelial cell dysfunction**

Juan Jin^{1*}, Yan Jing Fan^{1,2*}, Thi Van Nguyen¹, Zhen Nan Yu¹, Chang Ho Song^{1,3}, So-Young Lee^{4,5}, Hee Soon Shin^{4,5}, Ok Hee Chai^{1,3}

¹ Department of Anatomy, Jeonbuk National University Medical School, Jeonju, 54896, Jeonbuk, Korea

² School of Medicine, Liaocheng University, Liaocheng, Shandong, People's Republic of China.

³ Institute for Medical Sciences, Jeonbuk National University Medical School, Jeonju, 54896, Jeonbuk, Korea

⁴ Department of Food Biotechnology, University of Science and Technology (UST), Daejeon 34113, Korea

⁵ Division of Food Functionality Research, Korea Food Research Institute (KFRI), Wanju 55365, Korea

Correspondence: Ok Hee Chai, Department of Anatomy and Institute for Medical Sciences, Jeonbuk National University Medical School, Jeonju, 54896, Jeonbuk, Korea. Tel: +82-63-2703109, Fax: +82-63-2749880. Email: okchai1004@jbnu.ac.kr.

* These authors contributed equally to this work.




Figure S1. Full-length gels and blots of IL-33 signaling-related proteins in lung tissues (A) Western blot data. Original Western blot picture of (B) IL-33 and (C) β-actin. Membrane C was stripped from membrane B.




Figure S2. Full-length gels and blots of ST2 signaling-related proteins in lung tissues (A) Western blot data. Original Western blot picture of (B) IL-33 and (C) β-actin. Membrane C was stripped from membrane B.